



PERTH MODERN SCHOOL
Exceptional schooling. Exceptional students.
Independent Public School

Course Specialist

Year 11

Student name: _____

Teacher name: _____

Date: 18 Sep 2020

Task type: Response

Time allowed for this task: 45 mins

Number of questions: 6

Materials required: Calculator-Free

Standard items: Pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: Drawing instruments, templates

Marks available: 45 marks

Task weighting: 16 %

Formula sheet provided: Yes

Note: All part questions worth more than 2 marks require working to obtain full marks.

Question 1 (2.2.1, 2.2.2, 2.2.3)**(6 marks)**

If $A = \begin{bmatrix} 1 & 0 \\ -2 & -3 \end{bmatrix}$, O is the 2×2 zero matrix and I is the 2×2 identity matrix, find

a) Matrix B given that $A - B = I$

(1 mark)

$$\begin{aligned} B &= A - I = \begin{bmatrix} 1 & 0 \\ -2 & -3 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ -2 & -4 \end{bmatrix} \checkmark \end{aligned}$$

b) Matrix C given that $2A + C = O$

(1 mark)

$$\begin{aligned} C &= -2A \\ &= \begin{bmatrix} -2 & 0 \\ 4 & 6 \end{bmatrix} \checkmark \end{aligned}$$

c) Matrix D given that $D = B - AD$, where B is from part a)

(4 marks)

$$\begin{aligned} D + AD &= B \\ (I + A)D &= B \checkmark \\ D &= (I + A)^{-1} B \checkmark \\ &= \begin{bmatrix} 2 & 0 \\ -2 & -2 \end{bmatrix}^{-1} \begin{bmatrix} 0 & 0 \\ -2 & -4 \end{bmatrix} \\ &= -\frac{1}{4} \begin{bmatrix} -2 & 0 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ -2 & -4 \end{bmatrix} \checkmark \\ &= -\frac{1}{4} \begin{bmatrix} 0 & 0 \\ -4 & -8 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ 1 & 2 \end{bmatrix} \end{aligned}$$

} either \checkmark

or Let $D = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

$$\begin{aligned} a &= 0 \checkmark \\ b &= 0 \checkmark \quad R/W \\ c &= 1 \checkmark \\ d &= 2 \checkmark \end{aligned}$$

Question 2 (2.2.11)**(8 marks)**

If $A = \begin{bmatrix} 7 & 5 \\ -4 & -3 \end{bmatrix}$ and $B = \begin{bmatrix} 6 & 10 \\ -8 & -14 \end{bmatrix}$

a) Determine AB .

(2 marks)

$$\begin{bmatrix} 42-40 & 70-70 \\ -24+24 & -40+42 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

b) Express A^{-1} in terms of B .

(3 marks)

$$AB = 2I \quad \checkmark$$

$$A^{-1}AB = 2A^{-1}I \quad \checkmark$$

$$B = 2A^{-1}$$

$$A^{-1} = \frac{B}{2} \quad \checkmark$$

c) Solve the system $\begin{cases} 7x + 5y = 1 \\ -4x - 3y = 1 \end{cases}$, clearly showing your use of A^{-1} . (3 marks)

Award full mark if student used $4x + 3y = 1$

$$\begin{bmatrix} 7 & 5 \\ -4 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \checkmark$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 7 & 5 \\ -4 & -3 \end{bmatrix}^{-1} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \checkmark$$

$$= \begin{bmatrix} 3 & 5 \\ -4 & -7 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} 8 \\ -11 \end{bmatrix} \quad \checkmark$$

$$x = 8, \quad y = -11$$


$$\begin{bmatrix} 7 & 5 \\ 4 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \checkmark$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 7 & 5 \\ 4 & 3 \end{bmatrix}^{-1} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \checkmark$$

$$= \frac{1}{21-20} \begin{bmatrix} 3 & -5 \\ -4 & 7 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} -2 \\ 3 \end{bmatrix} \quad \checkmark$$

$$\therefore x = -2, \quad y = 3$$

Question 3 (2.1.4, 2.1.7)**(7 marks)**a) On the axes below, sketch the graph of $y = 5 \sec(x - \pi)$, $0 \leq x \leq 2\pi$.**(3 marks)**

✓ Asymptotes
 ✓ Maximum & Minimum turning points
 ✓ Shape with smooth curves

b) Find the general solution for $\sqrt{3} \cos(x) - \sin(x) = 1$.**(4 marks)**


$$\frac{\sqrt{3}}{2} \cos x - \frac{1}{2} \sin x = \frac{1}{2}$$

$$\cos x \cos \frac{\pi}{6} - \sin x \sin \frac{\pi}{6} = \frac{1}{2}$$

$$\cos \left(x + \frac{\pi}{6} \right) = \frac{1}{2} \quad \checkmark$$

$$\therefore x + \frac{\pi}{6} = \pm \frac{\pi}{3} + 2k\pi, \quad k \in \mathbb{Z} \quad \checkmark$$

$$\therefore x = \frac{\pi}{6} + 2k\pi \quad \checkmark \quad \text{or} \quad -\frac{\pi}{2} + 2k\pi, \quad k \in \mathbb{Z} \quad \checkmark$$



or

$$\sin x - \sqrt{3} \cos x = -1$$

$$\frac{1}{2} \sin x - \frac{\sqrt{3}}{2} \cos x = -\frac{1}{2}$$

$$\sin x \cos \frac{\pi}{3} - \cos x \sin \frac{\pi}{3} = -\frac{1}{2}$$

$$\therefore \sin \left(x - \frac{\pi}{3} \right) = -\frac{1}{2} \quad \checkmark$$

$$\therefore x - \frac{\pi}{3} = -\frac{\pi}{6} + 2k\pi \quad \text{or} \quad -\frac{5\pi}{6} + 2k\pi, \quad k \in \mathbb{Z} \quad \checkmark$$

$$\therefore x = \frac{\pi}{6} + 2k\pi \quad \checkmark \quad \text{or} \quad -\frac{\pi}{2} + 2k\pi, \quad k \in \mathbb{Z} \quad \checkmark$$

Question 4 (2.2.3, 2.1.3)**(6 marks)**

Let $A = \begin{bmatrix} \cos(\alpha) & \sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{bmatrix}$ and $B = \begin{bmatrix} \cos(\beta) \\ \sin(\beta) \end{bmatrix}$, such that $AB = \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix}$.

Find two different sets of possible values for α and β given that $0 \leq \alpha < \frac{\pi}{2}$ and $0 \leq \beta < \frac{\pi}{2}$

$$AB = \begin{bmatrix} \cos\alpha\cos\beta + \sin\alpha\sin\beta \\ \sin\alpha\cos\beta + \cos\alpha\sin\beta \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix}$$

$$\therefore \cos(\alpha - \beta) = \frac{1}{2}, \quad -\frac{\pi}{2} < \alpha - \beta < \frac{\pi}{2}$$

$$\sin(\alpha + \beta) = \frac{\sqrt{3}}{2}, \quad 0 \leq \alpha + \beta < \pi$$

$$\therefore \begin{cases} \alpha - \beta = \frac{\pi}{3} \\ \alpha + \beta = \frac{\pi}{3} \end{cases} \quad \text{or} \quad \begin{cases} \alpha - \beta = \frac{\pi}{3} \\ \alpha + \beta = \frac{2\pi}{3} \end{cases} \quad \text{or} \quad \begin{cases} \alpha - \beta = -\frac{\pi}{3} \\ \alpha + \beta = \frac{\pi}{3} \end{cases} \quad \text{or} \quad \begin{cases} \alpha - \beta = -\frac{\pi}{3} \\ \alpha + \beta = \frac{2\pi}{3} \end{cases}$$

$$\begin{cases} \alpha = \frac{\pi}{3} \\ \beta = 0 \end{cases} \quad \text{or} \quad \begin{cases} \alpha = \frac{\pi}{2} \\ \beta = \frac{\pi}{6} \end{cases} \quad \text{(exclude)} \quad \text{or} \quad \begin{cases} \alpha = 0 \\ \beta = \frac{\pi}{3} \end{cases} \quad \text{or} \quad \begin{cases} \alpha = \frac{\pi}{6} \\ \beta = \frac{\pi}{2} \end{cases} \quad \text{(exclude)}$$

* Award 1 mark if student gives $\alpha - \beta = \frac{\pi}{3}$ or $\alpha + \beta = \frac{\pi}{3}$ only

Question 5 (2.1.3, 2.1.5)**(6 marks)**

a) First show that

$$\tan\left(\theta - \frac{\pi}{4}\right) = \frac{\sin \theta - \cos \theta}{\cos \theta + \sin \theta}$$

(2 marks)

$$\begin{aligned}
 \text{LHS} &= \frac{\tan \theta - 1}{1 + \tan \theta} \quad \checkmark \\
 &= \frac{\frac{\sin \theta}{\cos \theta} - 1}{1 + \frac{\sin \theta}{\cos \theta}} \\
 &= \frac{\frac{\sin \theta - \cos \theta}{\cos \theta}}{\frac{\cos \theta + \sin \theta}{\cos \theta}} \quad \left. \begin{array}{l} \text{either } \checkmark \text{ or} \end{array} \right\} = \frac{\frac{\sin \theta}{\cos \theta} - \frac{\cos \theta}{\cos \theta}}{\frac{\cos \theta}{\cos \theta} + \frac{\sin \theta}{\cos \theta}} \quad \text{or} = \frac{\frac{\sin \theta}{\cos \theta} - 1}{1 + \frac{\sin \theta}{\cos \theta}} \times \frac{\cos \theta}{\cos \theta} \\
 &= \frac{\sin \theta - \cos \theta}{\cos \theta} \times \frac{\cos \theta}{\cos \theta + \sin \theta} \\
 &= \frac{\sin \theta - \cos \theta}{\cos \theta + \sin \theta}
 \end{aligned}$$

b) Hence (or otherwise) prove the following identity:

$$\tan\left(\theta - \frac{\pi}{4}\right) = \frac{\sin(2\theta) - 1}{1 - 2 \sin^2 \theta}$$

(4 marks)

$$\begin{aligned}
 \text{LHS} &= \frac{\sin \theta - \cos \theta}{\cos \theta + \sin \theta} \times \frac{\cos \theta - \sin \theta}{\cos \theta - \sin \theta} \quad \checkmark \\
 &= \frac{\sin \theta \cos \theta - \sin^2 \theta - \cos^2 \theta + \sin \theta \cos \theta}{\cos^2 \theta - \sin^2 \theta} \\
 &= \frac{2 \sin \theta \cos \theta - (\sin^2 \theta + \cos^2 \theta)}{\cos 2\theta} \quad \begin{array}{l} \checkmark \text{ Uses double angle identity for sine} \\ \checkmark \text{ Uses Pythagorean identity} \end{array} \\
 &= \frac{\sin 2\theta - 1}{1 - 2 \sin^2 \theta} \quad \checkmark \text{ Uses double angle identities for cosine}
 \end{aligned}$$

Question 6 (2.2.4, 2.2.5, 2.2.6, 2.2.7, 2.2.8, 2.2.9, 2.2.10)**(12 marks)**

a) Determine the matrices that produce each of the transformations described below:

- i. a rotation clockwise about the origin by
- 90°
- (1 mark)

$$\begin{bmatrix} \cos(-90^\circ) & -\sin(-90^\circ) \\ \sin(-90^\circ) & \cos(-90^\circ) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \quad \checkmark$$

- ii. a dilation parallel to the y-axis by a scale factor of 2 (1 mark)

$$\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \quad \checkmark$$


- iii. a reflection in the line
- $y = x$
- (1 mark)

$$\begin{bmatrix} \cos 90^\circ & \sin 90^\circ \\ \sin 90^\circ & -\cos 90^\circ \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \checkmark$$

- b) Show how to obtain the single transformation matrix
- T
- , given that
- T
- is the result of applying the transformations given in part a) in the order listed [i.e. a rotation clockwise about the origin by
- 90°
- , followed by a dilation parallel to the y-axis by a scale factor of 2, then a reflection in the line
- $y = x$
-]. (2 marks)

$$\begin{aligned} T &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \times \left(\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \times \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right) \quad \checkmark \\ &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \times \begin{bmatrix} 0 & 1 \\ -2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} -2 & 0 \\ 0 & 1 \end{bmatrix} \quad \checkmark \end{aligned}$$

- c) $\triangle ABC$ is first translated left by 1 unit and down by 2 units, then transformed by the transformation matrix T in part b). The final image $\triangle A'B'C'$ is shown below:



- i. Determine the coordinates of points A, B and C. (5 marks)

$$\begin{aligned} & \begin{bmatrix} -2 & 0 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} -2 & -4 & -4 \\ 0 & 0 & -2 \end{bmatrix} \checkmark \\ &= -\frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} -2 & -4 & -4 \\ 0 & 0 & -2 \end{bmatrix} \\ &= -\frac{1}{2} \begin{bmatrix} -2 & -4 & -4 \\ 0 & 0 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 2 & 2 \\ 0 & 0 & -2 \end{bmatrix} \checkmark \end{aligned}$$

$$\begin{aligned} \therefore A & (2, 2) \checkmark \\ B & (3, 2) \checkmark \\ C & (3, 0) \checkmark \end{aligned}$$

Allow follow through here

- ii. $\triangle A'B'C'$ is now transformed by a matrix

$$M = \begin{bmatrix} 2\sqrt{5} & \sqrt{5} \\ 3 & -1 \end{bmatrix}$$

Determine the exact area of the image of $\triangle A'B'C'$ under this transformation.

(2 marks)

$$|\det(M)| = |-2\sqrt{5} - 3\sqrt{5}| = 5\sqrt{5} \checkmark$$

$$\begin{aligned} \therefore A &= 5\sqrt{5} \times \left(\frac{1}{2} \times 2 \times 2\right) \\ &= 10\sqrt{5} \text{ units}^2 \checkmark \end{aligned}$$

Additional working space

Question number: _____