



MATHEMATICS SPECIALIST ATAR COURSE

FORMULA SHEET

2017

Copyright

© School Curriculum and Standards Authority, 2016

This document – apart from any third party copyright material contained in it – may be freely copied, or communicated on an intranet, for non-commercial purposes in educational institutions, provided that it is not changed and that the School Curriculum and Standards Authority is acknowledged as the copyright owner, and that the Authority's moral rights are not infringed.

Copying or communication for any other purpose can be done only within the terms of the *Copyright Act 1968* or with prior written permission of the School Curriculum and Standards Authority. Copying or communication of any third party copyright material can be done only within the terms of the *Copyright Act 1968* or with permission of the copyright owners.

Any content in this document that has been derived from the Australian Curriculum may be used under the terms of the Creative Commons [Attribution 4.0 International \(CC BY\)](#) licence.

This document is valid for teaching and examining until 31 December 2017.

Index

Differentiation and integration **3**

Applications of calculus
Functions
Statistical inference **4**

Mensuration
Vectors in 3D **5**

Complex numbers **6**

Trigonometry **7**

Differentiation and integration

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{x^{n+1}}{n+1} + c, \quad n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$
$\frac{d}{dx}(\ln x) = \frac{1}{x}$	$\int \frac{1}{x} dx = \ln x + c$
$\frac{d}{dx}(\ln f(x)) = \frac{f'(x)}{f(x)}$	$\int \frac{f'(x)}{f(x)} dx = \ln f(x) + c$
$\frac{d}{dx}(\sin f(x)) = f'(x) \cos(f(x))$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos f(x)) = -f'(x) \sin(f(x))$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan f(x)) = f'(x) \sec^2(f(x)) = \frac{f'(x)}{\cos^2 f(x)}$	$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$
Product rule	<div> <div> <p>If $y = uv$</p> <p>then</p> $\frac{d}{dx}(uv) = u \frac{dv}{dx} + \frac{du}{dx} v$ </div> <div> <p>If $y = f(x) g(x)$</p> <p>then</p> $y' = f'(x) g(x) + f(x) g'(x)$ </div> </div>
Quotient rule	<div> <div> <p>If $y = \frac{u}{v}$</p> <p>then</p> $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$ </div> <div> <p>If $y = \frac{f(x)}{g(x)}$</p> <p>then</p> $y' = \frac{f'(x) g(x) - f(x) g'(x)}{(g(x))^2}$ </div> </div>
Chain rule	<div> <div> <p>If $y = f(u)$ and $u = g(x)$</p> <p>then</p> $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$ </div> <div> <p>If $y = f(g(x))$</p> <p>then</p> $y' = f'(g(x)) g'(x)$ </div> </div>
Fundamental theorem	<div> $\frac{d}{dx}\left(\int_a^x f(t) dt\right) = f(x)$ </div> <div>and</div> <div> $\int_a^b f'(x) dx = f(b) - f(a)$ </div>

Applications of calculus

Growth and decay	
Exponential equation	$\frac{dP}{dt} = kP \Leftrightarrow P = P_0 e^{kt}$
Logistic equation	$\frac{dP}{dt} = rP(k - P) \Leftrightarrow P = \frac{kP_0}{P_0 + (k - P_0)e^{-rkt}}$
Volumes of solids of revolution	
About the x -axis	$V = \pi \int_a^b [f(x)]^2 dx$
About the y -axis	$V = \pi \int_c^d [f(y)]^2 dy$
Simple harmonic motion	
<p>If $\frac{d^2x}{dt^2} = -k^2x$ then $x = A \sin(kt + \alpha)$ or $x = A \cos(kt + \beta)$</p> <p>where A is the amplitude, α and β are phase angles, v is the velocity and x is the displacement</p>	
$v^2 = k^2(A^2 - x^2)$ Period: $T = \frac{2\pi}{k}$ Frequency: $f = \frac{1}{T}$	
Incremental formula	$\delta y \approx \frac{dy}{dx} \times \delta x$
Acceleration	$\frac{dv}{dt}$ or $v \frac{dv}{dx}$ or $\frac{d}{dx} \left(\frac{1}{2} v^2 \right)$

Functions

Quadratic function	If $f(x) = ax^2 + bx + c$ and $f(x) = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Absolute value function	$ x = \begin{cases} x, & \text{for } x \geq 0 \\ -x, & \text{for } x < 0 \end{cases}$

Statistical inference

Confidence interval for the mean of the population:	$\bar{X} - z \frac{s}{\sqrt{n}} \leq \mu \leq \bar{X} + z \frac{s}{\sqrt{n}}$
Sample size:	$n = \left(\frac{z \times s}{d} \right)^2$

Mensuration

Parallelogram	$A = bh$	
Triangle	$A = \frac{1}{2}bh$ or $A = \frac{1}{2}ab \sin C$	
Trapezium	$A = \frac{1}{2}(a + b)h$	
Circle	$A = \pi r^2$ and $C = 2\pi r = \pi d$	
Prism	$V = Ah$, where A is the area of the cross section	
Pyramid	$V = \frac{1}{3}Ah$, where A is the area of the cross section	
Cylinder	$V = \pi r^2 h$	$S = 2\pi r h + 2\pi r^2$
Cone	$V = \frac{1}{3}\pi r^2 h$	$S = \pi r s + \pi r^2$, where s is the slant height
Sphere	$V = \frac{4}{3}\pi r^3$	$S = 4\pi r^2$

Vectors in 3D

Magnitude	$ (a_1, a_2, a_3) = \sqrt{a_1^2 + a_2^2 + a_3^2}$
Dot product	$\mathbf{a} \cdot \mathbf{b} = \mathbf{a} \mathbf{b} \cos\theta = a_1b_1 + a_2b_2 + a_3b_3$
Cross product	$\mathbf{a} \times \mathbf{b} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \times \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{pmatrix}$
Equation of a line	One point and direction $\mathbf{r} = \mathbf{a} + \lambda\mathbf{u}$
	Two points A and B $\mathbf{r} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$
Equation of a plane	$\mathbf{r} = \mathbf{a} + \lambda\mathbf{u}_1 + \mu\mathbf{u}_2$ or $\mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$
Equation of a sphere	$ \mathbf{r} - \mathbf{d} = r$ or $(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2$
Cartesian equation of a line	$\frac{x - a_1}{u_1} = \frac{y - a_2}{u_2} = \frac{z - a_3}{u_3}$
Cartesian equation of a plane	$ax + by + cz = d$
Parametric equation of a line	$x = a_1 + \lambda u_1 \dots \dots (1)$ $y = a_2 + \lambda u_2 \dots \dots (2)$ $z = a_3 + \lambda u_3 \dots \dots (3)$

Complex numbers

Cartesian form	
$z = a + bi$	$\bar{z} = a - bi$
$\text{Mod}(z) = z = \sqrt{a^2 + b^2} = r$	$\text{Arg}(z) = \theta, \quad \tan \theta = \frac{b}{a}, \quad -\pi < \theta \leq \pi$
$ z_1 z_2 = z_1 z_2 $	$\left \frac{z_1}{z_2} \right = \frac{ z_1 }{ z_2 }$
$\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$	$\arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2)$
$z \bar{z} = z ^2$	$z^{-1} = \frac{1}{z} = \frac{\bar{z}}{ z ^2}$
$\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$	$\overline{z_1 z_2} = \bar{z}_1 \bar{z}_2$
Polar form	
$z = a + bi = r(\cos \theta + i \sin \theta) = r \text{cis } \theta$	$\bar{z} = r \text{cis } (-\theta)$
$z_1 z_2 = r_1 r_2 \text{cis } (\theta_1 + \theta_2)$	$\frac{z_1}{z_2} = \frac{r_1}{r_2} \text{cis } (\theta_1 - \theta_2)$
$\text{cis}(\theta_1 + \theta_2) = \text{cis } \theta_1 \text{cis } \theta_2$	$\text{cis}(-\theta) = \frac{1}{\text{cis } \theta}$
De Moivre's theorem	
$z^n = z ^n \text{cis } (n\theta)$	$(\text{cis } \theta)^n = \cos n\theta + i \sin n\theta$
$z^{\frac{1}{q}} = r^{\frac{1}{q}} \left(\cos \frac{\theta + 2\pi k}{q} + i \sin \frac{\theta + 2\pi k}{q} \right), \quad \text{for } k \text{ an integer}$	

Trigonometry

$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$	Length of arc = $r\theta$
$a^2 = b^2 + c^2 - 2bc \cos A$	Area of segment = $\frac{1}{2} r^2 (\theta - \sin \theta)$
$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$	Area of sector = $\frac{1}{2} r^2 \theta$
Identities	
$\cos^2 x + \sin^2 x = 1$	$1 + \tan^2 x = \sec^2 x$
$\cos (x \pm y) = \cos x \cos y \mp \sin x \sin y$	$\cos 2x = \cos^2 x - \sin^2 x$ $= 2 \cos^2 x - 1$ $= 1 - 2 \sin^2 x$
$\sin (x \pm y) = \sin x \cos y \pm \cos x \sin y$	$\sin 2x = 2 \sin x \cos x$
$\tan (x \pm y) = \frac{\tan x \pm \tan y}{1 \mp \tan x \tan y}$	$\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$
$\cos A \cos B = \frac{1}{2} (\cos(A - B) + \cos(A + B))$	$\sin A \cos B = \frac{1}{2} (\sin(A + B) + \sin(A - B))$
$\sin A \sin B = \frac{1}{2} (\cos(A - B) - \cos(A + B))$	$\cos A \sin B = \frac{1}{2} (\sin(A + B) - \sin(A - B))$

Note: Any additional formulas identified by the examination panel as necessary will be included in the body of the particular question.

