

Computational Fluid Dynamics

HW2

Almog Dobrescu

ID 214254252

March 12, 2025

Contents

1	Problem Definition	3
1.1	Governing Equations	3
1.2	Physical Domain	3
1.3	Initial Conditions	3
1.4	Boundary Conditions	4
2	Normalizing The Navier-Stokes Equations	4
3	The Computational Domain	5
3.1	Discretization	5
3.2	Boundary Conditions	6
4	The Numerical Schemes	7
4.1	Jacobian Matrices of The Navier-Stokes Equations	7
4.2	Linearization In Time	7
4.2.1	\tilde{E}_i^{n+1} Estimation	7
4.2.2	\tilde{V}_{1i}^{n+1} Estimation	8
4.3	First Order Approximate Riemann Roe Method	8
4.4	First Order Steger-Warming – Explicit	8
4.4.1	Finite Volume Formulation	9
4.4.2	Calculating $\tilde{V}_{1,i+\frac{1}{2}}$	10
4.5	First Order Steger-Warming – Implicit	10

List of Figures

1	Initial conditions	3
---	------------------------------	---



Nomenclature

Δx	size of each cell in the domain
$\Delta \tilde{t}$	normalized step size in time
$\Delta \tilde{x}$	normalized step size in space
γ	ratio of specific heats
κ	coefficient of thermal conductivity
μ	coefficient of viscosity
ρ	fluid density
c_p	constant specific heat capacity for a constant pressure
c_v	constant specific heat capacity for a constant volume
E	inviscid convective vector
e	total energy
E_ν	viscous convective vector
L	characteristic length
p	pressure
Q	conservation state space
R	gas constant
T	temperature
t	time
u	fluid velocity
x	spatial coordinate
x_F	x coordinate of the end of the domain
x_i	x coordinate of the i-th cell

Diagonal

Φ	main diagonal
Ψ	upper off-diagonal
Θ	lower off-diagonal

Far-Away Properties

κ_∞	coefficient of thermal conductivity far away
μ_∞	coefficient of viscosity far away



ρ_∞	density far away
a_∞	speed of sound far away
M_∞	mach number far away
p_∞	pressure far away
T_∞	temperature far away

Matrices

$\tilde{\Lambda}$	normalized eigenvalues matrix
\tilde{A}	normalized jacobian matrix of E w.r.t. Q
\tilde{P}	normalized jacobian matrix of E_ν w.r.t. Q
\tilde{R}	normalized jacobian matrix of E_ν w.r.t. Q_x
\tilde{T}	normalized eigenvectors matrix

Dimensionless Numbers

Pr_∞	Prandtl number far away
$Re_{L\infty}$	Reynolds number with respect to L far away



1 Problem Definition

1.1 Governing Equations

Consider the one-dimensional Navier-Stokes Equations:

$$\frac{\partial Q}{\partial t} + \frac{\partial E}{\partial x} = \frac{\partial E_\nu}{\partial x} \quad (1)$$

Where:

$$Q = \begin{pmatrix} \rho \\ \rho u \\ e \end{pmatrix}, \quad E = \begin{pmatrix} \rho u \\ p + \rho u^2 \\ (e + p)u \end{pmatrix}, \quad E_\nu = \begin{pmatrix} 0 \\ \tau_{xx} \\ u\tau_{xx} - q_x \end{pmatrix} = \begin{pmatrix} 0 \\ \frac{4}{3}\mu \frac{\partial u}{\partial x} \\ \frac{4}{3}\mu u \frac{\partial u}{\partial x} + \kappa \frac{\partial T}{\partial x} \end{pmatrix} \quad (2)$$

$$p = (\gamma - 1) \left(e - \frac{1}{2} \rho u^2 \right), \quad T = \frac{p}{\rho R},$$

$$\mu = 1.458 \cdot 10^{-6} \frac{T^{\frac{3}{2}}}{T + 110.4}, \quad \kappa = 2.495 \cdot 10^{-3} \frac{T^{\frac{3}{2}}}{T + 194}$$

$$R = c_p - c_v, \quad \gamma = \frac{c_p}{c_v}$$

The constants are:

- $\gamma = 1.4$ for air under standard atmospheric conditions
- $R = 287.0$ for air

1.2 Physical Domain

The physical domain is a tube extended between $x = 0.2$ and $x = 1.0$. At both ends there are impermeable walls.

1.3 Initial Conditions

The initial conditions are shown in Fig.1:




Figure 1: Initial conditions



1.4 Boundary Conditions

On each side of the tube there is an adiabatic, solid wall boundary conditions.

$$u_{(x=0.2)} = u_{(x=1.0)} = 0 \quad \left\| \quad \frac{\partial p}{\partial x} \Big|_{x=0.2} = \frac{\partial p}{\partial x} \Big|_{x=1.0} = 0 \quad \right\| \quad \frac{\partial T}{\partial x} \Big|_{x=0.2} = \frac{\partial T}{\partial x} \Big|_{x=1.0} = 0$$

2 Normalizing The Navier-Stokes Equations

Since the initial conditions are normalized, there is a need to normalize the N-S equations. We will use the following normalizations:

$$\rho = \rho_\infty \tilde{\rho}, \quad u = a_\infty \tilde{u}, \quad p = \gamma p_\infty \tilde{p}, \quad T = \gamma T_\infty \tilde{T}, \quad x = L \tilde{x}, \quad t = \frac{L}{a_\infty} \tilde{t}, \quad \mu = \mu_\infty \tilde{\mu}, \quad \kappa = \kappa_\infty \tilde{\kappa} \quad (3)$$

The normalization of the temperature was chosen to cancel out the γ in the normalization of the pressure:

$$\begin{aligned} p &= \rho R T \\ \gamma p_\infty \tilde{p} &= \rho_\infty \tilde{\rho} R \gamma T_\infty \tilde{T} \\ \tilde{p} &= \tilde{\rho} \tilde{T} \end{aligned} \quad (4)$$

The pressure normalization can be written also as:

$$p = \gamma p_\infty \tilde{p} = \gamma \rho_\infty R T_\infty \tilde{p} = \rho_\infty a_\infty^2 \tilde{p} \quad (5)$$

From equations 2 and 5 we can derive the normalization for the energy:

$$\begin{aligned} e &= \frac{p}{\gamma - 1} + \frac{1}{2} \rho u^2 \\ e &= \frac{\rho_\infty a_\infty^2 \tilde{p}}{\gamma - 1} + \frac{1}{2} \rho_\infty \tilde{\rho} a_\infty^2 \tilde{u}^2 \\ e &= \rho_\infty a_\infty^2 \left(\frac{\tilde{p}}{\gamma - 1} + \frac{1}{2} \tilde{\rho} \tilde{u}^2 \right) \\ e &= \rho_\infty a_\infty^2 \tilde{e} \end{aligned} \quad (6)$$

After substituting the normalizations in the N-S equations we get:

$$\frac{\partial}{\partial \frac{L}{a_\infty} \tilde{t}} \begin{pmatrix} \rho_\infty \tilde{\rho} \\ \rho_\infty a_\infty \tilde{\rho} \tilde{u} \\ \rho_\infty a_\infty^2 \tilde{e} \end{pmatrix} + \frac{\partial}{\partial L \tilde{x}} \begin{pmatrix} \rho_\infty a_\infty \tilde{\rho} \tilde{u} \\ \rho_\infty a_\infty^2 \tilde{p} + \rho_\infty a_\infty^2 \tilde{\rho} \tilde{u}^2 \\ \rho_\infty a_\infty^3 (\tilde{e} + \tilde{p}) \tilde{u} \end{pmatrix} = \frac{\partial}{\partial L \tilde{x}} \begin{pmatrix} 0 \\ \frac{4}{3} \mu_\infty a_\infty \tilde{\mu} \frac{\partial \tilde{u}}{\partial L \tilde{x}} \\ \frac{4}{3} \mu_\infty a_\infty^2 \tilde{\mu} \tilde{u} \frac{\partial \tilde{u}}{\partial L \tilde{x}} + \frac{\kappa_\infty a_\infty^2}{R} \tilde{\kappa} \frac{\partial \tilde{T}}{\partial L \tilde{x}} \end{pmatrix} \quad (7)$$

Rearranging:

$$\frac{\rho_\infty a_\infty}{L} \frac{\partial}{\partial \tilde{t}} \begin{pmatrix} \tilde{\rho} \\ a_\infty \tilde{\rho} \tilde{u} \\ a_\infty^2 \tilde{e} \end{pmatrix} + \frac{\rho_\infty a_\infty}{L} \frac{\partial}{\partial \tilde{x}} \begin{pmatrix} \tilde{\rho} \tilde{u} \\ a_\infty \tilde{p} + a_\infty \tilde{\rho} \tilde{u}^2 \\ a_\infty^2 (\tilde{e} + \tilde{p}) \tilde{u} \end{pmatrix} = \frac{\mu_\infty}{L^2} \frac{\partial}{\partial \tilde{x}} \begin{pmatrix} 0 \\ \frac{4}{3} a_\infty \tilde{\mu} \frac{\partial \tilde{u}}{\partial \tilde{x}} \\ \frac{4}{3} a_\infty^2 \tilde{\mu} \tilde{u} \frac{\partial \tilde{u}}{\partial \tilde{x}} + \frac{\kappa_\infty a_\infty^2}{\mu_\infty R} \tilde{\kappa} \frac{\partial \tilde{T}}{\partial \tilde{x}} \end{pmatrix} \quad (8)$$

Dividing the second equation by a_∞ , the third equation by a_∞^2 , and the whole set of equations by $\frac{\rho_\infty a_\infty}{L}$ we get:

$$\frac{\partial}{\partial \tilde{t}} \begin{pmatrix} \tilde{\rho} \\ \tilde{\rho} \tilde{u} \\ \tilde{e} \end{pmatrix} + \frac{\partial}{\partial \tilde{x}} \begin{pmatrix} \tilde{\rho} \tilde{u} \\ \tilde{p} + \tilde{\rho} \tilde{u}^2 \\ (\tilde{e} + \tilde{p}) \tilde{u} \end{pmatrix} = \frac{\mu_\infty}{L \rho_\infty a_\infty} \frac{\partial}{\partial \tilde{x}} \begin{pmatrix} 0 \\ \frac{4}{3} \tilde{\mu} \frac{\partial \tilde{u}}{\partial \tilde{x}} \\ \frac{4}{3} \tilde{\mu} \tilde{u} \frac{\partial \tilde{u}}{\partial \tilde{x}} + \frac{\kappa_\infty}{\mu_\infty R} \tilde{\kappa} \frac{\partial \tilde{T}}{\partial \tilde{x}} \end{pmatrix} \quad (9)$$

The Reynolds number and the mach number far away are defined as:

$$\begin{aligned} M_\infty &= \frac{u_\infty}{a_\infty} & Re_{L\infty} &= \frac{\rho_\infty u_\infty L}{\mu_\infty} \\ &\Downarrow & & \\ \frac{\mu_\infty}{L \rho_\infty a_\infty} &= \frac{M_\infty}{Re_{L\infty}} \end{aligned} \quad (10)$$

The Prandtl number far away is defined as:

$$\begin{aligned} Pr_\infty &= \frac{c_p \mu_\infty}{\kappa_\infty} \\ &\Downarrow \\ \frac{\kappa_\infty}{\mu_\infty R} &= \frac{c_p}{Pr_\infty (c_p - c_v)} = \frac{\gamma}{Pr_\infty (\gamma - 1)} \end{aligned} \quad (11)$$

Substituting into the normalized N-S equations:

$$\frac{\partial \tilde{Q}}{\partial \tilde{t}} + \frac{\partial \tilde{E}}{\partial \tilde{x}} = \frac{M_\infty}{Re_{L\infty}} \frac{\partial \tilde{E}_\nu}{\partial \tilde{x}} \quad (12)$$

Where:

$$\tilde{Q} = \begin{pmatrix} \tilde{\rho} \\ \tilde{\rho} \tilde{u} \\ \tilde{e} \end{pmatrix}, \quad \tilde{E} = \begin{pmatrix} \tilde{\rho} \tilde{u} \\ \tilde{p} + \tilde{\rho} \tilde{u}^2 \\ (\tilde{e} + \tilde{p}) \tilde{u} \end{pmatrix}, \quad \tilde{E}_\nu = \begin{pmatrix} 0 \\ \frac{4}{3} \tilde{\mu} \frac{\partial \tilde{u}}{\partial \tilde{x}} \\ \frac{4}{3} \tilde{\mu} \tilde{u} \frac{\partial \tilde{u}}{\partial \tilde{x}} + \frac{\gamma}{Pr_\infty (\gamma - 1)} \tilde{\kappa} \frac{\partial \tilde{T}}{\partial \tilde{x}} \end{pmatrix} \quad (13)$$

The normalized Navier-Stokes equations are:

$$\frac{\partial \tilde{Q}}{\partial \tilde{t}} + \frac{\partial \tilde{E}}{\partial \tilde{x}} = \frac{M_\infty}{Re_{L\infty}} \frac{\partial \tilde{V}_1}{\partial \tilde{x}} \quad (14)$$

Where:

$$\tilde{V}_1 = \tilde{V}_1(\tilde{Q}, \tilde{Q}_x) = \tilde{E}_\nu$$

3 The Computational Domain

3.1 Discretization

The physical domain $[x_I, x_F]$ is discretized into N equispaced cells. The size of each cell is there for:

$$\Delta x = \frac{x_F - x_I}{N} = \frac{L}{N} \quad (15)$$

so the x coordinate of the i-th cell x_i is:

$$x_i = x_I + \frac{1}{2} \Delta x + \Delta x \cdot (i - 1) \quad \text{when starting from } i = 1 \quad (16)$$



3.2 Boundary Conditions

In order to set the boundary conditions on the edge faces we will define ghost cells that will be calculated as follows:

$$\begin{aligned} u_{(i=0)} &= -u_{(i=1)} \\ u_{(i=N+1)} &= -u_{(i=N)} \end{aligned} \quad (17)$$

in order to maintain velocity zero on the boundary and like so:

$$\begin{aligned} T_{(i=0)} &= T_{(i=1)} \\ T_{(i=N+1)} &= T_{(i=N)} \end{aligned} \quad (18)$$

in order to maintain adiabatic boundary conditions. Since the gradient of the pressure on the wall is zero, we get:

$$\begin{aligned} p_{(i=0)} &= p_{(i=1)} \\ p_{(i=N+1)} &= p_{(i=N)} \end{aligned} \quad (19)$$

From equations 2, 18, and 19 we can conclude:

$$\begin{aligned} \rho_{(i=0)} &= \rho_{(i=1)} \\ \rho_{(i=N+1)} &= \rho_{(i=N)} \end{aligned} \quad (20)$$

and from equations 2, 17, 19, and 20 we can conclude:

$$\begin{aligned} e_{(i=0)} &= e_{(i=1)} \\ e_{(i=N+1)} &= e_{(i=N)} \end{aligned} \quad (21)$$



4 The Numerical Schemes

4.1 Jacobian Matrices of The Navier-Stokes Equations

We can rewrite Eq.14 as:

$$\begin{aligned}\frac{\partial \tilde{Q}}{\partial \tilde{t}} &= -\frac{\partial \tilde{E}}{\partial \tilde{x}} - \frac{M_\infty}{Re_{L\infty}} \frac{\partial \tilde{V}_1}{\partial \tilde{x}} \\ \frac{\partial \tilde{Q}}{\partial \tilde{t}} &= -\underbrace{\frac{\partial \tilde{E}}{\partial \tilde{Q}}}_{\tilde{A}} \frac{\partial \tilde{Q}}{\partial \tilde{x}} - \frac{M_\infty}{Re_{L\infty}} \left(\underbrace{\frac{\partial \tilde{V}_1}{\partial \tilde{Q}}}_{\tilde{P}} \frac{\partial \tilde{Q}}{\partial \tilde{x}} + \underbrace{\frac{\partial \tilde{V}_1}{\partial \tilde{Q}_x}}_{\tilde{R}} \frac{\partial \tilde{Q}_x}{\partial \tilde{x}} \right)\end{aligned}\quad (22)$$

Where:

$$\begin{aligned}\tilde{A} &= \begin{pmatrix} 0 & 1 & 0 \\ \frac{\gamma-3}{2} \tilde{u}^2 & (3-\gamma) \tilde{u} & \gamma-1 \\ -\frac{\gamma \tilde{e} \tilde{u}}{\tilde{\rho}} - (\gamma-1) \tilde{u}^3 & \frac{\gamma \tilde{e}}{\tilde{\rho}} - \frac{3(\gamma-1)}{2} \tilde{u}^2 & \gamma \tilde{u} \end{pmatrix} \\ \tilde{P} - \tilde{R}_x &= -\frac{1}{\rho} \begin{pmatrix} 0 & 0 & 0 \\ -\tilde{u} \left(\frac{4}{3} \tilde{\mu} \right)_x & \left(\frac{4}{3} \tilde{\mu} \right)_x & 0 \\ -\tilde{u}^2 \left(\frac{4}{3} \tilde{\mu} \right)_x & \tilde{u} \left(\frac{4}{3} \tilde{\mu} \right)_x & 0 \end{pmatrix} \\ \tilde{R} &= -\frac{1}{\rho} \begin{pmatrix} 0 & 0 & 0 \\ \frac{4}{3} \tilde{u} \tilde{\mu} & -\frac{4}{3} \tilde{\mu} & 0 \\ \left(\frac{4}{3} \tilde{\mu} - \alpha \frac{\tilde{\kappa}}{c_v} \right) \tilde{u}^2 + \alpha \frac{\tilde{\kappa}}{c_v} \frac{\tilde{e}}{\tilde{\rho}} & -\left(\frac{4}{3} \tilde{\mu} - \alpha \frac{\tilde{\kappa}}{c_v} \right) \tilde{u} & -\alpha \frac{\tilde{\kappa}}{c_v} \end{pmatrix}\end{aligned}\quad (23)$$

and α is:

$$\alpha = \frac{\gamma}{Pr_\infty(\gamma-1)}$$

4.2 Linearization In Time

4.2.1 \tilde{E}_i^{n+1} Estimation

$$\tilde{E}_i^{n+1} = \tilde{E}_i^n + \underbrace{\frac{\partial \tilde{E}}{\partial \tilde{Q}} \bigg|_i}_{\tilde{A}_i^n} \Delta \tilde{Q}_i^n + \text{H.O.T}\quad (24)$$

$$\tilde{E}_i^{n+1} = \tilde{E}_i^n + \tilde{A}_i^n \Delta \tilde{Q}_i^n$$



4.2.2 \tilde{V}_i^{n+1} Estimation

$$\tilde{V}_{1,i}^{n+1} = \tilde{V}_{1,i}^n + \underbrace{\left. \frac{\partial \tilde{V}_1}{\partial \tilde{Q}} \right|_i^n}_{\tilde{P}_i^n} \Delta \tilde{Q}_i^n + \underbrace{\left. \frac{\partial \tilde{V}_1}{\partial \tilde{Q}_x} \right|_i^n}_{\tilde{R}_i^n} \Delta \tilde{Q}_{x_i}^n + \text{H.O.T} \quad (25)$$

$$\tilde{V}_{1,i}^{n+1} = \tilde{V}_{1,i}^n + \tilde{P}_i^n \Delta \tilde{Q}_i^n + \tilde{R}_i^n \Delta \tilde{Q}_{x_i}^n$$

The difficulty stems from the fact that the solution vector is $\Delta \tilde{Q}$ and not $\Delta \tilde{Q}_x$. This can be solved by a linearization of the term $\Delta \tilde{Q}_x$ which can be conducted using the following relation:

$$\begin{aligned} \frac{\partial \left(\tilde{R} \Delta \tilde{Q} \right)_i^n}{\partial \tilde{x}} &= \frac{\partial \tilde{R}_i^n}{\partial \tilde{x}} \Delta \tilde{Q}_i^n + \tilde{R}_i^n \frac{\partial \Delta \tilde{Q}_i^n}{\partial \tilde{x}} = \frac{\partial \tilde{R}}{\partial \tilde{x}} \Delta \tilde{Q} + \tilde{R}_i^n \Delta \tilde{Q}_{x_i}^n \\ &\Downarrow \\ \tilde{V}_{1,i}^{n+1} &= \tilde{V}_{1,i}^n + \left(\tilde{P} - \tilde{R}_x \right)_i^n \Delta \tilde{Q}_i^n + \frac{\partial}{\partial \tilde{x}} \left(\tilde{R} \Delta \tilde{Q} \right)_i^n \end{aligned} \quad (26)$$

4.3 First Order Approximate Riemann Roe Method

4.4 First Order Steger-Warming – Explicit

The A matrix (from Eq.23) is a diagonalizable matrix and can be written as:

$$\begin{aligned} \tilde{A} &= \tilde{T} \tilde{\Lambda} \tilde{T}^{-1} \\ \tilde{T} &= \begin{pmatrix} 1 & \frac{\tilde{\rho}}{2\tilde{a}} & -\frac{\tilde{\rho}}{2\tilde{a}} \\ \tilde{u} & \frac{\tilde{\rho}}{2\tilde{a}} (\tilde{u} + \tilde{a}) & -\frac{\tilde{\rho}}{2\tilde{a}} (\tilde{u} - \tilde{a}) \\ \frac{\tilde{u}^2}{2} & \frac{\tilde{\rho}}{2\tilde{a}} \left(\frac{\tilde{u}^2}{2} + \tilde{u}\tilde{a} + \frac{\tilde{a}^2}{\gamma-1} \right) & -\frac{\tilde{\rho}}{2\tilde{a}} \left(\frac{\tilde{u}^2}{2} - \tilde{u}\tilde{a} + \frac{\tilde{a}^2}{\gamma-1} \right) \end{pmatrix} \\ \tilde{\Lambda} &= \begin{pmatrix} \tilde{u} & 0 & 0 \\ 0 & \tilde{u} + \tilde{a} & 0 \\ 0 & 0 & \tilde{u} - \tilde{a} \end{pmatrix} \\ \tilde{T}^{-1} &= \begin{pmatrix} 1 - \frac{\gamma-1}{2} \frac{\tilde{u}^2}{\tilde{a}^2} & (\gamma-1) \frac{\tilde{u}^2}{\tilde{a}^2} & -\frac{\gamma-1}{\tilde{a}^2} \\ \frac{1}{\tilde{\rho}\tilde{a}} ((\gamma-1) \tilde{u}^2 - \tilde{u}\tilde{a}) & \frac{1}{\tilde{\rho}\tilde{a}} (\tilde{a} - (\gamma-1) \tilde{u}) & \frac{\gamma-1}{\tilde{\rho}\tilde{a}} \\ -\frac{1}{\tilde{\rho}\tilde{a}} ((\gamma-1) \tilde{u}^2 + \tilde{u}\tilde{a}) & \frac{1}{\tilde{\rho}\tilde{a}} (\tilde{a} + (\gamma-1) \tilde{u}) & -\frac{\gamma-1}{\tilde{\rho}\tilde{a}} \end{pmatrix} \end{aligned} \quad (27)$$

Where:

$$\tilde{a} = \sqrt{\frac{\gamma \tilde{p}}{\tilde{\rho}}}$$



Let the $\tilde{\Lambda}^\pm$ matrix be defined as:

$$\tilde{\Lambda}^\pm = \begin{pmatrix} \frac{\tilde{u} \pm |\tilde{u}|}{2} & 0 & 0 \\ 0 & \frac{\tilde{u} + \tilde{a} \pm |\tilde{u} + \tilde{a}|}{2} & 0 \\ 0 & 0 & \frac{\tilde{u} - \tilde{a} \pm |\tilde{u} - \tilde{a}|}{2} \end{pmatrix} \quad (28)$$

Where the matrix $\tilde{\Lambda}^+$ contains only positive eigenvalues and the matrix $\tilde{\Lambda}^-$ contains only negative eigenvalues.

Define:

$$\begin{aligned} \tilde{A}^+ &\triangleq \tilde{T} \tilde{\Lambda}^+ \tilde{T}^{-1} & \tilde{A} &= \tilde{A}^+ + \tilde{A}^- \\ \tilde{A}^- &\triangleq \tilde{T} \tilde{\Lambda}^- \tilde{T}^{-1} & |\tilde{A}| &\triangleq \tilde{A}^+ - \tilde{A}^- \end{aligned} \quad (29)$$

Assuming a perfect gas, the flux vector $\tilde{E}_{(Q)}$ is a homogeneous function of degree one in \tilde{Q} , meaning:

$$\forall \alpha \quad \tilde{E}_{(\alpha \tilde{Q})} = \alpha \tilde{E}_{(\tilde{Q})}$$

The homogeneity allows to rewrite the flux vector \tilde{E} using Eq.22 as:

$$\tilde{E} = \tilde{A} \tilde{Q} = (\tilde{A}^+ + \tilde{A}^-) \tilde{Q} = \underbrace{\tilde{A}^+ \tilde{Q}}_{\tilde{E}^+} + \underbrace{\tilde{A}^- \tilde{Q}}_{\tilde{E}^-} = \tilde{E}^+ + \tilde{E}^- \quad (30)$$

There is a discontinuity and deference between \tilde{E}^+, \tilde{E}^- . To eliminate the discontinuities and guarantee a smooth transition through critical points (sonic points or stagnation points), a blending function is introduced together with a blending parameter ε . An appropriate choice of the blending parameter has to be chosen.

$$\begin{aligned} \tilde{\lambda}^+ &= \frac{\tilde{\lambda} + |\tilde{\lambda}|}{2} & \tilde{\lambda}^{+'} &= \frac{\tilde{\lambda} + \sqrt{\tilde{\lambda}^2 + \varepsilon^2}}{2} \\ & \Rightarrow & & \\ \tilde{\lambda}^- &= \frac{\tilde{\lambda} - |\tilde{\lambda}|}{2} & \tilde{\lambda}^{-'} &= \frac{\tilde{\lambda} - \sqrt{\tilde{\lambda}^2 + \varepsilon^2}}{2} \end{aligned} \quad (31)$$

Rewriting the conservation law form of the N-S equations Eq.14 using Eq.30:

$$\frac{\partial \tilde{Q}}{\partial \tilde{t}} = -\frac{\partial \tilde{E}^+}{\partial \tilde{x}} - \frac{\partial \tilde{E}^-}{\partial \tilde{x}} + \frac{M_\infty}{Re_{L\infty}} \frac{\partial \tilde{V}_1}{\partial \tilde{x}} \quad (32)$$

A simple, explicit, first order (in space and time) scheme in delta form is obtained using:

$$\Delta \tilde{Q}_i^n = -\frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\nabla \tilde{E}_i^{+n} + \Delta \tilde{E}_i^{-n} - \frac{M_\infty}{Re_{L\infty}} \delta \tilde{V}_{1,i}^n \right) \quad (33)$$

And advancing the solution by:

$$\tilde{Q}_i^{n+1} = \Delta \tilde{Q}_i^n + \tilde{Q}_i^n \quad (34)$$

4.4.1 Finite Volume Formulation

Rearranging Eq.33 using the finite volume notation:

$$\begin{aligned} \Delta \tilde{Q}_i^n &= -\frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\tilde{E}_i^{+n} - \tilde{E}_{i-1}^{+n} + \tilde{E}_{i+1}^{-n} - \tilde{E}_i^{-n} - \frac{M_\infty}{Re_{L\infty}} \left(\tilde{V}_{1,i+\frac{1}{2}}^n - \tilde{V}_{1,i-\frac{1}{2}}^n \right) \right) \\ \Delta \tilde{Q}_i^n &= -\frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\left(\tilde{E}_i^{+n} + \tilde{E}_{i+1}^{-n} \right) - \left(\tilde{E}_{i-1}^{+n} + \tilde{E}_i^{-n} \right) - \frac{M_\infty}{Re_{L\infty}} \left(\tilde{V}_{1,i+\frac{1}{2}}^n - \tilde{V}_{1,i-\frac{1}{2}}^n \right) \right) \end{aligned} \quad (35)$$



Define:

$$\begin{aligned}
 \tilde{E}_{i+\frac{1}{2}} &\triangleq \tilde{E}_i^+ + E_{i+1}^- \\
 &= \tilde{A}_i^+ \tilde{Q}_i + \tilde{A}_{i+1}^- \tilde{Q}_{i+1} \\
 &\equiv \tilde{\tilde{E}}_{i+\frac{1}{2}}
 \end{aligned} \tag{36}$$

Finally we get:

$$\Delta \tilde{Q}_i^n = -\frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\tilde{\tilde{E}}_{i+\frac{1}{2}}^n - \tilde{\tilde{E}}_{i-\frac{1}{2}}^n - \frac{M_\infty}{Re_{L\infty}} \left(\tilde{V}_{1,i+\frac{1}{2}}^n - \tilde{V}_{1,i-\frac{1}{2}}^n \right) \right) \tag{37}$$

4.4.2 Calculating $\tilde{V}_{1,i+\frac{1}{2}}$

$$\tilde{V}_{1,i+\frac{1}{2}} = \begin{pmatrix} 0 \\ \frac{4}{3} \tilde{\mu}|_{i+\frac{1}{2}} \frac{\partial \tilde{u}}{\partial \tilde{x}} \Big|_{i+\frac{1}{2}} \\ \frac{4}{3} \tilde{\mu}|_{i+\frac{1}{2}} \tilde{u}|_{i+\frac{1}{2}} \frac{\partial \tilde{u}}{\partial \tilde{x}} \Big|_{i+\frac{1}{2}} + \frac{\gamma}{Pr_\infty(\gamma-1)} \tilde{\kappa}|_{i+\frac{1}{2}} \frac{\partial \tilde{T}}{\partial \tilde{x}} \Big|_{i+\frac{1}{2}} \end{pmatrix} \tag{38}$$

Where:

$$\begin{aligned}
 \frac{\partial \tilde{u}}{\partial \tilde{x}} \Big|_{i+\frac{1}{2}} &= \frac{\tilde{u}_{i+1} - \tilde{u}_i}{\Delta \tilde{x}}, & \tilde{\mu}|_{i+\frac{1}{2}} &= \frac{\tilde{\mu}_{i+1} + \tilde{\mu}_i}{2} \\
 \frac{\partial \tilde{T}}{\partial \tilde{x}} \Big|_{i+\frac{1}{2}} &= \frac{\tilde{T}_{i+1} - \tilde{T}_i}{\Delta \tilde{x}}, & \tilde{\kappa}|_{i+\frac{1}{2}} &= \frac{\tilde{\kappa}_{i+1} + \tilde{\kappa}_i}{2}
 \end{aligned} \tag{39}$$

$$\tilde{u}|_{i+\frac{1}{2}} = \frac{\tilde{u}_{i+1} + \tilde{u}_i}{2}$$

4.5 First Order Steger-Warming – Implicit

The Implicit Steger-Warming scheme starts from:

$$\begin{aligned}
 \frac{\Delta \tilde{Q}_i^n}{\Delta \tilde{t}} &= -\frac{\partial \tilde{E}_i^{n+1}}{\partial \tilde{x}} + \frac{M_\infty}{Re_{L\infty}} \frac{\partial \tilde{V}_{1,i}^{n+1}}{\partial \tilde{x}} \\
 \Delta \tilde{Q}_i^n &= -\Delta \tilde{t} \frac{\partial}{\partial \tilde{x}} \left(\tilde{E}_i^n + \tilde{A}_i^n \Delta \tilde{Q}_i^n \right) + \Delta \tilde{t} \frac{M_\infty}{Re_{L\infty}} \frac{\partial}{\partial \tilde{x}} \left(\tilde{V}_{1,i}^n + \left(\tilde{P} - \tilde{R}_x \right)_i^n \Delta \tilde{Q}_i^n + \frac{\partial}{\partial \tilde{x}} \left(\tilde{R} \Delta \tilde{Q} \right)_i^n \right)
 \end{aligned} \tag{40}$$

Rearranging in delta form:

$$\begin{aligned}
 \left(I + \Delta \tilde{t} \left(\frac{\partial}{\partial \tilde{x}} \left[\tilde{A} - \left(\tilde{P} - \tilde{R}_x \right) \right]_i^n - \frac{\partial^2 \tilde{R}_i^n}{\partial \tilde{x}^2} \right) \right) \Delta \tilde{Q}_i^n &= \text{RHS}_i^n \\
 \text{RHS}_i^n &= -\Delta \tilde{t} \frac{\partial}{\partial \tilde{x}} \left(\tilde{E}^+ + \tilde{E}^- - \frac{M_\infty}{Re_{L\infty}} \tilde{V}_1 \right)_i^n
 \end{aligned} \tag{41}$$



$$\begin{aligned} \left(I + \frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\left[\nabla \tilde{A}^+ + \Delta \tilde{A}^- - \frac{D_0}{2} (\tilde{P} - \tilde{R}_x) \right]_i^n - \frac{\delta^2}{\Delta \tilde{x}} \tilde{R}_i^n \right) \right) \Delta \tilde{Q}_i^n &= \text{RHS}_i^n \\ \text{RHS}_i^n &= -\frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\nabla \tilde{E}^+ + \Delta \tilde{E}^- - \frac{M_\infty}{Re_{L\infty}} \delta \tilde{V}_1 \right)_i^n \end{aligned} \quad (42)$$

Rewrite using the finite volume notation for the inviscid terms like in Eq.37:

$$\begin{aligned} \left(I + \frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\left[\nabla \tilde{A}^+ + \Delta \tilde{A}^- - \frac{D_0}{2} (\tilde{P} - \tilde{R}_x) \right]_i^n - \frac{\delta^2}{\Delta \tilde{x}} \tilde{R}_i^n \right) \right) \Delta \tilde{Q}_i^n &= \text{RHS}_i^n \\ \text{RHS}_i^n &= -\frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\tilde{E}_{i+\frac{1}{2}}^n - \tilde{E}_{i-\frac{1}{2}}^n - \frac{M_\infty}{Re_{L\infty}} (\tilde{V}_{1,i+\frac{1}{2}}^n - \tilde{V}_{1,i-\frac{1}{2}}^n) \right) \end{aligned} \quad (43)$$

and opening the delta form on the LHS:

$$\begin{aligned} \text{LHS}_i^n &= \Delta \tilde{Q}_i^n + \frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\nabla \tilde{A}_i^{+n} \Delta \tilde{Q}_i^n + \Delta \tilde{A}_i^{-n} \Delta \tilde{Q}_i^n - \frac{D_0}{2} (\tilde{P} - \tilde{R}_x)_i^n \Delta \tilde{Q}_i^n - \frac{\delta^2}{\Delta \tilde{x}} \tilde{R}_i^n \Delta \tilde{Q}_i^n \right) \\ &= \Delta \tilde{Q}_i^n + \frac{\Delta \tilde{t}}{\Delta \tilde{x}} \left(\tilde{A}_i^{+n} \Delta \tilde{Q}_i^n - \tilde{A}_{i-1}^{+n} \Delta \tilde{Q}_{i-1}^n + \tilde{A}_{i+1}^{-n} \Delta \tilde{Q}_{i+1}^n - \tilde{A}_i^{-n} \Delta \tilde{Q}_i^n - \right. \\ &\quad \left. - \frac{1}{2} \left[(\tilde{P} - \tilde{R}_x)_{i+1}^n \Delta \tilde{Q}_{i+1}^n - (\tilde{P} - \tilde{R}_x)_{i-1}^n \Delta \tilde{Q}_{i-1}^n \right] - \right. \\ &\quad \left. - \frac{1}{\Delta \tilde{x}} \left[\tilde{R}_{i+1}^n \Delta \tilde{Q}_{i+1}^n - 2\tilde{R}_i^n \Delta \tilde{Q}_i^n + \tilde{R}_{i-1}^n \Delta \tilde{Q}_{i-1}^n \right] \right) \\ &= \Theta_i^n \Delta \tilde{Q}_{i-1}^n + \Phi_i^n \Delta \tilde{Q}_i^n + \Psi_i^n \Delta \tilde{Q}_{i+1}^n \end{aligned} \quad (44)$$

Where:

$$\begin{aligned} \Theta_i^n &= -\frac{\Delta \tilde{t}}{\Delta \tilde{x}} \tilde{A}_{i-1}^{+n} + \frac{\Delta \tilde{t}}{2\Delta \tilde{x}} (\tilde{P} - \tilde{R}_x)_{i-1}^n - \frac{\Delta \tilde{t}}{\Delta \tilde{x}^2} \tilde{R}_{i-1}^n \\ \Phi_i^n &= I + \frac{\Delta \tilde{t}}{\Delta \tilde{x}} (\tilde{A}_i^{+n} - \tilde{A}_i^{-n}) + 2\frac{\Delta \tilde{t}}{\Delta \tilde{x}^2} \tilde{R}_i^n \\ \Psi_i^n &= \frac{\Delta \tilde{t}}{\Delta \tilde{x}} \tilde{A}_{i+1}^{-n} - \frac{\Delta \tilde{t}}{2\Delta \tilde{x}} (\tilde{P} - \tilde{R}_x)_{i+1}^n - \frac{\Delta \tilde{t}}{\Delta \tilde{x}^2} \tilde{R}_{i+1}^n \end{aligned} \quad (45)$$

For the implicit Steger-Warming scheme a matrix inversion is needed as follows:

$$\begin{pmatrix} \Phi_1^n & \Psi_1^n & 0 & \cdots & \cdots & \cdots & 0 \\ \Theta_2^n & \Phi_2^n & \Psi_2^n & 0 & \cdots & \cdots & 0 \\ 0 & \ddots & \ddots & \ddots & 0 & \cdots & 0 \\ 0 & 0 & \Theta_i^n & \Phi_i^n & \Psi_i^n & 0 & 0 \\ 0 & \cdots & 0 & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & \cdots & 0 & \Theta_{N-1}^n & \Phi_{N-1}^n & \Psi_{N-1}^n \\ 0 & \cdots & \cdots & \cdots & 0 & \Theta_N^n & \Phi_N^n \end{pmatrix} \begin{pmatrix} \Delta \tilde{Q}_1^n \\ \Delta \tilde{Q}_2^n \\ \cdots \\ \cdots \\ \cdots \\ \Delta \tilde{Q}_{N-1}^n \\ \Delta \tilde{Q}_N^n \end{pmatrix} = \begin{pmatrix} \text{RHS}_1^n \\ \text{RHS}_2^n \\ \cdots \\ \cdots \\ \cdots \\ \text{RHS}_{N-1}^n \\ \text{RHS}_N^n \end{pmatrix} \quad (46)$$