



הטכניון – מכון טכנולוגי לישראל

**NUMERICAL METHODS IN AEROSPACE
ENGINEERING**

HOMEWORK ASSIGNMENT x

סמסטר אביב תשפ"ה

SPRING SEMESTER 2025

GRADE	OUT OF	CHAPTER
	2	ABSTRACT
	2	CONTENTS, STYLE &C.
	4	PHYSICAL PROBLEM
	4	MATHEMATICAL MODEL
	26	NUMERICAL METHODS
	20	INFLUENCE OF NUMERICAL METHODS
	20	RESULTS
	2	SUMMARY & CONCLUSIONS
	20	COMPUTER PROGRAM
	100	TOTAL

Almog Dobrescu

ID 214254252

June 2, 2025

Contents

1	The Physical Problem	1
2	The Mathematical Model	1
3	The Numerical Methods	1
3.1	Finite Differencing	1
3.2	Convergence Criteria	2
3.3	Integral Calculation	2
4	Influence of The Numerical Methods	2
4.1	Influence of Number of Elements N	2
4.2	Influence of Convergence Criteria ε	3
4.3	Influence of The Numerical Parameter R	3
5	Results and Discussion	4
5.1	Temperature Distribution	4
5.2	Strain Calculation	5
6	Summary and Conclusion	5
A	Listing of The Computer Program	6
A.1	Parameters	6
A.2	Main Code	6

List of Figures

1	Influence of the number of elements N	2
2	FD - Influence of the convergence criteria ε	3
3	FD - Influence of the convergence criteria R	3
4	Temperature as a Function of r	4
5	Temperature as a Function of r over time	4

Listings

1	Parameters file	6
2	The main file	6

Nomenclature

Δr	difference between two points in space
Δt	difference between two points in time
ε	convergence criteria
i	index in coordinate r
J	final time coordinate
j	index in time
K	diffusivity coefficient
N	number of cell
r	radial coordinate
T	temperature
t	time coordinate



1 The Physical Problem

Heated tubing has many applications, from water cooling in computers to transporting liquid gas. Research has been conducted to investigate the stress-strain relationships and the material properties of a tube subjected to heating and cooling.

2 The Mathematical Model

The following heat equation was in use:

$$\frac{1}{4K} \frac{\partial T}{\partial t} = \frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} \quad , \quad \frac{1}{2} \leq r \leq 1 \quad , \quad 0 < t \quad (1)$$

The boundary and initial conditions of the problem:

$$\begin{aligned} T_{(0.5,t)} &= t \\ T_{(1.0,t)} &= 100 + 40t \\ T_{(r,0)} &= 200 \left(r - \frac{1}{2} \right) \\ K &= 0.1 \end{aligned} \quad (2)$$

The strain I is given by the equation:

$$I = \int_{0.5}^1 \alpha T_{(r,t)} r dr \quad (3)$$

- $\alpha = 10.7$

3 The Numerical Methods

3.1 Finite Differencing

The Heat equation can be rewritten as:

$$\frac{\partial T}{\partial t} = 4K \left(\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} \right) \quad (4)$$

By using central differencing for the spatial derivatives and forward differencing for the time derivative we get the explicit scheme:

$$\begin{aligned} \frac{T_{i,j+1} - T_{i,j}}{\Delta t} + O(\Delta t) &= 4K \left(\frac{T_{i-1,j} - 2T_{i,j} + T_{i+1,j}}{\Delta r^2} + O(\Delta r^2) + \frac{1}{r} \frac{T_{i+1,j} - T_{i-1,j}}{2\Delta r} + O(\Delta r^2) \right) \\ &\Downarrow \\ T_{i,j+1} &= T_{i,j} + 4K\Delta t \left(\frac{T_{i-1,j} - 2T_{i,j} + T_{i+1,j}}{\Delta r^2} + \frac{1}{r} \frac{T_{i+1,j} - T_{i-1,j}}{2\Delta r} \right) + O(\Delta t, \Delta r^2) \end{aligned} \quad (5)$$

- $$T_{i,j+1} = T_{i,j} + 4KR \left(T_{i-1,j} - 2T_{i,j} + T_{i+1,j} + \frac{\Delta r}{2r} (T_{i+1,j} - T_{i-1,j}) \right) + O(\Delta t, \Delta r^2)$$
- $i = 1, 2, \dots, N$
 - $j = 1, 2, \dots, J$
 - $R = \frac{\Delta t}{\Delta r^2}$
 - $T_{(i=0,j)} = j \cdot \Delta t$
 - $T_{(i=N+1,j)} = 100 + 40 \cdot j \cdot \Delta t$



$$\bullet T_{(i,j=0)} = 200 \left(r - \frac{1}{2} \right), \quad 0 \leq r \leq N + 1$$

Δr was calculated as follows:

$$\Delta r = \frac{r_{\max} - r_{\min}}{N + 1} \quad (6)$$

The step size in time Δt was chosen somewhat arbitrary, only to maintain stability.

The system of equation will be solved by Jacobi method for every j . This method adds an iterative index n :

$$T_{i,j+1}^{n+1} = T_{i,j}^n + 4KR \left(T_{i-1,j}^n - 2T_{i,j}^n + T_{i+1,j}^n + \frac{\Delta r}{2r} (T_{i+1,j}^n - T_{i-1,j}^n) \right) \quad (7)$$

3.2 Convergence Criteria

In order determined if the iterative method for solving the system of equation has converged, we will check if the temperature vector at a specific time has changed from step n to step $n+1$ in the following way:

$$\left| T_{i,j}^{n+1} - T_{i,j}^n \right| < \varepsilon \quad (8)$$

3.3 Integral Calculation

The integral to calculate the strain I will be calculated using trapezoid integration:

$$I = \alpha \frac{h}{2} \sum_{i=0}^N T_{(i+1)} + T_{(i)} \quad (9)$$

4 Influence of The Numerical Methods

NOTE - the temperature will be presented in logarithmic scale.

4.1 Influence of Number of Elements N




Figure 1: Influence of the number of elements N

In Fig.1 we can see that for N bigger than 20, the solution does not really change. We can conclude that $N = 20$ is a sufficient number of elements.



4.2 Influence of Convergence Criteria ε




Figure 2: FD - Influence of the convergence criteria ε

From Fig.2 we can conclude that for a convergence criteria smaller than $1e^{-5}$, the solution stays the same. We can determine that $\varepsilon = 1e^{-5}$ is a good choice.

4.3 Influence of The Numerical Parameter R





Figure 3: FD - Influence of the convergence criteria R

From Fig.3 we can conclude that the value of the numerical parameter R has close to no effect on the error of the solution.


5 Results and Discussion

5.1 Temperature Distribution

Using the parameters chosen above, the temperature distribution is therefor:



We can also plot the temperature distribution over time for all the radiuses:





5.2 Strain Calculation

When using the chosen parameters, the calculated strain is:

$$I_{N=20} = 1224.8 \quad (10)$$

However, when using and $N = 40$ $N = 60$, the strain is:

$$I_{N=40} = 1234.0 \quad I_{N=60} = 1237.0 \quad (11)$$

We observed that while $N = 20$ is adequate for convergence of the temperature, a higher number of elements is necessary to accurately calculate the strain. This is due to the order of the integration being $O(h)$.

6 Summary and Conclusion



A Listing of The Computer Program

A.1 Parameters

```

1  N          = 20;
2  epsilon    = 1e-5;
3  max_iteration = 1e6;
4
5  r_max      = 1;
6  r_min      = 0.5;
7  t_start    = 0;
8  t_end      = 10;
9  K          = 0.1;
10 alpha      = 10.7;
11
12 h          = (r_max - r_min) / (N+1);
13 R          = 0.49;
14 % R        = delta_t / h^2;
15 delta_t    = R * h^2;
16 r          = r_min+h*(0:1:N+1);

```

Listing 1: Parameters file

A.2 Main Code

```

1  clc; clear; close all;
2
3  %% Influenc of N =====
4  % =====
5  % =====
6
7  parameters
8  % Ns = [5, 10, 15, 20, 40, 60, 100];
9  Ns = [3, 5, 7, 10];
10
11 results_vec = {};
12 r_vec = {};
13 for Ns_index = 1:length(Ns)
14     N      = Ns(Ns_index);
15     h      = (r_max - r_min) / (N+1);
16     R      = 0.45;
17     delta_t = R * h^2;
18     r      = r_min+h*(0:1:N+1);
19
20     results_vec{Ns_index,1} = solver(r_min, r, K, h, delta_t, t_start, t_end, N, epsilon,
                                     max_iteration);
21 end
22 %%
23 load("effect_of_N_to_60.mat")
24
25 fig1 = figure('Name', '1', 'Position', [50, 250, 900, 600]);
26 size = 15;
27
28 colors = cool(length(results_vec(:,1)))*0.9;
29 for i = length(results_vec(:,1))
30     plot(results_vec{i,1}.r, results_vec{i,1}.Ts(end,:), 'LineWidth', 2, 'Color', colors(
        i,:))

```



```

31 end
32 xlabel('r$\displaystyle\left[\frac{m}{sec}\right]$', 'FontSize',size, 'Interpreter','
    latex')
33 ylabel('T$\left[K\right]$', 'FontSize',size, 'Interpreter','latex')
34 title('Temperature as a Function of The Radius', 'FontSize',size, 'Interpreter','latex')
35 subtitle(sprintf('$N=%g$ $||$ $t=%f[s]$ $||$ $\backslash\varepsilon=%g$ $||$ $R=%g$', results_vec{
    end,1}.N, results_vec{end,1}.t_vec(end), results_vec{end,1}.epsilon, results_vec{end
    ,1}.R), 'FontSize', size, 'Interpreter','latex')
36 grid on
37 grid minor
38 box on
39 % exportgraphics(fig1, 'images/T over r.png','Resolution',400);
40
41 % //////////////////////////////////////
42 % //////////////////////////////////////
43
44 fig2 = figure('Name', '2','Position', [150, 250, 900, 600]);
45 size = 15;
46
47 colors = cool(length(results_vec(:,1))*0.9;
48 lg = {};
49 for i = 1:length(results_vec(:,1))
50     semilogy(results_vec{i,1}.r, results_vec{i,1}.Ts(end,:), 'LineWidth', 2, 'Color',
        colors(i,:))
51     hold on;
52     lg{end+1} = sprintf('N=%g, $\Delta t=%f$', results_vec{i,1}.N, results_vec{i,1}.
        delta_t);
53 end
54 xlabel('r$\displaystyle\left[\frac{m}{sec}\right]$', 'FontSize',size, 'Interpreter','
    latex')
55 ylabel('T$\left[K\right]$', 'FontSize',size, 'Interpreter','latex')
56 title('Temperature as a Function of The Radius For Different N', 'FontSize',size, '
    Interpreter','latex')
57 subtitle(sprintf('$t=%f[s]$ $||$ $\backslash\varepsilon=%g$ $||$ $R=%g$', results_vec{end,1}.t_vec(
    end), results_vec{end,1}.epsilon, results_vec{end,1}.R), 'FontSize', size, '
    Interpreter','latex')
58 legend(lg, 'FontSize',size-2, 'Location','eastoutside','Interpreter','latex')
59 grid on
60 grid minor
61 box on
62 % exportgraphics(fig1, 'images/T over r.png','Resolution',400); exportgraphics(fig2, '
    images/Influenc of N.png','Resolution',400);
63
64 % //////////////////////////////////////
65 % //////////////////////////////////////
66
67 fig3 = figure('Name', '3','Position', [250, 250, 900, 600]);
68 rms_vec = [];
69 Ns = [];
70 for i = 1:length(results_vec(:,1))-1
71     rms_vec(i) = RMS(results_vec{i,1}.r, results_vec{i,1}.Ts(end,:), results_vec{i+1,1}.r
        , results_vec{i+1,1}.Ts(end,:));
72     Ns(i) = results_vec{i,1}.N;
73 end
74
75 plot(Ns, rms_vec, 'LineWidth', 2, 'Color', colors(end,:))
76
77 xlabel('N', 'FontSize',size, 'Interpreter','latex')

```



```

78 ylabel('rms$\left[K\right]$', 'FontSize',size, 'Interpreter','latex')
79 title('The RMS between two solution as a function of N', 'FontSize',size, 'Interpreter','
    latex')
80 subtitle(sprintf('$t=%f[s] $ $||$ $\varepsilon=%g$ $||$ $ R=%g$', results_vec{end,1}.t_vec(
    end), results_vec{end,1}.epsilon, results_vec{end,1}.R), 'FontSize', size, '
    Interpreter','latex')
81 grid on
82 grid minor
83 box on
84 % exportgraphics(fig1, 'images/T over r.png','Resolution',400); exportgraphics(fig2, '
    images/Influenc of N.png','Resolution',400); exportgraphics(fig3, 'images/Influenc of
    N — error.png','Resolution',400);
85
86 %% Influenc of epsilon =====
87 % =====
88 % =====
89
90 parameters
91
92 epsilon_vec = logspace(-1, -10, 10);
93 % epsilon_vec = logspace(-1, -6, 6);
94
95 results_vec = {};
96 r_vec = {};
97 for eps_index = 1:length(epsilon_vec)
98     epsilon = epsilon_vec(eps_index);
99     results_vec{eps_index,1} = solver(r_min, r, K, h, delta_t, t_start, t_end, N, epsilon
        , max_iteration);
100 end
101 %%
102 load("effect_of_epsilon_—1 to —10.mat")
103
104 fig4 = figure('Name', '4','Position', [350, 250, 900, 600]);
105 size = 15;
106
107 colors = cool(length(results_vec(:,1)))*0.9;
108 lg = {};
109 for i = 1:length(results_vec(:,1))
110     semilogy(results_vec{i,1}.r, results_vec{i,1}.Ts(end,:), 'LineWidth', 2, 'Color',
        colors(i,:))
111     hold on;
112     lg{end+1} = sprintf('$\varepsilon=%g$', results_vec{i,1}.epsilon);
113 end
114 xlabel('$r\displaystyle\left[\frac{m}{sec}\right]$', 'FontSize',size, 'Interpreter','
    latex')
115 ylabel('$T\left[K\right]$', 'FontSize',size, 'Interpreter','latex')
116 title('Temperature as a Function of The Radius For Different $\varepsilon$', 'FontSize',
    size, 'Interpreter','latex')
117 subtitle(sprintf('$t=%f[s] $ $||$ $ N=%g$ $||$ $ R=%g$', results_vec{end,1}.t_vec(end),
    results_vec{end,1}.N, results_vec{end,1}.R), 'FontSize', size, 'Interpreter','latex')
118 legend(lg, 'FontSize',size-2, 'Location','eastoutside','Interpreter','latex')
119 grid on
120 grid minor
121 box on
122 % exportgraphics(fig4, 'images/Influenc of epsilon.png','Resolution',400);
123
124 fig5 = figure('Name', '5','Position', [450, 250, 900, 600]);
125 rms_vec = [];

```



```

126 eps_vec = [];
127 for i = 1:length(results_vec(:,1))-1
128     rms_vec(i) = RMS(results_vec{i,1}.r, results_vec{i,1}.Ts(end,:), results_vec{i+1,1}.r
        , results_vec{i+1,1}.Ts(end,:));
129     eps_vec(i) = results_vec{i,1}.epsilon;
130 end
131
132 loglog(eps_vec, rms_vec, 'LineWidth', 2, 'Color', colors(end,:))
133
134 xlabel('$\varepsilon$', 'FontSize',size, 'Interpreter','latex')
135 ylabel('rms$\left[K\right]$', 'FontSize',size, 'Interpreter','latex')
136 title('The RMS between two solution as a function of $\varepsilon$', 'FontSize',size, '
    Interpreter','latex')
137 subtitle(sprintf('$t=f[s]$$$ $N=%g$ $$$ R=%g', results_vec{end,1}.t_vec(end),
    results_vec{end,1}.N, results_vec{end,1}.R), 'FontSize', size, 'Interpreter','latex')
138 grid on
139 grid minor
140 box on
141 % exportgraphics(fig4, 'images/Influenc of epsilon.png','Resolution',400); exportgraphics
    (fig5, 'images/Influenc of epsilon — error.png','Resolution',400);
142
143 %% Influenc of R =====
144 % =====
145 % =====
146
147 parameters
148
149 % Rs = linspace(0.05,0.49,10);
150 Rs = linspace(0.45, 0.3, 4);
151
152 results_vec = {};
153 r_vec = {};
154 for R_index = 1:length(Rs)
155     R = Rs(R_index);
156     delta_t = R * h^2;
157     results_vec{R_index,1} = solver(r_min, r, K, h, delta_t, t_start, t_end, N, epsilon,
        max_iteration);
158 end
159 %%
160 load("effect_of_R_0.05_to_0.49.mat")
161
162 fig6 = figure('Name', '6','Position', [550, 250, 900, 600]);
163 size = 15;
164
165 colors = cool(length(results_vec(:,1)))*0.9;
166 lg = {};
167 for i = 1:length(results_vec(:,1))
168     semilogy(results_vec{i,1}.r, results_vec{i,1}.Ts(end,:), 'LineWidth', 2, 'Color',
        colors(i,:))
169     hold on;
170     lg{end+1} = sprintf('$R=%g$', results_vec{i,1}.R);
171 end
172 xlabel('$r\displaystyle\left[\frac{m}{sec}\right]$', 'FontSize',size, 'Interpreter','
    latex')
173 ylabel('$T\left[K\right]$', 'FontSize',size, 'Interpreter','latex')
174 title('Temperature as a Function of The Radius For Different R', 'FontSize',size, '
    Interpreter','latex')

```




```

226 load("effect_of_N_to_60.mat")
227 result = results_vec{end};
228 sum_N60 = 0;
229 for i = [0:result.N+1-1]+1
230
231     sum_N60 = sum_N60 + (result.Ts(end,i)+result.Ts(end,i+1))*result.r(i);
232 end
233 sum_N60 = sum_N60 * 10.7 * 0.5 * result.h
234
235
236
237
238 %% Functions =====
239 % =====
240 % =====
241
242 function T = init_fuild(r_min, h, N)
243     i = 0:1:N+1;
244     r = r_min+h * i;
245     T = 200 * (r - 0.5);
246 end
247
248 function T = set_BC(T, t)
249     T(1) = t;
250     T(end) = 100 + 40 * t;
251 end
252
253 function T_next = step_space_jacobi(T_current, r, K, h, delta_t, N)
254     T_next(1) = T_current(1);
255     T_next(N+1+1) = T_current(N+1+1);
256     for i = [1:N]+1
257         T_next(i) = T_current(i) + 4 * K * delta_t * ((T_current(i+1) - 2 * T_current(i)
                + T_current(i-1)) / (h^2) + 1 / r(i) * (T_current(i+1) - T_current(i-1)) /
                (2*h));
258         % T_next(i) = T_current(i) + 4 * K * delta_t * ((T_current(i+1) - 2 * T_current(i)
                + T_next(i-1)) / (h^2) + 1 / r(i) * (T_current(i+1) - T_next(i-1)) / (2*h))
                ;
259     end
260 end
261
262 function converged = check_convergence(T_next, T_current, epsilon, N)
263     converged = true;
264     for i = [1:N]+1
265         if abs(T_next(i) - T_current(i)) > epsilon
266             converged = false;
267             return
268         end
269     end
270 end
271
272 function T_next_t = solve_for_specific_t_jacobi(T_current_t, r, K, h, delta_t, N, epsilon
    , max_iteration)
273     T_current = T_current_t;
274     for n = 1:max_iteration
275         T_next = step_space_jacobi(T_current, r, K, h, delta_t, N);
276         if check_convergence(T_next, T_current, epsilon, N)
277             break
278         end

```



```

279     T_current = T_next;
280 end
281 T_next_t = T_next;
282 end
283
284 function results = solver(r_min, r, K, h, delta_t, t_start, t_end, N, epsilon,
    max_iteration)
285     T_current_t = init_fuild(r_min, h, N);
286     t = t_start;
287     Ts(1, :) = T_current_t;
288
289     while t <= t_end
290         fprintf('N: %d || R: %g || epsilon: %g || delta_t: %g || t: %6.4f/%g\n', N,
            delta_t / h^2, epsilon, delta_t, t, t_end)
291         T_current_t = set_BC(T_current_t, t);
292         T_next_t = solve_for_specific_t_jacobi(T_current_t, r, K, h, delta_t, N, epsilon,
            max_iteration);
293         Ts(end+1, :) = T_next_t;
294
295         t = t + delta_t;
296         T_current_t = T_next_t;
297     end
298     results.Ts      = Ts;
299     results.N       = N;
300     results.r       = r;
301     results.h       = h;
302     results.delta_t = delta_t;
303     results.epsilon = epsilon;
304     results.R       = delta_t / h^2;
305     results.t_vec   = delta_t*(0:1:t_end/delta_t);
306 end
307
308 function y = interpolate(x, x_vec, y_vec)
309 % This function assume an ordered x_vec from low to high
310 if x > x_vec(end) || x < x_vec(1)
311     fprintf('value out of bounds\n');
312     return;
313 end
314
315 index_i = 0;
316 for i = 1:length(x_vec)
317     if x <= x_vec(i)
318         index_i = i;
319         break;
320     end
321 end
322
323 if x == x_vec(index_i)
324     y = y_vec(index_i);
325     return;
326 end
327
328 m = (y_vec(index_i+1) - y_vec(index_i)) / (x_vec(index_i+1) - x_vec(index_i));
329 b = y_vec(index_i) - x_vec(index_i) * m;
330
331 y = m * x + b;
332 end
333

```



```
334 function rms = RMS(x1, y1, x2, y2)
335 % This functions calculatlates the RMS value for 1 compared to 2
336     y2_len_y1 = [];
337     for i = 1:length(x1)
338         y2_len_y1(i) = interpolate(x1(i), x2, y2);
339     end
340
341     rms = sqrt(sum((y1-y2_len_y1).^2)/length(y1));
342 end
```

Listing 2: The main file