



הטכניון – מכון טכנולוגי לישראל
NUMERICAL METHODS IN AEROSPACE
ENGINEERING

HOMEWORK ASSIGNMENT x
סמסטר אביב תשפ"ה
SPRING SEMESTER 2025

GRADE	OUT OF	CHAPTER
	2	ABSTRACT
	2	CONTENTS, STYLE &C.
	4	PHYSICAL PROBLEM
	4	MATHEMATICAL MODEL
	26	NUMERICAL METHODS
	20	INFLUENCE OF NUMERICAL METHODS
	20	RESULTS
	2	SUMMARY & CONCLUSIONS
	20	COMPUTER PROGRAM
	100	TOTAL

Almog Dobrescu ID 214254252

April 24, 2025

Abstract

hello

Contents

1	The Physical Problem	1
2	The Mathematical Model	1
3	The Numerical Methods	1
3.1	Finite Difference Method	1
3.2	Shooting Method	2
4	Influence of The Numerical Methods	2
5	Results and Discussion	2
6	Summary and Conclusion	2
A	Listing of The Computer Program	3

List of Figures

Nomenclature

α	mass ratio between the liquid fuel and environment oxygen
β	ratio between the latent heat of the liquid fuel and between the chemical reaction heat of liquid vapor with oxygen
$.(n)$	value at time step n
Λ	empirical constant of droplet vaporation
h	size of each cell in the domain
i	cell index
m_d	mass fraction of liquid fuel in droplets
N	number of elements
s	temporery variable
T	temperature
T_u	environment temperature of the liquid fuel upstream
T_v	vaporization temperature of the liquid fuel
x	spatial coordinate



1 The Physical Problem

The physical problem at hand is the location of the flame front which depends on the point at which the fuel spray evaporates.

2 The Mathematical Model

The evaporation front of fuel spray can be described by solving the following equations:

$$\begin{aligned} \frac{d^2 T}{d\zeta^2} &= \Lambda e^T \left(T_v - T_u + \alpha\beta - \frac{dT}{d\zeta} \right) \\ m_d &= (\alpha\beta\Lambda e^T)^{-1} \frac{d^2 T}{d\zeta^2} \end{aligned} \quad (1)$$

The boundary condition of the problem:

$$\begin{aligned} \zeta \rightarrow -\infty : \quad & m_d \rightarrow 1 & T &\rightarrow \zeta \cdot (T_v - T_u) \\ \zeta \rightarrow +\infty : \quad & T \rightarrow \zeta \cdot (T_v - T_u + \alpha\beta) & m_d &\rightarrow 0 \end{aligned} \quad (2)$$

- According to the defenition of T : $T|_{\zeta=0} = 0$

3 The Numerical Methods

Eq.1 can be rewrite as:

$$\frac{d^2 T}{d\zeta^2} + \Lambda e^T \frac{dT}{d\zeta} - \Lambda e^T (T_v - T_u + \alpha\beta) = 0$$

In our case:

- ∞ is at around 30
- $\Lambda = 0.1$
- $T_v = 0.203$
- $T_u = 0.152$
- $\alpha\beta = 0.0234$

3.1 Finite Difference Method

Using central difference we can write the difference equations:

$$\begin{aligned} \frac{T_{i+1} - 2T_i + T_{i-1}}{h^2} + \Lambda e^{T_i} \frac{T_{i+1} - T_{i-1}}{2h} - \Lambda e^{T_i} (T_v - T_u + \alpha\beta) &= 0 & i = 1, 2, \dots, N \\ h &= \frac{\zeta|_{i=N+1} - \zeta|_{i=0}}{N+1-0} \\ &\vdots \end{aligned} \quad (3)$$



3.2 Shooting Method

Let's rewrite Eq.3 as a system of 2 ODE:

$$\begin{cases} \frac{dT}{d\zeta} = s & T|_{\zeta \rightarrow -\infty} \rightarrow \zeta \cdot (T_v - T_u) \\ \frac{ds}{d\zeta} = -\Lambda e^T s - \Lambda e^T (T_v - T_u + \alpha\beta) & T|_{\zeta \rightarrow +\infty} \rightarrow \zeta \cdot (T_v - T_u + \alpha\beta) \end{cases} \quad (4)$$

To solve the system of equations using the shooting method, we will guess $s_{(i=0)}^{(n)}$ and solve the system of equations using forward difference. Namely:

$$\begin{cases} T_{i+1}^{(n)} = s_i^{(n)} \cdot h + T_i^{(n)} \\ s_{i+1}^{(n)} = \left(-\Lambda e^{T_i^{(n)}} s_i^{(n)} - \Lambda e^{T_i^{(n)}} (T_v - T_u + \alpha\beta) \right) \cdot h + s_i^{(n)} \end{cases} \quad (5)$$

To correct the guess of $s_{(i=0)}^{(n)}$, let's define:

$$F_{(s_{(i=0)})} = T_{(i=N+1)}^{(n)} - T|_{\zeta \rightarrow +\infty} \quad (6)$$

- When $F = 0$, the guess of $s_{(i=0)}^{(n)}$ is correct

The next guess of $s_{(i=0)}^{(n+1)}$ we will use a numerical method to find the root of an equation. Namely:

$$s_{(i=0)}^{(n+1)} = s_{(i=0)}^{(n)} - F_{(s_{(i=0)})}^{(n)} \cdot \frac{s_{(i=0)}^{(n)} - s_{(i=0)}^{(n-1)}}{F_{(s_{(i=0)})}^{(n)} - F_{(s_{(i=0)})}^{(n-1)}} \quad (7)$$

4 Influence of The Numerical Methods

5 Results and Discussion

6 Summary and Conclusion



A Listing of The Computer Program