

Assignment 2. ST661 2019
Catherine Hurley
Due Monday October 28, 6pm

You should complete this assignment by filling in the answers into assign2.Rmd (on Moodle). You should knit the file to html and upload the html file to Moodle. The upload must be completed by time and date given above.

Marks will be deducted for unnecessary use of `for`.

1. Download the `cdc1.Rdata` file from moodle and load with

```
load("cdc1.Rdata")  
# or if you are working on the Rstudio server  
load("rstudio_files/ST661/cdc1.Rdata")
```

which gives a dataset called `cdc1`, a subset of the data used in class.

- (a) Change `exerany`, `smoke100` and `hlthplan` to be factors with suitable levels.
 - (b) Calculate the proportion of men that have health very good or better. Do the same for women. Who has the better health?
 - (c) Use `subset` to extract the smokers. For the smokers, calculate the proportion of men that have health very good or better. Do the same for women.
 - (d) Repeat (c) for non smokers.
 - (e) Based on your calculations of (c) and (d), compare the health of men and women, and the health of smokers and non smokers.
2. The definition of sample skewness of a set of data x_1, x_2, \dots, x_n is

$$\frac{m_3}{m_2^{3/2}}$$

where

$$m_3 = \sum_1^n (x_i - \bar{x})^3 / n$$

and

$$m_2 = \sum_1^n (x_i - \bar{x})^2 / n$$

- (a) Write function called `skew` to calculate the skewness of a numeric vector.
 - (b) Test your function on `mtcars$wt`.
 - (c) Test it also on a set of 50 numbers generated from the standard normal distribution.
 - (d) Using `replicate`, replicate the calculation of (c) 1000 times. Draw a histogram of the results and calculate the mean skewness value.
 - (e) Repeat (c) and (d), this time using a set 50 numbers generated from the exponential distribution with parameter 1.
 - (f) Compare the results of (d) and (e).
 - (g) Write code which calculates the skewness for every numeric variable in `cdc1`. Your code should construct a named vector of skewness values, with NA's for non numeric variables. Your code should not give any warnings.
3. (a) Make a plot of weight-`wt`desire versus age for the `cdc1` data. Use `pch=20` and a colour vector which assigns colours as follows:
weight > `wt`desire and age > 30 red
weight > `wt`desire and age <= 30 blue
weight <= `wt`desire and age > 30 orange
weight <= `wt`desire and age <= 30 cyan
Hint: use `ifelse`.

- (b) Recall in a previous lecture we used `boxplot.stats` to find indices of outliers. Write a function using `boxplot.stats` called `boxplot.out` that given a numeric vector returns “low” if there are low outliers, “high” if there are high outliers, “both” if there are high and low outliers, and “none” if there are no outliers.
- (c) For the `cdc1` data, make boxplots showing `weight-wtdesire` versus `genhlth`.
- (d) Using `boxplot.out` and `tapply`, construct a vector indicating the type of outlier (high, low, both or none) in each boxplot.
- (e) Construct a color vector that is “red” for boxplots that have high and low outliers, “blue” for just high outliers, “green” for just low outliers, and “yellow” otherwise. Use this vector to colour the boxplots.

4. Download `kobe.csv` from Moodle and load it in with

```
load("kobe.csv")
```

In the data frame `kobe`, every row records a shot taken by Kobe Bryant. If he hit the shot (made a basket), a hit, H, is recorded in the column named `basket`, otherwise a miss, M, is recorded.

- (a) Write a while loop to calculate how many throws are required to reach 3 hoops.
- (b) Write another while loop to calculate how many throws are required to reach 3 more hoops hoops.

Remaining questions are optional.

- (c) Construct a vector containing the number of throws required to get 3 hoops.
- (d) Construct a vector containing the number of throws required to get 3 hoops. Do not use `for`, `if`, `while`. Instead use `cumsum`, `diff`.
- (e) Define the length of a shooting streak to be the number of consecutive baskets made until a miss occurs. For example, in Game 1 Kobe had the following sequence of hits and misses from his nine shot attempts in the first quarter:

H M | M | H H M | M | M | M

Within the nine shot attempts, there are six streaks, which are separated by a | above. Their lengths are one, zero, two, zero, zero, zero (in order of occurrence). Write a function called `streaklen` that calculates the streak lengths in Kobe’s baskets. Tabulate the results and draw a barplot.