



Computer
Science

CSC380: Principles of Data Science

Probability Primer

Alon Efrat

Acknowledgement and thanks: Materials Built on previous product by

- Jason Pacheco,
- Kwang-Sung Jun,
- Chicheng Zhang
- Xincheng yu


- Please read
 - Ch. 6 ([WJ: Watkins, J., “An Introduction to the Science of Statistics: From Theory to Implementation”](#))

- Random Events and Probability
- Axioms of Probability
- Set Theory
- Law of Total Probability

Random Events and Probability

Suppose we roll two fair dice...

- ❑ What are the possible outcomes?
- ❑ What is the *probability* of rolling **even** numbers?
- ❑ What is the *probability* of having two numbers sum to 6?
- ❑ If one die rolls 1, then what is the probability of the second die also rolling 1?



...this is a random process.

How to mathematically formulate outcomes
and compute these probabilities?

Random Events and Probability


6

Probability of a random event

\approx

Simulate the random process n times, the fraction of times this event happens

- How large should n be?
- Simulation results vary from trails?



Background: Numpy in Python

7

Numpy: numerical computing package

```
import numpy as np
np.random.randint(1,1+6,size=10)
=> array([5, 4, 1, 1, 1, 5, 5, 2, 4, 6])
```

```
randint(low,high,size)
: generate `size` random numbers
in {low, low+1, ..., high-1}
```

Numpy array

- Replaces python's list in numpy.
- More numerical functionality
- It's a 'vector' in mathematics.

```
a=np.array([1,2]); b=np.array([4,5])
a+b
=> np.array([5,7]) // elementwise addition
np.dot(a,b)
=> 14           // dot product
```

Random Events and Probability

8

Consider: What is the probability of having two numbers sum to 6?

```
import numpy as np
for n in [10,100,1_000,10_000,100_000]:
    res_dice1 = np.random.randint(1,6+1,size=n)
    res_dice2 = np.random.randint(1,6+1,size=n)
    res = [(res_dice1[i], res_dice2[i]) for i in range(len(res_dice1))]

    cnt = len(list(filter(lambda x: x[0] + x[1] == 6, res)))
    print("n=%6d, result: %.4f " % (n, cnt/n))
```

```
n=    10, result: 0.1000
n=   100, result: 0.1200
n=  1000, result: 0.1350
n= 10000, result: 0.1365
n=100000, result: 0.1388
n=1000000, result: 0.1385
```

```
n=    10, result: 0.1000
n=   100, result: 0.1900
n=  1000, result: 0.1540
n= 10000, result: 0.1366
n=100000, result: 0.1371
n=1000000, result: 0.1394
```

every time you run, you
get a different result

however, the number
seems to converge to
0.138-0.139

There seems to be a precise value that it will converge to.. what is it?


Each outcome is equally likely.
by the **independence**
(will learn this concept later)
=> $1/36$

answer:
 $(1/36) * 5 = 0.13888..$

Random Events and Probability

10

- **Theoretical probability** describes how likely an event is going to occur based on math.
- **Experimental probability** describes how frequently an event actually occurred in an experiment.




- **Probability** is a real-world phenomenon.
- But under what mathematical framework can we formulate **probability** so we can solve practical problems?
 - e.g., weather prediction, predicting the election outcome
- **Disclaimer**: not all mathematics correspond to real-world phenomenon (e.g., Banach–Tarski paradox). Fortunately, we will not talk about this in our lecture 😊

Random Events and Probability

12

Consider: What is the probability of having two numbers sum to 6?



Some examples of events...

- Both even numbers

Q: how many such pairs?

9

$$E^{\text{even}} = \{(2, 2), (2, 4), \dots, (6, 4), (6, 6)\}$$

- The sum of both dice is even,

$$E^{\text{sum even}} = \{(1, 1), (1, 3), (1, 5), \dots, (2, 2), (2, 4), \dots\}$$

- The sum is greater than 12,

$$E^{\text{sum} > 12} = \emptyset$$

We can talk about
impossible outcomes

- The product is even (How many events????)

Axioms of Probability

Random Events and Probability

15

But, what is probability, really?

(e.g., can explain the probability of seeing an event when throwing two dice)

Mathematicians have found a set of conditions that 'makes sense'.


- Probability is a map P on a set Ω .
 \Rightarrow i.e., takes in an event, spits out a real value
- P must map events to a real value in interval $[0,1]$.
- P is a (valid) **probability distribution** if it satisfies the following **axioms of probability**,

1. For any event E , $P(E) \geq 0$

2. $P(\Omega) = 1$

3. For any sequence of disjoint events E_1, E_2, E_3, \dots

$$P\left(\bigcup_{i \geq 1} E_i\right) = \sum_{i \geq 1} P(E_i)$$



disjoint: intersection is empty

Random Events and Probability

16

- Many properties follows (i.e., can be proved mathematically)

$$\mathbb{P}(\emptyset) = 0$$

$$A \subset B \implies \mathbb{P}(A) \leq \mathbb{P}(B) \quad \text{E.g., throw a die. } A = \text{getting 1, } B = \text{getting an odd number}$$


$$0 \leq \mathbb{P}(A) \leq 1$$

$$\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

$$A \cap B = \emptyset \implies \mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B). \quad \text{E.g., } A = \text{getting 1, } B = \text{getting 3 or 5}$$

A^c

the complement of A. All outcomes not in A



(I recommend that you maintain your own version of cheat sheet!)

Random Events and Probability

17


Special case

Assume each outcome is equally likely, and sample space is finite, then the probability of event is:

$$P(E) = \frac{|E|}{|\Omega|}$$

Number of elements in event set

Number of possible outcomes (36)



This is called uniform probability distribution

Q: What axiom we are using?
=> Axiom 3

(Fair) Dice Example: Probability that we roll even numbers,


$$P((2, 2) \cup (2, 4) \cup \dots \cup (6, 6)) = P((2, 2)) + P((2, 4)) + \dots + P((6, 6))$$
$$= \frac{1}{36} + \frac{1}{36} + \dots + \frac{1}{36} = \frac{9}{36}$$

9 Possible outcomes, each with equal probability of occurring

Random Events and Probability

18

Consider: What is the probability of having two numbers sum to 6?



Each outcome is equally likely.
by the **independence**
(will learn this concept later)
=> 1/36

of outcomes that sum to 6:
=> 5

answer:
(1/36) * 5 = 0.13888..

$$P(E) = \frac{|E|}{|\Omega|}$$

Set Theory

Set Theory

20

Two dice example: Suppose

E_1 : First die equals 1

E_2 : Second die equals 1

$$E_1 = \{(1, 1), (1, 2), \dots, (1, 6)\}$$

$$E_2 = \{(1, 1), (2, 1), \dots, (6, 1)\}$$

Operators on events:

Operation	Value	Interpretation
$E_1 \cup E_2$	$\{(1, 1), (1, 2), \dots, (1, 6), (2, 1), \dots, (6, 1)\}$	Any die rolls 1
$E_1 \cap E_2$	$\{(1, 1)\}$	Both dice roll 1
$E_1 \setminus E_2$	$\{(1, 2), (1, 3), (1, 4), (1, 5), (1, 6)\}$	Only the first die rolls 1
$\overline{E_1 \cup E_2}$	$\{(2, 2), (2, 3), \dots, (2, 6), (3, 2), \dots, (6, 6)\}$	No die rolls 1


$$E_1 \setminus E_2 = E_1 - E_2 := E_1 \cap E_2^c$$

$$\overline{E_1 \cup E_2} = (E_1 \cup E_2)^c$$

Set Theory

21

Can interpret these operations as a Venn diagram...



Set Theory

More results

- $\neg(\bigcup_n A_n) = \bigcap_n \neg A_n$, $\neg(\bigcap_n A_n) = \bigcup_n \neg A_n$ DEMORGAN

Special case: $\neg(A \cup B) = \neg A \cap \neg B$

Notation: $\neg A := A^c$


- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ and $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$.

$$A \cap (\bigcup_i B_i) = \bigcup_i (A \cap B_i), \quad A \cup (\bigcap_i B_i) = \bigcap_i (A \cup B_i)$$

- $B = \Omega \cap B = (A \cup \neg A) \cap B = (A \cap B) \cup (\neg A \cap B)$

// by distributive law


TIP: always draw a picture to visualize these identities!



Lemma: (inclusion-exclusion rule) For any two events E_1 and E_2 ,

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2)$$

Graphical Proof:



Subtract from both sides

Alternative Proof

24

Lemma: For any two events E_1 and E_2 ,

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2)$$

Alternative proof:



$$\begin{aligned} A &= E_1 - (E_1 \cap E_2) \\ B &= E_1 \cap E_2 \\ C &= E_2 - (E_1 \cap E_2). \end{aligned}$$

$$\begin{aligned} P(E_1 \cup E_2) &= P(A \cup B \cup C) \\ &= P(A) + P(B) + P(C) \\ &= P(A) + P(B) + P(B) + P(C) - P(B) \\ &= P(A \cup B) + P(B \cup C) - P(B) \end{aligned}$$

(by axiom 3)

(by axiom 3)

Exercise:

Quiz candidate

- Consider rolling two fair dice
- E_1 : two dice sum to 6
- E_2 : second die is even
- Compute the numerical value of $P(E_1 \cup E_2)$. Hint: Use inclusion-exclusion rule.

$$P(E_1) = 5/36$$

$$P(E_2) = 18/36$$

$$P(E_1 \cap E_2) = 2/36$$


answer: 21/36

Law of Total Probability

Random Events and Probability

27

[Def] The set of events $\{B_i\}_{i=1}^n$ **partitions** outcome space $C \Leftrightarrow \cup_i B_i = C$ and B_1, B_2, \dots are disjoint.




Q: Why is this true?

A: [Axiom 3!](#)

$$A = A \cap \Omega = A \cap (\cup_i B_i) = \cup_i (A \cap B_i)$$

Now, $\{A \cap B_i\}_{i=1}^n$ partitions A



Random Events and Probability

28

Law of total probability: Let A be an event. For any events B_1, B_2, \dots that partitions Ω , we have

$$P(A) = \sum_i P(A \cap B_i)$$

Example Roll two fair dice. Let X be the outcome of the first die. Let Y be the sum of both dice. What is the probability that both dice sum to 6 (i.e., $Y=6$)?

quiz candidate

$$p(Y = 6) = \sum_{x=1}^6 p(Y = 6, X = x)$$

$$P(A, B) := P(A \cap B)$$

$$= p(Y = 6, X = 1) + p(Y = 6, X = 2) + \dots + p(Y = 6, X = 6)$$

$$= \frac{1}{36} + \frac{1}{36} + \frac{1}{36} + \frac{1}{36} + \frac{1}{36} + 0 = \frac{5}{36}$$

Summary So Far

29

- Most of the rules we learned is basically set theory + axiom 3

So, here is a generic workflow for computing $P(A)$.

1. Use set theory and slice and dice A into a manageable partition of A where $P(\text{each piece of partition})$ is easy to compute.
2. Apply Axiom 3.

Conditional Probability

30


- Two fair dice example:

- Suppose I roll two dice secretly and tell you that one of the dice is 2. C
- In this situation, find the probability of two dice summing to 6. E

```
import numpy as np
for n in [10,100,1000,10_000,100_000, 1_000_000]:
    res_dice1 = np.random.randint(6,size=n) + 1
    res_dice2 = np.random.randint(6,size=n) + 1
    res = [(res_dice1[i], res_dice2[i]) for i in range(len(res_dice1))]
```

```
conditioned = list(filter(lambda x: x[0] == 2 or x[1] == 2, res))
n_eff = len(conditioned)
```

```
cnt = len(list(filter(lambda x: x[0] + x[1] == 6, conditioned)))
print("n=%9d, n_eff=%9d, result: %.4f " % (n, n_eff, cnt/n_eff))
```



compare:
without conditioning,
it was 0.138..

n=	10, n_eff=	4, result: 0.0000
n=	100, n_eff=	32, result: 0.2500
n=	1000, n_eff=	300, result: 0.1733
n=	10000, n_eff=	3002, result: 0.1742
n=	100000, n_eff=	30590, result: 0.1823
n=	1000000, n_eff=	305616, result: 0.1818

n=	10, n_eff=	3, result: 0.3333
n=	100, n_eff=	32, result: 0.0625
n=	1000, n_eff=	343, result: 0.2245
n=	10000, n_eff=	3062, result: 0.1897
n=	100000, n_eff=	30651, result: 0.1811
n=	1000000, n_eff=	305580, result: 0.1808