

University of Utah

Spring 2025

# MATH 3210-001

## Midterm 1 Questions

Instructor: Alp Uzman

February 7, 2025, 9:40 AM - 10:30 AM

Surname:

First Name:

uNID:

KEY

**Before turning the page  
make sure to read and  
sign the exam policy  
document, distributed  
separately.**

1. [20 points] Prove or disprove the following statement: Let

$f : \mathbb{R} \rightarrow \mathbb{R}$  be a function,  $A, B \subseteq \mathbb{R}$  be two subsets. Then

$$f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B).$$

$$(\subseteq) A \cap B \subseteq A, B \Rightarrow f^{-1}(A \cap B) \subseteq f^{-1}(A), f^{-1}(B)$$

$$\Rightarrow f^{-1}(A \cap B) \subseteq f^{-1}(A) \cap f^{-1}(B) \quad \square$$

$$(\supseteq) \text{ Let } x \in f^{-1}(A) \cap f^{-1}(B). \text{ Then } x \in f^{-1}(A) \text{ AND}$$

$$x \in f^{-1}(B) \Rightarrow f(x) \in A \text{ AND } f(x) \in B$$

$$\Rightarrow f(x) \in A \cap B \Rightarrow x \in f^{-1}(A \cap B)$$

$$\Rightarrow f^{-1}(A) \cap f^{-1}(B) \subseteq f^{-1}(A \cap B). \quad \square$$

**Extra Space** If you do not want the work on this page to be graded, cross out this page before turning in your booklet. Otherwise, ensure your work is properly numbered and organized.

2. [60 points] Consider the set  $S$  that consists of all positive rational numbers  $x$  with the property that  $x^2 \leq 2$ .

$$S = \{x \in \mathbb{Q} \mid 0 < x, x^2 \leq 2\}.$$

- (a) [20 points] Prove or disprove the following statement:

The supremum of  $S$  is  $\sqrt{2}$ .

$$x \in S \Rightarrow x^2 \leq 2 \Rightarrow x \leq \sqrt{2}$$

$\Rightarrow \sqrt{2}$  is an upper bound for  $S$

$$(1^2 = 1 \leq 2 \Rightarrow 1 \in S \Rightarrow S \neq \emptyset)$$

$$\Rightarrow \inf(S) \leq \sqrt{2}.$$

say  $\sup(S) \leq \sqrt{2}$ . ~~then  $0 < \sqrt{2} - \sup(S)$~~

Then by the density of rationals,

$$\exists q \in \mathbb{Q}: \text{ ~~$0 < q < \sqrt{2} - \sup(S) < \sqrt{2}$~~ }$$

$$\text{ ~~$q \in S$~~ } \quad \left[ \begin{array}{l} \sup(S) \leq \max\{\sup(S), 0\} \\ < q < \sqrt{2} \end{array} \right]$$

- (b) [20 points] Prove or disprove the following statement:

The infimum of  $S$  is  $\sqrt{2}$ .

$$0 < 1, \quad 1^2 = 1 \leq 2$$

$$\Rightarrow 1 \in S.$$

$$\Rightarrow \inf(S) \leq 1 < \sqrt{2}$$

$$\Rightarrow \inf(S) \neq \sqrt{2}. \quad \square$$

$$\Rightarrow 0 < q, \quad q^2 < 2$$

$$\Rightarrow q \in S, \text{ a contradiction!}$$

$$\text{Hence } \sup(S) = \sqrt{2}. \quad \square$$

(c) [10 points] Prove or disprove the following statement:

The set of all upper bounds of  $S$  is  $[\sqrt{2}, \infty[$ .

after part (a),  $\sup(S) = \sqrt{2}$ .

By definition,  $\sup(S)$  is the minimum of all upper bounds for  $S$ .

$\Rightarrow \forall U > \sup(S), \forall x \in S: x \leq U$

$\Rightarrow [\sqrt{2}, \infty[$  indeed is the set of all upper bounds for  $S$ .  $\square$

(d) [10 points] Prove or disprove the following statement:

The smallest closed interval containing  $S$  is  $[0, \sqrt{2}]$ .

Claim:  $\inf(S) = 0$ .

Pf of Claim: By def,  $\forall x \in S, 0 \leq x$

$\Rightarrow 0 \leq \inf(S)$ . If  $0 < \inf(S)$ , by density of rationals,  $\exists q \in \mathbb{Q}$ ,

$0 < q < \min\{\inf(S), \sqrt{2}\}$

$\Rightarrow q \in S \Rightarrow q < \inf(S)$ , a contradiction.

We also know from part (a) that  $\sup(S) = \sqrt{2}$

$\hookrightarrow$



**Extra Space** If you do not want the work on this page to be graded, cross out this page before turning in your booklet. Otherwise, ensure your work is properly numbered and organized.

(2.(d) continued)

$$\text{Hence } \forall x \in S : 0 = \inf(S) \leq x \leq \sup(S) = \sqrt{2} \\ \Rightarrow S \subseteq [0, \sqrt{2}].$$

Let  $[\lambda, p]$ ,  $-\infty < \lambda \leq p < \infty$  be such that  $[\lambda, p] \supseteq S \Rightarrow \lambda$  is a lower bound for  $S$  and  $p$  is an upper bound for  $S$ . Then by def,

$$\lambda \leq \inf(S), \quad \sup(S) \leq p$$

$$\Leftrightarrow [\inf(S), \sup(S)] = [0, \sqrt{2}] \subseteq [\lambda, p] \\ \square.$$

3. [16 points] Prove or disprove the following statement: Let  $x_n$  be a sequence of real numbers that converges to zero. Then for any bounded sequence  $b_n$  of real numbers,

Have:

$$\lim_{n \rightarrow \infty} x_n b_n = 0.$$

Since  $b_n$  is bounded,

$$\exists M \in \mathbb{R}_{>0}, \forall n \in \mathbb{Z}_{\geq 0}: |b_n| \leq M.$$

Since  $x_n \rightarrow 0$ ,

$$\forall \gamma \in \mathbb{R}_{>0}, \exists N = N(\gamma) \in \mathbb{Z}_{\geq 0}, \forall n \in \mathbb{Z}_{\geq 0}:$$

"eta"  $n \geq N(\gamma) \Rightarrow |x_n| < \gamma.$

Want:  $\forall \varepsilon \in \mathbb{R}_{>0}, \exists N' = N'(\varepsilon) \in \mathbb{Z}_{\geq 0}, \forall n \in \mathbb{Z}_{\geq 0}:$   
 $n \geq N'(\varepsilon) \Rightarrow |x_n b_n| < \varepsilon.$

Let  $\varepsilon \in \mathbb{R}_{>0}$ . Put  $\gamma = \frac{\varepsilon}{2M} > 0$ . Then  $\forall n$ :

$$n \geq N(\gamma) = N\left(\frac{\varepsilon}{2M}\right) \Rightarrow |x_n| < \frac{\varepsilon}{2M}$$

$$\Rightarrow |x_n b_n| = |x_n| |b_n| \leq \frac{\varepsilon}{2M} M = \frac{\varepsilon}{2} < \varepsilon.$$

Thus for  $N'(\varepsilon) = N\left(\frac{\varepsilon}{2M}\right)$ , we ~~have~~ obtain what we want.  $\square$ .

**Extra Space** If you do not want the work on this page to be graded, cross out this page before turning in your booklet. Otherwise, ensure your work is properly numbered and organized.



4. [4 points] Let  $x_\bullet : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}$  be a sequence of real numbers and  $x_*$  be a real number.

- (a) [1 point] **Prove or disprove the following statement:** If  $x_\bullet$  converges to  $x_*$ , then any subsequence of  $x_\bullet$  converges to  $x_*$ .

Let  $x'_\bullet \leq x_\bullet$ . Then  $\exists$  strictly increasing  $j$ ,  $\forall n: x'_n = x_{j(n)}$ . Note:  $\forall n: n \leq j(n)$ .

(\*)

Have:  $x_\bullet \rightarrow x_*$ , i.e.,

$$\left[ \forall \gamma \in \mathbb{R}_{>0}, \exists N = N(\gamma) \in \mathbb{Z}_{\geq 0}, \forall n \in \mathbb{Z}_{\geq 0}: \right. \\ \left. n \geq N = N(\gamma) \Rightarrow |x_n - x_*| < \gamma. \right.$$

Want:  $x'_\bullet \rightarrow x_*$ , i.e.,

$$\left[ \forall \varepsilon \in \mathbb{R}_{>0}, \exists N' = N'(\varepsilon) \in \mathbb{Z}_{\geq 0}, \forall n \in \mathbb{Z}_{\geq 0}: \right. \\ \left. n \geq N' = N'(\varepsilon) \Rightarrow |x'_{j(n)} - x_*| < \varepsilon. \right.$$

Let  $\varepsilon \in \mathbb{R}_{>0}$ , put  $\gamma = \varepsilon$ . Then  $\exists N = N(\varepsilon)$ ,  $\forall n: n \geq N(\varepsilon) \Rightarrow |x_n - x_*| < \varepsilon$ .

But by (\*), if  $n \geq N(\varepsilon)$ , we also have  $j(n) \geq n \geq N(\varepsilon)$ , hence defining  $N'(\varepsilon) = N(\varepsilon)$  we are done.  $\square$ .

(b) [1 point] Prove or disprove the following statement: If

$$\forall \varepsilon \in \mathbb{R}_{>0}, \forall N \in \mathbb{Z}_{\geq 0}, \exists n \in \mathbb{Z}_{\geq N} : |x_n - x_*| < \varepsilon,$$

then  $x_n$  converges to  $x_*$ .

Consider  $x : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}, n \mapsto (-1)^n$ ,

and put  $x_* = 1$ .

Let  $\varepsilon \in \mathbb{R}_{>0}, N \in \mathbb{Z}_{\geq 0}$ . Then for  
 $n = j(N) = 2N \geq N$ , and

$$|x_n - x_*| = |x_{2N} - 1| = 0 < \varepsilon.$$

Thus  $x$  and  $x_*$  satisfy the hypothesis.

However  $x$  does not converge to  $x_*$

(proved in class).  $\square$ .

(c) [1 point] Prove or disprove the following statement: If

$$\forall \varepsilon \in \mathbb{R}_{>0}, \forall N \in \mathbb{Z}_{\geq 0}, \exists n \in \mathbb{Z}_{\geq N} : |x_n - x_*| < \varepsilon,$$

then some subsequence of  $x_\bullet$  converges to  $x_*$ .

$$\varepsilon = 1, N = 1 \Rightarrow \exists n_1 : n_1 \geq 1, |x_{n_1} - x_*| < 1.$$

$$\varepsilon = \frac{1}{2}, N = n_1 + 1 \Rightarrow \exists n_2 > n_1, |x_{n_2} - x_*| < \frac{1}{2}.$$

$$\varepsilon = \frac{1}{3}, N = n_2 + 1 \Rightarrow \exists n_3 > n_2 : |x_{n_3} - x_*| < \frac{1}{3}.$$

$$\vdots$$

$$\forall k \in \mathbb{Z}_{\geq 1}, \text{ for } \varepsilon = \frac{1}{k}, N = n_{k-1} + 1, \exists n_k > n_{k-1}:$$

(\*)  $|x_{n_k} - x_*| < \frac{1}{k}$  by the Principle of Mathematical Induction.

$$1 \leq n_1 < n_2 < \dots < n_k < \dots$$

$\Rightarrow x'_\bullet$  defined by  $k \mapsto x_{n_k}$  is a subsequence of  $x_\bullet$ .

Let  $\gamma \in \mathbb{R}_{>0}$ . Then  $\exists K \in \mathbb{Z}_{\geq 1} : 0 < \frac{1}{K} < \gamma$  by the Archimedean Principle.  $\forall k : k \geq K$   
 $\Rightarrow 0 < \frac{1}{k} \leq \frac{1}{K} < \gamma \stackrel{(*)}{\Rightarrow} |x'_k - x_*| \leq \frac{1}{k} < \gamma \Rightarrow x'_\bullet \rightarrow x_*$   
 T.B.

(Or you can use the squeeze theorem for sequences to conclude that  $x'_\bullet \rightarrow x_*$ )



(d) [1 point] Prove or disprove the following statement: If some subsequence of  $x_n$  converges to  $x_*$ , then

$$\forall \varepsilon \in \mathbb{R}_{>0}, \forall N \in \mathbb{Z}_{\geq 0}, \exists n \in \mathbb{Z}_{\geq N} : |x_n - x_*| < \varepsilon.$$

Let  $x'_n \leq x_n$  be such that  $x'_n \rightarrow x_*$ .

Then  $\exists$  strictly increasing  $j$ ,  $\forall n$ :

$$x'_n = x_{j(n)}.$$

Let  $\varepsilon \in \mathbb{R}_{>0}$ ,  $N \in \mathbb{Z}_{\geq 0}$ . Since

$$x'_n \rightarrow x_*,$$

$$\exists N' = N'(\varepsilon) \in \mathbb{Z}_{\geq 0}, \forall n \in \mathbb{Z}_{\geq 0}:$$

$$n \geq N' = N'(\varepsilon) \Rightarrow |x_{j(n)} - x_*| < \varepsilon. (*)$$

Want:  $\boxed{\exists n = n(\varepsilon, N) : n \geq N \text{ and } |x_n - x_*| < \varepsilon.}$

Put  $n(\varepsilon, N) = j(\max\{N, N'\})$ .

Then  $n(\varepsilon, N) \geq N, N'$  (as  $j$  is strictly increasing) and by  $(*)$ ,  $|x_{n(\varepsilon, N)} - x_*| < \varepsilon$ .  $\square$

**Extra Space** If you do not want the work on this page to be graded, cross out this page before turning in your booklet. Otherwise, ensure your work is properly numbered and organized.



**Extra Space** If you do not want the work on this page to be graded, cross out this page before turning in your booklet. Otherwise, ensure your work is properly numbered and organized.