

Exact 6-cut rigidity and small-order superconnectivity for the 6-regular case of Dirac’s $k = 4$ problem

Alper Ferudun*

June 2026

Abstract

Dirac asked in 1970 whether for every $k \geq 4$ there is a k -vertex-critical graph without critical edges; Jensen settled all $k \geq 5$, and only $k = 4$ remains open. Following Skottová and Steiner, call a graph G a $(4, 1)$ -graph if $\chi(G) = 4$, $\chi(G - v) = 3$ for every vertex v , and $\chi(G - e) = 4$ for every edge e ; they proved $\delta(G) \geq 6$ and $\lambda(G) \geq 6$ for every $(4, 1)$ -graph and asked whether a 6-regular $(4, 1)$ -graph exists. We prove two results about this 6-regular case. **Theorem A** (computational, double-verified): there is no 6-regular 4-vertex-critical graph on $n \leq 12$ or $n = 14$ vertices; on $n = 13$ there is exactly one, and its 13 critical edges form a Hamilton cycle. Consequently any 6-regular $(4, 1)$ -graph has at least 15 vertices. **Theorem B**: in a 6-regular $(4, 1)$ -graph, every 6-edge-cut is either the edge star of a vertex or has both shores of size at least 15; consequently every 6-regular $(4, 1)$ -graph on at most 29 vertices is super-6-edge-connected. The proof of Theorem B combines an exact classification of the 3×3 cut matrices of 6-edge-cuts in $(4, 1)$ -graphs (exactly 21 matrices, five types up to row/column permutations) with four local necessary conditions on a hypothetical shore, the strongest being a *boundary-shortfall lemma* derived from vertex-deleted colourings; a finite, exhaustively enumerated candidate space then collapses. The unique near-miss is $K_{3,3,3}$ minus a rainbow perfect matching of size 3, which survives all counting filters and is excluded by the boundary-shortfall lemma alone. Key lemmas are machine-checked in Lean 4/Mathlib, and all enumerations were performed by two independent implementations.

1 Introduction

A graph G is k -vertex-critical if $\chi(G) = k$ and $\chi(G - v) = k - 1$ for every vertex v . An edge set $R \subseteq E(G)$ is *critical* if $\chi(G - R) < \chi(G)$. Dirac (see Erdős [2]) asked in 1970 whether for every $k \geq 4$ there exists a k -vertex-critical graph with no critical edge. Jensen [3] resolved all cases $k \geq 5$; the case $k = 4$ remains open, the only prior partial result being that of Jensen and Siggers [4], who constructed 4-vertex-critical graphs in which fewer than an ε -fraction of the edges are critical. The problem is #944 in Bloom’s database of Erdős problems [1].

Following Skottová and Steiner [5] we call a graph G a $(4, 1)$ -graph (for short, a *target*) if

$$\chi(G) = 4, \quad \chi(G - v) = 3 \quad \forall v \in V(G), \quad \chi(G - e) = 4 \quad \forall e \in E(G).$$

Skottová and Steiner proved that every $(4, 1)$ -graph satisfies $\delta(G) \geq 6$ and $\lambda(G) \geq 6$ (their Proposition 5.1) and posed as an explicit subproblem (their Problem 5.2) whether a 6-regular

*alper@mercurycodelab.com. Code, certificates and machine-checked Lean files accompany this note as ancillary files; development history at github.com/AlperTheKing.

$(4,1)$ -graph exists — the extremal, sparsest possible case. This note contributes two verified results on that subproblem.

Theorem 1.1 (A). *There is no 6-regular 4-vertex-critical graph on $n \leq 12$ or on $n = 14$ vertices. On $n = 13$ vertices there is exactly one 6-regular 4-vertex-critical graph G_{13} (given in graph6 form in Section 6); its critical edges number 13 and form a Hamilton cycle. Consequently every 6-regular $(4,1)$ -graph has at least 15 vertices.*

Theorem 1.2 (B). *In every 6-regular $(4,1)$ -graph, every 6-edge-cut is either the edge star of a single vertex or has both shores of size at least 15. Consequently every 6-regular $(4,1)$ -graph on at most 29 vertices is super-6-edge-connected.*

Theorem B is obtained by an *exact finite obstruction method*: the equality case of the Skottová–Steiner cut bound forces every 6-edge-cut of a target to carry one of exactly 21 cut matrices (five types up to row and column permutations, Theorem 3.2); this, together with three further local necessary conditions — the strongest being the boundary-shortfall lemma (Lemma 4.2) — confines a hypothetical small shore to a finite, explicitly enumerable family which then collapses entirely. We consider the method itself, and the structural near-miss it uncovers ($K_{3,3,3}$ minus a rainbow 3-matching, Section 5), to be of independent interest.

We are careful to separate known from new. The bounds $\delta \geq 6$, $\lambda \geq 6$ and the cut-averaging argument are due to Skottová and Steiner [5]; the exclusion of comparable nonadjacent vertices in vertex-critical graphs is folklore. The equality-case classification (Theorem 3.2), the boundary-shortfall lemma, the shore-exclusion theorems, and the computational closure of $n = 13, 14$ appear to be new. None of this comes close to resolving Problem 5.2: a target could still be a super-6-edge-connected 6-regular graph on $n \geq 15$ vertices, and our results do not constrain that core case.

All computations were performed by two independent implementations (C++ and Python) with exact agreement, with generator-side completeness guards, and the central counting lemmas are machine-checked in Lean 4/Mathlib; see Section 6 and the ancillary files.

2 Local colouring lemmas

Throughout, colourings are proper maps to $\{1, 2, 3\}$ (not counted up to colour permutation). For $v \in V(G)$ and a 3-colouring φ of $G - v$ write $a_i(v, \varphi) = |\{u \in N(v) : \varphi(u) = i\}|$.

Lemma 2.1 (singleton lemma). *Let G be 4-vertex-critical, $v \in V(G)$, and φ a 3-colouring of $G - v$. Then $a_i(v, \varphi) \geq 1$ for every colour i ; and if $a_i(v, \varphi) = 1$, say with unique colour- i neighbour u , then the edge vu is critical: $\chi(G - vu) = 3$.*

Proof. If $a_i = 0$, extend φ by $\varphi(v) = i$; this 3-colours G , contradicting $\chi(G) = 4$. If $a_i = 1$ with witness u , keep φ on $G - v$ and set $\varphi(v) = i$ in $G - vu$: the only monochromatic edge would be vu , which has been deleted. $\square$

Corollary 2.2. *In a target, $a_i(v, \varphi) \geq 2$ for all v, φ, i ; hence $\delta(G) \geq 6$, and in a 6-regular target every neighbourhood splits exactly $2 + 2 + 2$ under every vertex-deleted colouring. (This is the local form of [5, Prop. 5.1].)*

Corollary 2.3. *Every 4-vertex-critical graph G has at least $\frac{1}{2} \sum_v \max(0, 6 - d(v))$ critical edges.*

Proof. Fix for each v one colouring φ_v of $G - v$. The three positive integers $a_i(v, \varphi_v)$ sum to $d(v)$, so at least $\max(0, 6 - d(v))$ of them equal 1; each singleton yields a critical edge at v by Lemma 2.1, and each critical edge is counted at most twice. $\square$

Lemma 2.4 (folklore). *A k -vertex-critical graph contains no two nonadjacent vertices u, v with $N(u) \subseteq N(v)$.*

Proof. Take a $(k - 1)$ -colouring φ of $G - u$ and set $\varphi(u) = \varphi(v)$: every neighbour of u is a neighbour of v , so no conflict arises and $\chi(G) \leq k - 1$. $\square$

We will also use the following observation, which gives a very effective computational prefilter.

Lemma 2.5. *In a 4-vertex-critical graph G on $n \geq 8$ vertices, every induced neighbourhood $G[N(v)]$ is bipartite.*

Proof. An odd cycle $C \subseteq N(v)$ together with v forms a 4-chromatic wheel on at most 7 vertices; any vertex u outside it gives $\chi(G - u) \geq 4$. $\square$

3 Exact rigidity of 6-edge-cuts

Let G be a target and $V(G) = A \sqcup B$ a nontrivial partition with cut $F = E(A, B)$. Both $G[A]$ and $G[B]$ are 3-colourable (each is contained in some $G - x$). For 3-colourings α of $G[A]$ and β of $G[B]$ define the *cut matrix* $M = (m_{ij})$ by $m_{ij} = |\{ab \in F : \alpha(a) = i, \beta(b) = j\}|$, and for $\pi \in S_3$ the diagonal sum $D_\pi(M) = \sum_i m_{i\pi(i)}$, the number of monochromatic cut edges after permuting the colours of B by π .

Lemma 3.1. *In a target: $D_\pi(M) = 0$ is impossible; if $D_\pi(M) = 1$ then the unique monochromatic cut edge is critical. Hence $D_\pi(M) \geq 2$ for all π , and since $\frac{1}{6} \sum_\pi D_\pi(M) = |F|/3$, every cut satisfies $|F| \geq 6$ ([5, Prop. 5.1]), and for $|F| = 6$,*

$$D_\pi(M) = 2 \quad \text{for all } \pi \in S_3 \text{ and all pairs } (\alpha, \beta).$$

Proof. $D_\pi = 0$ gives a 3-colouring of G ; $D_\pi = 1$ gives a 3-colouring of $G - e$ for the unique monochromatic cut edge e . Each cut edge lies on exactly two of the six permutation diagonals, giving the average $|F|/3$; for $|F| = 6$ the average equals the minimum, forcing equality everywhere. $\square$

Theorem 3.2 (21-matrix classification). *Let M be a 3×3 matrix with nonnegative integer entries, total sum 6, and $D_\pi(M) = 2$ for all $\pi \in S_3$. Then*

$$m_{ij} = \frac{R_i + C_j - 2}{3}$$

where R_i, C_j are the row and column sums; all R_i are congruent mod 3, likewise all C_j ; and M is, up to row and column permutations, one of the five types

$$\begin{pmatrix} 2 & 2 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 0 & 0 \\ 2 & 0 & 0 \\ 2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix},$$

with orbit sizes 3, 3, 9, 3, 3: exactly 21 matrices in all. In particular the row-sum vector is a permutation of $(6, 0, 0)$, $(3, 3, 0)$, $(4, 1, 1)$ or $(2, 2, 2)$.

Proof. Comparing two permutations that differ by a transposition and cancelling the common entry gives $m_{rc} + m_{sd} = m_{rd} + m_{sc}$ for every 2×2 submatrix; hence $m_{pq} = m_{pj} + m_{iq} - m_{ij}$ for all p, q, i, j , and summing over all (p, q) yields $6 = 3R_i + 3C_j - 9m_{ij}$, i.e. the displayed formula. Integrality forces $R_i + C_j \equiv 2 \pmod{3}$ for all i, j , whence the congruence claims; the nonnegative solutions of $\sum R_i = 6$ with all R_i congruent mod 3 are the four listed vectors, and the case analysis over (R, C) produces exactly the five displayed types. The count 21 is also machine-checked (Lean, by exhaustive enumeration of the 3003 weak compositions of 6 into 9 parts together with a completeness lemma for the enumeration; see Section 6). $\square$

Since the row sums of $M(\alpha, \beta)$ do not depend on β , we obtain:

Corollary 3.3. *For every 3-colouring α of a 6-cut shore $G[A]$ in a target, the vector $(\sum_{\alpha(v)=i} b(v))_{i=1,2,3}$, where $b(v)$ is the number of cut edges at v , is a permutation of $(6, 0, 0)$, $(3, 3, 0)$, $(4, 1, 1)$ or $(2, 2, 2)$.*

4 Shore constraints

From now on G is a 6-regular target and A a shore of a nontrivial 6-edge-cut, $a = |A| \geq 2$, $H = G[A]$, $b(v) = 6 - d_H(v)$.

Lemma 4.1. *$e(H) = 3a - 3$; H is connected; $0 \leq b(v) \leq 5$ and $\sum_{v \in A} b(v) = 6$.*

Proof. Degree sum: $6a = 2e(H) + 6$. If H had components $A_1, \dots, A_t$, the cuts $\partial_G(A_i)$ are disjoint subsets of $\partial_G(A)$, each of size at least 6 by $\lambda(G) \geq 6$ [5], and their sizes sum to 6; so $t = 1$. A vertex with $b(v) = 6$ would be isolated in H . $\square$

Lemma 4.2 (boundary-shortfall lemma). *For every $v \in A$ there exists a 3-colouring ψ of $H - v$ with*

$$\sum_{i=1}^3 \max(0, 2 - c_i(\psi)) \leq b(v), \quad \text{where } c_i(\psi) = |\{u \in N_A(v) : \psi(u) = i\}|.$$

Proof. Take a 3-colouring φ of $G - v$ and let $\psi = \varphi|_{A-v}$. By Corollary 2.2, every colour appears at least twice on $N_G(v)$, which is the disjoint union of $N_A(v)$ and the set C_v of outside cut-neighbours of v , $|C_v| = b(v)$. Writing o_i for the number of vertices of C_v coloured i we get $c_i + o_i \geq 2$, hence $\max(0, 2 - c_i) \leq o_i$, and summing over i gives the bound since $\sum_i o_i = b(v)$. $\square$

For v with $b(v) = 0$ this says: $H - v$ must admit a colouring in which $N(v)$ splits exactly $2 + 2 + 2$ — full-degree vertices must be “deletion-unfrozen”. Note also that for $u, v \in A$ nonadjacent with $b(u) = 0$, Lemma 2.4 localizes: $N_A(u) \subseteq N_A(v)$ is impossible, because then $N_G(u) = N_A(u) \subseteq N_G(v)$.

5 Small shores

Theorem 5.1. *No 6-edge-cut in a target (6-regular or not) has a shore of size $2 \leq a \leq 8$.*

Proof. By $\delta(G) \geq 6$, $e(H) \geq 3a - 3$, while H is 3-colourable, so $e(H) \leq \lfloor a^2/3 \rfloor$. For $2 \leq a \leq 7$ these conflict ($\lfloor a^2/3 \rfloor < 3a - 3$; this numeric fact is also Lean-checked). For $a = 8$ both bounds equal 21, forcing equality: H is the Turán graph $K_{3,3,2}$ and every vertex of A has G -degree exactly 6. The two vertices of the part of size 2 then have all six edges inside A , so they are nonadjacent vertices with equal neighbourhoods in G , contradicting Lemma 2.4. $\square$

Theorem 5.2. *In a 6-regular target, no 6-edge-cut has a shore of size $9 \leq a \leq 14$.*

Proof (computer-assisted). By Lemma 4.1 a shore of size a induces a connected graph on a vertices with exactly $3a - 3$ edges and maximum degree at most 6. We enumerated all such graphs with **geng** (nauty 2.8.9) and applied the necessary conditions of Sections 3–4: 3-colourability; Corollary 3.3 for *every* 3-colouring; the localized Lemma 2.4; and Lemma 4.2 for every vertex. The outcome:

a	generated	not 3-col.	fail Cor. 3.3	fail Lem. 2.4	fail Lem. 4.2	survive
9	729	711	9	8	1	0
10	18 655	18 345	197	86	27	0
11	696 208	687 377	6 013	1 300	1 518	0
12	32 833 744	32 484 081	241 863	27 322	80 478	0
13	1 839 349 287	1 822 133 664	11 944 366	788 481	4 482 776	0
14	119 236 283 370	118 201 012 144	719 933 200	28 146 103	287 191 923	0

No graph survives, proving the theorem. Verification methodology (independent double implementation, generator completeness guards, per-graph certificates) is described in Section 6. $\square$

Remark 5.3 (the near-miss at $a = 9$). Exactly one graph survives all the *counting* filters at $a = 9$: $H_9 = K_{3,3,3} \setminus M_3$, where M_3 is a perfect rainbow matching of size 3 (one missing edge between each pair of colour classes, with six distinct endpoints; graph6 **HEzftz**{). H_9 has a unique 3-colouring up to colour permutation; its three full-degree vertices — one in each class — have all neighbours inside A . For each such vertex v , all six colourings of $H_9 - v$ leave some colour with at most one occurrence on $N(v)$, so Lemma 4.2 (with $b(v) = 0$) excludes H_9 . This is the only point at $a \leq 13$ where the boundary-shortfall lemma is essential for a graph that the other filters miss, and we verified the kill additionally by brute force over all 3^8 assignments.

Proof of Theorem 1.2. A nontrivial 6-edge-cut has shores of sizes ≥ 2 ; Theorems 5.1 and 5.2 exclude sizes 2..14, so both shores have size at least 15, forcing $n \geq 30$. For $n \leq 29$ every 6-edge-cut is therefore trivial, i.e. a vertex star. $\square$

6 Computations, certificates, and machine-checked lemmas

Theorem A. All 6-regular graphs on $n \leq 14$ vertices were enumerated with **geng -d6 -D6** (nauty 2.8.9) and classified by a C++ backtracking 3-colourability checker into: 3-colourable; $\chi \geq 4$ but not vertex-critical; 4-vertex-critical with a critical edge; target. Counts:

n	total	3-colourable	not vertex-critical	4-VC with critical edge	target
11	266	3	263	0	0
12	7 849	50	7 799	0	0
13	367 860	849	367 010	1	0
14	21 609 301	42 667	21 566 634	0	0

The $n = 14$ run was executed twice with different residue-class chunkings (mod 110 and mod 73) with identical aggregates, and the total matches the known count of 6-regular graphs on 14 vertices. The unique $n = 13$ graph is

Its 13 critical edges form a Hamilton cycle. Across its 13 vertex-deleted subgraphs it admits exactly 78 proper 3-colourings (as maps to $\{1, 2, 3\}$, not modulo S_3), and *none* of them splits the open neighbourhood $2 + 2 + 2$: every colouring leaves a singleton class in $N(v)$, so Lemma 2.1 predicts critical edges at every vertex; the 156 singleton predictions hit exactly the 13 critical edges (no false positives). Thus the unique smallest 6-regular 4-vertex-critical graph fails the target conditions everywhere locally, not marginally.

Theorem B. Candidate shores were generated by `geng -c -D6 a E:E` with $E = 3a - 3$; large cases ran as residue classes with a generator-side completeness guard (each class must end with `geng's >Z` terminator line, and the generated count must equal the classifier's count). The filter battery was implemented twice independently (C++ and Python) with identical counts at $a = 9, 10, 11$; the $a = 9$ kill was additionally brute-forced. Per-graph certificates (graph6 plus the failed filter, and for shortfall kills the killing vertex) are included for $a \leq 11$ in the ancillary files.

Machine-checked lemmas. The following are verified in Lean 4 / Mathlib (no `sorry`, no `native_decide`; axioms at most `propext`, `Classical.choice`, `Quot.sound`): the recolouring core of Lemma 2.1 (axiom-free); the count of 21 in Theorem 3.2, via an enumeration of the 3003 weak compositions of 6 into 9 parts together with a proved completeness lemma for the enumeration, and a bridge lemma stating the matrix form with $\forall \pi \in S_3$ diagonal bounds; and the numeric inequality $\lfloor a^2/3 \rfloor < 3a - 3$ for $2 \leq a \leq 7$ used in Theorem 5.1.

7 Concluding remarks

Theorem B turns the Skottová–Steiner edge-connectivity threshold into an exact small-cut impossibility statement, but it says nothing about super-6-edge-connected candidates, which is where Problem 5.2 now lives: a 6-regular target must have $n \geq 15$ (Theorem A), and if $n \leq 29$ it has no nontrivial 6-edge-cut at all (Theorem B). The shore-exclusion machine is open-ended — each further size a is a finite computation; the $a = 14$ case (119,236,283,370 candidates, every residue class closed with a generator-side completeness guard) completed in under a day on a workstation — but we do not currently see an argument that it terminates for all a : the row-sum filter can be neutralized by small colour-forcing gadgets, and the burden falls entirely on the boundary-shortfall lemma. Proving that Corollary 3.3 forces enough colour rigidity for Lemma 4.2 to fail at some full-degree vertex, for every a , would upgrade Theorem B to: *every 6-regular $(4, 1)$ -graph is super-6-edge-connected*. We leave this as the natural next question.

References

- [1] T. Bloom, *Erdős problem #944*, Online database of Erdős problems, <https://www.erdosproblems.com/944>.
- [2] P. Erdős, *On some aspects of my work with Gabriel Dirac*, in: Graph Theory in Memory of G. A. Dirac (Sandbjerg, 1985), Ann. Discrete Math. **41**, North-Holland, 1989, 111–116.
- [3] T. R. Jensen, *Dense critical and vertex-critical graphs*, Discrete Math. **258** (2002), 63–84.

- [4] T. R. Jensen and M. Siggers, *On a question of Dirac on critical and vertex-critical graphs*, Sib. Elektron. Mat. Izv. **9** (2012), 156–160.
- [5] E. Skottová and R. Steiner, *Critical edge sets in vertex-critical graphs*, arXiv:2508.08703 (2025).