


# 3DNA (v1.5) — A Software Package for the Analysis, Rebuilding and Visualization of 3-Dimensional Nucleic Acid Structures



*by*

**Xiang-Jun Lu**

3dna.lu@gmail.com

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Installation</b>	<b>3</b>
2.1	BASEPARS . . . . .	4
2.2	FIBER . . . . .	5
2.3	bin . . . . .	5
2.4	Examples . . . . .	5
<b>3</b>	<b>How to run 3DNA</b>	<b>6</b>
<b>4</b>	<b>A tutorial introduction</b>	<b>10</b>
4.1	Base-pair geometry parameters . . . . .	10
4.2	Analysis of the Tc3 transposon DNA . . . . .	10
4.3	Fiber models . . . . .	15
4.4	Input for <code>render</code> in Raster3D etc . . . . .	16
4.5	Build a DNA structure with sugar-phosphate backbone . . . . .	16
<b>5</b>	<b>Some technical details</b>	<b>19</b>
5.1	Least-squares fitting procedures . . . . .	19
5.2	Base reference frames . . . . .	22
5.3	General rotation matrix . . . . .	23
5.4	Base-pair parameters . . . . .	23
5.5	Step parameters . . . . .	24
5.6	Local helical parameters . . . . .	26
5.7	Rebuilding based on local step parameters . . . . .	29
5.8	Rebuilding based on local helical parameters . . . . .	31
5.9	Relation between local helical and step parameters . . . . .	33
<b>6</b>	<b>Citation</b>	<b>35</b>
<b>7</b>	<b>Acknowledgments</b>	<b>35</b>
<b>8</b>	<b>Appendix: 3DNA output for Tc3 transposon DNA</b>	<b>36</b>

## 1 Introduction

Our recent comparison studies (Lu & Olson, 1999; Lu *et al.*, 1999) on seven popular analyzing programs for nucleic acid structures demonstrated that the choice of reference frame rather the mathematical formulation has the greater effect on the calculated parameters. Broadly speaking, there are two classes of reference frames. One is defined by the purine (R) C8 atom and pyrimidine (Y) C6 atom, and the mean base-pair plane as in FreeHelix/NewHelix (Dickerson, 1998). The other uses the average of the two constituent base reference frames in a base-pair as in Curves (Lavery & Sklenar, 1989).

While each has its own advantages and limitations, the base-centered one was recommended at a workshop held on January 13-14, 1999 at the AIST-NIBHT Structural Biology Center in Tsukuba, Japan (Olson *et al.*, 2001).

3DNA is a comprehensive software package for the analysis and rebuilding of nucleic acid structures based on the new recommended reference frame. Mathematically, it follows the CEHS definition (El Hassan & Calladine, 1995) as implemented in SCHNAaP/SCHNArP (Lu *et al.*, 1997a; Lu *et al.*, 1997b) with the global helical parameters replaced by a new set of *local* ones. This matrix based algorithm is rigorous and reversible in that it allows for exact reconstruction of the structure from a set of parameters which can be deduced from the structure. The parameters such defined also have simple geometrical meaning thus the distance between base-pair centers is  $\sqrt{\text{Shift}^2 + \text{Slide}^2 + \text{Rise}^2}$ , the bending angle between two base-pairs is  $\sqrt{\text{Roll}^2 + \text{Tilt}^2}$ , and the angle between base-pair normal and local helical axis is  $\sqrt{\text{Inclination}^2 + \text{Tip}^2}$ .

3DNA is a completely new written software package conforming strictly to the ANSI / ISO C. It is much more robust and efficient than SCHNAaP/SCHNArP and has new and improved features. Its analysis part can automatically classify a dinucleotide step as A-, B-, or TA-like (Guzikevich-Guerstein & Shakked, 1996) based on the positioning of phosphorus atoms with respect to the middle frame. This can be very useful for pinpointing conformational transitions in ligand-bound DNA, especially in a short fragment which is often neglected by other analyzing programs (Lu *et al.*, 2000). High quality “standardized” base-stacking diagrams can be generated by a utility program following the analysis. Authentic CEHS/SCHNAaP parameters are also available, which can be used for comparisons with the 3DNA recommended ones to see how a difference in reference frames can affect the calculated parameters, especially for Rise in heavily deformed DNA structures. Also note that FreeHelix/NewHelix parameters are very similar to the CEHS/SCHNAaP ones due to their similarity in reference frames. For completeness, local parameters using the seven methods (Lu & Olson, 1999; Lu *et al.*, 1999) based on the new recommended reference frame are also calculated. Obviously, they are all quite similar to 3DNA recommended ones.

The rebuilding part of 3DNA can be used for generating sequence-dependent atomic structures, without or with the sugar-phosphate backbone, suitable as starting point for molecular mechanics calculations and molecular dynamics simulations. It is also very convenient to generate publication quality Calladine-Drew style schematic representations of DNA, either in PS, XFIG<sup>1</sup>, or Raster3D (Merritt & Bacon, 1997)<sup>2</sup> format. With XFIG, the picture can be edited and annotated, combined with others, or exported to image formats such as GIF, JPEG and TIFF. With Raster3D, the Calladine-Drew style block representation of bases can be combined with the backbone and protein in either atomic or schematic presentations.

Fifty five different types of fiber DNA and RNA structures based on the work of Chandrasekaran & Arnott (1989) can also be conveniently generated. The atom naming and ordering conventions of the NDB are strictly followed which make it easy for direct comparison with X-ray crystal structures or theoretical models.

---

<sup>1</sup><http://www.xfig.org/>

<sup>2</sup><http://www.bmsc.washington.edu/raster3d/raster3d.html>

## 2 Installation

The latest version of 3DNA, in binary form for Linux, SGI and Windows, is available from the following URL: <http://rutchem.rutgers.edu/~xiangjun/3DNA/>. Download the version that fits your system and then do the following:

- `gunzip Linux_X3DNA_v1.5.tar.gz`, you get file `Linux_X3DNA_v1.5.tar`
- `tar xvf Linux_X3DNA_v1.5.tar`, you get directory `X3DNA` and four sub-directories underneath: `BASEPARS`, `FIBER`, `bin`, and `Examples`, the contents of which are detailed below.
- If `X3DNA` is not directly under your home directory, set environment variable `X3DNA` as follows:

```
in csh | tcsh:  setenv X3DNA Your_Directory_Containing_X3DNA
                e.g.,  setenv X3DNA /usr1/xiangjun/X3DNA
```

```
in bash | sh:  export X3DNA=Your_Directory_Containing_X3DNA
                e.g.,  export X3DNA=/usr1/xiangjun/X3DNA
```

```
in Windows (suppose 3DNA is installed under C:\X3DNA):
    set X3DNA=C:\X3DNA
```

Or do the followings (globally):

Start

Settings

Control Panel

System

Advanced

Environment Variables

at "User variables" section

Variable: `X3DNA`

Variable Value: `C:\X3DNA`

Then append to your command search path the `3DNA` bin subdirectory:

```
in csh | tcsh:  set path = ($path $X3DNA/bin)
```

```
in bash | sh:  export PATH=$PATH:$X3DNA/bin
```

in Windows:

Start

Settings

Control Panel

System

Advanced

Environment Variables

at "System variables" section

click on Path and "Edit" it

by appending `";C:\X3DNA\bin"` to it.

Another option for Windows users is to install CygWin from <http://www.cygwin.com/>.

## 2.1 BASEPARS

This directory contains standard residue geometry files and other parameters controlling various aspects of 3DNA. They are all in text format, so users can view the structures with `rasmol` or edit the parameters as they see fit.

- `Atomic_?.pdb` ( $? = A, C, G, T$  or  $U$ ) are the default standard residue geometries used by 3DNA for analyzing and rebuilding full atomic nucleic acid structures in PDB format.

Under its subdirectory `ATOMIC`, there are four sets of standard residue geometry files:

`ADNA_std?.pdb`, `BDNA_std?.pdb`, `NDB96_std?.pdb` and `RNA_std?.pdb`.

The base geometries by Clowney *et al.* (1996) are used (downloaded from the NDB archive: <http://ndbserver.rutgers.edu/NDB/archives/index.html>).

The NDB96 set includes base and C1' atoms, without sugar-phosphate backbone. ADNA set uses C3'-endo sugar-backbone conformation as defined by Leslie *et al.* (1980) fiber studies, but with a  $\chi$  torsion angle of  $-157^\circ$ , average of high resolution single crystal X-ray oligonucleotide structures. RNA set is the same as ADNA except for an additional O2' atom for each residue. BDNA set is similarly defined as ADNA except for a C3'-endo sugar fiber sugar-backbone conformation and the  $-108^\circ$   $\chi$  torsion angle.

The NDB96 set is the default. To use another data set, simply overwrite the corresponding `Atomic_?.pdb` in `BASEPARS` or copy them to your current working directory. A utility program, `cp_std` (see below), can do this automatically for you. You can also use other residue geometries with the help of the utility program `std_base`.

Note the standard set contains only the five common residues, A, C, G, T and U. Residue I can be got by deleting the N2 atom from G. Their modified counterparts, +A, +C, +G, +I, +T, +U which are changed to lower case by 3DNA, can be approximated by using their normal forms. For +C, for example, 3DNA requires file `Atomic_c.pdb`, which can be simply a copy of `Atomic_C.pdb`.

- `Block_BP.alc` defines the default base-pair rectangular block. It has a size of 10 Å (long) by 4.5 Å (wide) by 0.5 Å (thick) and is in `ALCHEMY` format. It is used for drawing the Calladine-Drew style schematic presentation of DNA structures. `Block_R.alc` is for the purine base (R) ( $4.5 \times 4.5 \times 0.5$ ). `Block_Y.alc` is for the pyrimidine base (Y) ( $3.0 \times 4.5 \times 0.5$ ).

Under its subdirectory `BLOCK`, there six block geometry files. `Block_M.alc` has half the size of `Block_BP.alc`, and can be used if you would like the two blocks consisting a base-pair to be of the same size. `Block_Ms.alc` is slightly smaller in length than `Block_M.alc` to avoid possible overlaps in a compressed base-pair (*i.e.* with negative `Stretch`). Furthermore, the blocks do not necessarily to be rectangular, as shown in `Block_R_nr.alc`.

- `Pxyz.dat` contains the *xyz* coordinates of phosphorus atoms with regard to the middle dinucleotide reference frame. Four sets were defined, corresponding respectively to average values in high resolution A- and B-DNA crystal structures, their intermediate and TA-DNA. New set can be added following the format. This file is used by `rebuild` for generating DNA structures with only base and phosphorus atoms. `PxyzH.dat` is the same as above except the coordinates are given in terms of the middle helical frame.
- `fig_image.par` contains parameters controlling the style of generated XFIG files, which can be edited by users to suit their liking. Similarly, `ps_image.par` holds parameters defining the drawing style for postscript images. Finally, `raster3d.par` and `my_header.r3d` are for Raster3D input.
- `misc_3dna.par` contains various parameters mainly for analyze and the utility program `find_pair`.
- `baselist.dat` contains a comprehensive list of currently known base residues and their standard counterpart. It makes analysis of unusual DNA and RNA structures straightforward.
- `trans_pep.pdb` & `trans_pep.alc` are trans peptide unit used for drawing peptide block in protein structures, in the same way as basepair blocks.

## 2.2 FIBER

This directory contains repeating unit for each type of the 55 fibers DNA and RNA structures. The original data as provided by Chandrasekaran & Arnott (1989) is given in subdirectory `Data`. Directories `Str01`–`Str55` store the “clean-up” version of each repeating unit in a format suitable for building the structure with utility program `fiber`.

## 2.3 bin

This directory contains executables of the 3DNA package. Most of which are utility programs with some in short Perl script. Detailed usage of each program is described in Section 3.

## 2.4 Examples

Four subdirectories are included to illustrate the various functionalities of the 3DNA package. You are strongly recommended to study these examples carefully in order to use 3DNA more effectively.

- `Analysis_Rebuild` contains the analysis/rebuilding results of four structures: `adh026` (A-DNA), `bd1084` (B-DNA), `pde0128` and `pd0001` (DNA-protein complexes). The `*.pdb` data files were downloaded from the NDB. The `*.inp` files are the corresponding input to the analysis routine (`analyze`) and `*.out` are the output containing various structural parameters.

Input file `multi_str.inp` illustrates how to analyze multiple structure from one input file.

README contains detailed information on how to run the `analyze/rebuild` programs to generate results in this directory, and the RMS deviations between 3DNA rebuilt structures and the experimental ones. For base atoms, the RMS is virtually zero, and with backbone atoms it less than 0.85Å even for the 146bp nucleosomal DNA.

- `Calladine_Drew` illustrates how to generate the two sets of DNA schematic pictures made popular by Calladine & Drew (1997) (Figures 1 and 2). The README file provides every detail. Note that the plots in these two figures are on the same scale and in the same orientation.




Figure 1: One complete *helical* turn of DNA having twist of  $36^\circ$ , showing the effects of introducing uniform roll and slide at each step (Calladine & Drew, 1997).

- NMR gives an example on how to analyze multiple NMR structures from the PDB. Follow the README file there.
- Stacking shows the procedures for generating “standardized” base-stacking diagrams (Figure 3). Check README file there.
- Triplex gives examples on how to analyze triplex and parallel duplex structures. Follow README file there for details.

### 3 How to run 3DNA

Running 3DNA is simple although it might take a while to use it more effectively. For each of the program, there is a simple on-line help illustrating its usage.

#### 1. Analyzing part




Figure 2: Two complete *helical* turns of DNA, with a curvature of  $45^\circ$  per turn, or  $4.5^\circ$  per step on average. Such tight curvature may be achieved, in principle by any of the distributions of roll angle shown in parts (a) to (d) (Calladine & Drew, 1997).

- `analyze [infile1 infile2 ...]`  
Sample input files are given in directory `Examples/Analysis`.
- `cehs [infile1 infile2 ...]`  
`cehs` gives the original CEHS/SCHNAaP parameters, to which `FreeHelix/NewHelix` ones should be quite similar. `cehs` is provide for completeness and comparison purpose.
- `find_pair [options] pdbfile infile`  
`find_pair` is used to generate an input file for `analyze/cehs`, starting directly from a PDB file. It can also generate input files for the popular nucleic acid analysis program `Curves` with option `-c`.
- `manalyze [-cehs] infile`  
A Perl script for analyzing multiple structures
- `nmr_strs [-cehs] infile n1 n2`  
A Perl script for analyzing multiple NMR structures

## 2. Rebuilding part

- `rebuild [options] [-negx] infile outfile`  
For rebuilding DNA structures of either atomic model in PDB format or schematic representation in `ALCHEMY` format.
- `regular_dna [options] outfile`  
A utility program to generate input file for `rebuild` for the construction of regular DNA structures.
- `frame_mol [options] -n1[,n2] reffile infile outfile`  
A utility program for attaching local helical frames and set the orientation of a structure.






Figure 3: “Standardized” base stacking diagrams of three consecutive dimer steps of the 1.4 Å B-DNA structure d(CGCGAATTCGCG)<sub>2</sub> (Shui *et al.*, 1998) (BDL084).

- `rotate_mol rotfile infile outfile`

A utility program for adjusting the orientation of a structure by rotations in MolScript style. In RasMol, a structure can be rotated etc, but the corresponding new coordinates can *not* be saved. `rotate_mol` is provided simply for this purpose: Rotate your structure in RasMol to the orientation you like, then write molscript rotfile.

- `alc2img [options] [-s=factor] infile outfile`  
Generating schematic representation of DNA from a ALCHEMY file.
- `pdb2img [options] [-s=factor] infile outfile`  
Generating schematic block representation of nucleic acid bases or protein peptide bonds directly from a PDB file.

### 3. Fiber models

- `fiber options pdbfile`

The options is `-num`, where num is between 1 and 38. `-a|-b|-c|-d|-z` can be used for A-DNA, B-DNA, C-DNA, D-DNA and Z-DNA models.

### 4. Stacking diagrams

- `stack2img [options] [-s=factor] infile outfile`  
Program to generate “standardized” base-stacking diagram and atomic representations of nucleic acid structures with filled base rings and hydrogen bonds.
- `mstack2img ['options'] stackfile n1 n2 prefix`  
A Perl utility calling `stack2img` to generate stacking diagrams for a set of dinucleotide structures.

### 5. Set standard bases

- `std_base [options] [angle1 distance angle2] infile outfile`  
A utility program for setting standard base geometries used by `analyze` and `rebuild`.

### 6. Other utilities

- `r3d_atom`  
A utility program for generating scenes of atomic CPK, ball-and-stick, filled-base-ring, hydrogen bonds between bases, colored by atomic type, base residue or in gray, for input to Raster3D. Some of its functionalities are similar to `rods`, `balls` and `ring3d` of the Raster3D distribution, but with a more general approach for locating base rings.
- `cp_std prefix`  
A Perl script for changing standard base geometry files.
- `ex_str -num [-nmr] pdbfile`  
A utility program for extracting a specific structure unit or an NMR structure from a multiple-structure PDB file.

- `get_part [options] infile outfile`  
Extracting protein, nucleic acid or other components from a PDB file.
- `olp_o2p infile [outfile]`  
Checking the designation of O1P and O2P atoms in a PDB file. Useful for RMS fittings.
- `block_atom`  
A simple Perl script for generating block plus atomic representation of a nucleic acid structure in `ALCHEMY` format, which can be displayed using `RasMol`. For example, with `block_atom bdl084` (assuming a PDB data file `bdl084.pdb` exists), one get a file `bdl084.alc`, which can be display with `rasmol -alchemy -noconnect bdl084.alc`. Note the undocumented option `-noconnect` is needed to avoid `RasMol` recalculating bond connections.
- `dcmnfile`  
A simple Perl script for clean-up. Various programs in `3DNA` generate auxiliary files with fixed names which can be deleted with this utility program.

## 4 A tutorial introduction

This section shows some detailed examples on how to use `3DNA` properly and effectively. Only some of the main features are covered for beginners to get started. As mentioned above, a serious user should go through the `Examples` directory provided with `3DNA` and play with it.

The universal molecular graphics program `rasmol` (version 2.6+, but not 2.7.1) is indispensable. Also, `gv` (`ghostview`), `xfig` (version 3.2+) and `Raster3D` (version 2.5+) are highly recommended. Most likely they might be already available on your machine, since they are all extremely valuable freewares.

### 4.1 Base-pair geometry parameters

Figures 4 and 5 illustrate the definition of various parameters describing the base-pair geometry in a nucleic acid structure. These two figures were themselves generated with the help of `3DNA`.

### 4.2 Analysis of the Tc3 transposon DNA

The crystal structure of Tc3 transposase in complex with transposon DNA was solved by van Pouderoyan *et al.* (1997). This transposon DNA structure is unusual in that it contains an A/B junction: the G+C rich end adopts the A-DNA conformation while the A+T rich part has the B-DNA conformation. As a consequence, this DNA is non-linear, *i.e.*, curved.

The NDB reference code for this structure is `pde0128`. Download the structure and display it with `rasmol`, you will find that protein and DNA are together. To analyze the helical DNA structure, the program (`analyze`) needs to know which base-




Figure 4: Pictorial definitions of parameters that relate complementary base pairs and sequential base-pair steps. The base-pair reference frame is constructed such that the  $x$ -axis points away from the (shaded) minor groove edge. Images illustrate positive values of the designated parameters (Dickerson *et al.*, 1989).




Figure 5: Helical parameters. Pictorial definitions of parameters that relate base pairs to its helical frame.

residue (sequential number in the PDB file) pairs with which. This information (in file `pde0128.inp`) can be obtained with the utility program `find_pair` as follows:

```
find_pair pde0128.pdb pde0128.inp
```

where `pde0128.pdb` is the data file in PDB format downloaded from the NDB and `pde0128.inp` is the file required by `analyze`.

```
analyze pde0128.inp
```

The above two steps can be combined as follows:

```
find_pair pde0128.pdb stdout | analyze
```

Among several other output files, the one named `pde0128.out` (see page 36) gives a detailed listing of the DNA structural parameters. Each part is briefly summarized as follows:

- `RMS deviation` ... gives the rms deviation between each experimental base and the standard reference. Normally the value is less than 0.05 Å. For each base, the corresponding PDB residue information is provided. A base-pair is classified as either Watson-Crick or non-Watson-Crick (denoted with a star).
- `H-bonding information` gives atom-list and their length in Å of all possible H-bonds in each of the base-pairs.
- `Overlap area` in Å<sup>2</sup> between polygons defined by atoms on successive bases. Polygons projected in the mean plane of the designed base-pair step.

Values in parentheses measure the overlap of base ring atoms only. Those outside parentheses include exocyclic atoms on the ring.

- `Origin (Ox, Oy, Oz)` ... gives the origins and mean base-pair normal vectors in the coordinate system of the given structure. Dickerson *et al.* has found it very useful by drawing the Ny vs. Nx normal plot to demonstrate the curvature of a DNA molecule.
- `Local base-pair parameters`, *i.e.*, Shear, Stretch, Stagger, Buckle, Propeller, and Opening (Figure 4). *Generally speaking*, Propeller always has a value of about  $-10^\circ$  in A- and B-DNA (Calladine & Drew, 1997), while Buckle can be either positive or negative of up to  $20^\circ$ . The other four show much less variations.
- `Local base-pair step parameters` gives the familiar Shift, Slide, Rise, Tilt, Roll, and Twist (Figure 4). Slide-Roll-Twist show more variations than the other three. Slide is the best single parameter among the six in discriminating A- from B-DNA.
- `Local base-pair helical parameters`, *i.e.*, X-displacement, Y-displacement, Helical rise, Inclination, Tip, and Helical twist, (Figure 5) are also commonly used. For canonical A-DNA, for example, the X-displacement characterizes the “hole” in the top view (Figure 6), and the Inclination means that the bases are non-perpendicular to the helical axis in the side view (see cover image).

- `lambda` ... gives the virtual angle between C1'-YN1 or C1'-RN9 glycosidic bonds and the base-pair C1'-C1' line. For mismatched base-pairs, the two  $\lambda$  angles are normally quite different.
- `Classification of each dinucleotide step` ... is based on the `Zp` and `ZpH` values. It applies to right-handed A-, B- and TA-DNA with Watson-Crick base-pair steps. This classification should be checked against the step/helical parameters and the backbone torsion angles etc for final assignment. Generally speaking, this classification can be taken as a good starting point for pinpointing possible structural transitions in large DNA structures. For `pde0128`, this section says that the transposon DNA is not simply a deformed B-DNA, but one with an A/B junction.
- `Minor and major groove widths` are calculated based on the method proposed by El Hassan & Calladine (1998). It uses simple cross-strand P-P distances with and without an angle correction, and assign the groove widths to each dinucleotide step. Please note that the sum of vdW radii (5.8 Å) of the two phosphate groups is **not** subtracted from the values given here.
- `Structure classification` gives the overall assignment of the structure as right-handed (A-, B-, TA-DNA), left-handed Z-DNA, and other two topologically possible forms.
- `Global linear helical axis` is defined by equivalent C1' and RN9/YN1 atom pairs as in `SCHNAaP`. It gives the unit vector along the helical axis and the two end points which can be used to locate the axis by adding these two points as `ATOM/HETATM` records in your PDB file and draw a line between them. While such a linear-fitting makes no much sense for a strongly curved DNA structure, the deviation from regular linear helix is a good measure of the overall structural deformation.
- `Main chain and chi torsion angles`.
- `Sugar conformational parameters` include five internal torsional angles, and amplitude and phase angle of pseudorotation of the sugar ring.
- `Same strand P-P and C1'-C1' virtual bond distances` have been used in the literature to discriminate A- and B-DNA conformations, although it is now clear that they are not as good as `Zp` and `Slide`.
- `Helix radius` gives the radial displacement of P, O4', and C1' atoms in local helix frame of each dimer.
- `Position (Px, Py, Pz)` and local helical axis vector (`Hx, Hy, Hz`) for each dinucleotide step.

The output file `bp_step.par` contains information for a rigorous reconstruction of the base-pair geometry. It will be overwritten each time when another analysis is performed unless it is renamed.

```

19 base-pairs
0 ***local base-pair & step parameters***
      Shear  Stretch  Stagger  Buckle  Prop-Tw  Opening  Shift  Slide  Rise  Tilt  Roll  Twist
G-C    -0.18   -0.27    0.58    8.57   -9.77   -8.26    0.00    0.00    0.00    0.00    0.00    0.00
G-C     0.16   -0.26    0.13   -5.72  -10.04   -3.52    0.19   -1.13    3.65    5.58    3.88   39.73
G-C    -0.10   -0.25    0.42   -1.54   -7.63   -0.67    0.43   -1.67    2.94   -3.14   10.32   29.99
G-C     0.02   -0.33    0.08   -3.56   -6.03    1.74    0.17   -2.24    3.13    1.31    6.72   31.31
G-C    -0.21   -0.24   -0.13   -3.78   -0.69    3.13    0.08   -1.92    3.33    3.67    8.74   27.62
G-C     0.04   -0.22   -0.35   -8.83  -11.08   -3.03    0.50   -1.91    3.41    4.23    5.60   31.56
G-C    -0.32   -0.40   -0.38  -16.06  -17.27   -3.53    0.18   -2.14    3.43    4.68   10.29   27.30
T-A     0.43   -0.40   -0.05  -11.97  -14.80   12.03   -0.07   -0.42    3.15   -0.69    5.91   35.56
C-G     0.52   -0.43   -0.34  -10.52    4.23    3.87    0.25    0.76    3.46    3.31    0.88   37.83
C-G     0.27   -0.33    0.52  -15.53   -4.19   -2.52   -1.73    0.69    3.77   -7.18   -1.10   31.39
T-A    -0.18   -0.23   -0.13   -9.98   -5.91   -1.56    0.35   -0.32    3.19    5.36    6.16   26.98
A-T    -0.57   -0.49    0.47    0.67   -7.27   -0.77    0.41    1.89    3.06   -6.05   -2.74   45.60
T-A    -0.34   -0.35    0.33   -3.99  -14.34   -5.83   -0.34   -0.18    3.44    1.73   -3.42   31.69
A-T    -0.69   -0.02   -0.04   -2.98   -0.45   12.91    0.29    2.18    3.37   -2.75   -6.21   49.54
G-C    -0.51   -0.07   -0.61    3.47   -9.27   12.05    0.33    1.29    3.25   -0.15   15.17   23.41
A-T    -0.66   -0.15   -0.12   12.32  -13.66    3.93   -0.68    0.30    2.96   -4.96    0.68   35.87
A-T    -0.41   -0.38    0.73   21.10  -23.76   -1.56   -0.06   -0.03    2.97   -6.65   -0.95   34.68
C-G     0.40   -0.55    0.33   -5.21  -10.24   -3.28    0.37   -1.03    3.93    4.89    1.28   38.27
T-A     0.56   -0.49    0.46  -21.17   -8.34   -4.72   -0.19   -1.07    3.61    1.91    4.78   37.16

```

Let's rebuild an atomic structure with the following command:

```
rebuild -atomic bp_step.par tc3_base.pdb
```

Depending on your setting of the standard base geometry (*i.e.* `Atomic_?.pdb` files), you will get a structure with either only base atoms (the default) or with an approximate sugar-phosphate backbone attached. Use `rasmol` to have a look.

You can also rebuild a schematic Calladine-Drew style picture with the following commands:

```
rebuild bp_step.par tc3_bp1.alc [for one block per base-pair]
```

```
rebuild -block2 bp_step.par tc3_bp2.alc [for one block
per base]
```

You need to use `rasmol` with the `-alchemy` command-line option to display files `tc3_bp1.alc` and `tc3_bp2.alc` since they are in `ALCHEMY` format. *e.g.*,

```
rasmol -alchemy tc3_bp2.alc
```

With this image, the Buckle and Propeller deformations are immediately obvious.

### 4.3 Fiber models

The 55 types of fiber nucleic acid models by Chandrasekaran & Arnott (1989) can be easily generated with the program `fiber`. Use `fiber -m` to get a list of all structures. Here I will use calf thymus A-DNA (number 1 in the list) as an example to illustrate its usage.

```
fiber -a fiber_A.pdb
```



Here `-a` is the same as `-l` and `fiber_A.pdb` is the output file name. You will then be asked to input your base sequence. It could either be from a data file (complete sequence) or from keyboard (enter only the repeating sequence, which is the default). Type `enter` means the default for input from keyboard. You are prompted for repeating unit with a default for polyA. Type `atcg` (either case is Okay and uncommon bases will be ignored) for a mixed A-T-C-G repeating sequence. Finally you are asked for the number of repeats (default is 10). Type 5 so you get  $(ATCG)_5$ , *i.e.*, 20 base-pairs. Display it with `rasmol fiber_A.pdb`.

High quality postscript picture (Figure 6) of `fiber_A.pdb` can be generated with the utility `stack2img` as follows:

```
stack2img -cao fiber_A.pdb fiber_A.ps
```

The option `-cao` means color-coded (c), atomic-model (a) with filled base-rings (o). If you add the option `-f`, you will get an image in XFIG format which you can easily edit. This is actually what these *atomic* options are intended for.

## 4.4 Input for render in Raster3D etc

The Calladine-Drew style base-pair representations are themselves quite useful as shown in Figures 1 and 2. They are, however, even more helpful when combined with atomic and schematic representations of ligands and proteins etc. This can be achieved with the `render` program of Raster3D as follows:

Use RasMol to find the view you want, and write `molscript molfile` at the command window. Then use the utility program `rotate_mol` to get a new PDB file with coordinates corresponding to your chosen view (RasMol does not write back coordinates in a new view). The new PDB coordinates are the *common reference* for scenes generated by different programs (3DNA, MolScript, Raster3D etc) to be properly rendered by Raster3D. Specifically, avoid any further coordinates transformation by MolScript or Raster3D.

You might use the utility `get_part` to divide a structure into nucleic acid part (default or `-n`), protein part (`-p`) and others (`-o`) to be used with different programs. For example, use 3DNA utilities `r3d_atom` and `pdb2img/stack2img` for nucleic acids, and MolScript for the protein part. You might need to delete the header part from MolScript with `del_ms -n`, a simple Perl script comes with 3DNA. The different scenes can then be concatenated together for render as detailed in Raster3D document.

Most of the images in 3DNA homepage were generated this way.

## 4.5 Build a DNA structure with sugar-phosphate backbone

The rebuilding of a DNA structure relies on its base sequence and associated base-pair and step parameters, and a set of standard `Atomic_?.pdb` data file.

The rebuilding algorithm is rigorous as far as the base-pair geometry is concerned. However, approximate sugar-phosphate conformations can be attached to the bases as a nucleic acid structure is built. This way, you get a full atomic structure with the sugar-phosphate backbone.




Figure 6: Top and side view images of the 20 base-pair long, (ATCG)<sub>5</sub>, fiber A-DNA color coded by residue: A-red, T-blue, G-green, and C-yellow.

In directory BASEPARS/ATOMIC, there are four sets of standard PDB files, corresponding to A-DNA (C3'-endo), B-DNA (C2'-endo), RNA (C3'-endo plus O2') and NDB96 (default, without backbone). The base geometry in each set is identical to ensure the exact same numerical values of base pair parameters no matter which set is used in the analysis procedure.

To use a set other than the default NDB96, simply overwrite these files to their corresponding `Atomic_?.pdb` counterparts (e.g., `BDNA_A.pdb` to `Atomic_A.pdb`). A simple Perl script `cp_std` helps with this process. For example: `cp_std BDNA` will put the B-DNA set into your current directory.

3DNA searches for `Atomic_?.pdb` files in the following order:

- current working directory
- `$X3DNA/BASEPARS` (default)
- `$HOME/X3DNA/BASEPARS` (if environment variable `$X3DNA` is not defined)

As an example, the procedure to rebuild `bd1084` with standard B-DNA C2'-endo sugar-phosphate backbone conformation is as follows:

```
find_pair -t bd1084.pdb stdout | analyze
```

You will get file `bp_step.par` which contains base-pair sequence and step parameters. (Another file, `bp_helical.par`, which contains helical parameters, is also suitable for the following rebuilding process.)

```
rebuild -atomic bp_step.par temp.pdb
```

Use RasMol to have a look at `temp.pdb`. It contains explicitly all valence bonds.

To verify that the algorithms used in 3DNA is rigorous and reversible, re-analyze the rebuilt structure `temp.pdb`, and you will get the same base-pair parameters as those directly from `bd1084.pdb`. Alternatively, you could superimpose `temp.pdb` to `bd1084.pdb`, and you will find that the RMSD (in Å) is virtually zero if only base atoms (i.e., excluding the backbone) are considered. If the sugar-phosphate backbone atoms (full-atom model) are also used, the results for three DNA structures in `Examples/Analyze_Rebuild` are as follows:

	pd0001	bd1084	adh026
base-atom	0.05	0.02	0.03
full-atom	0.82	0.73	0.52*

In rebuilding full-atomic models, regular sugar-phosphate backbone conformation was used: B-type for `pd0001` and `bd1084`, and A-type for `adh026`. It should be noted that the labeling of O1P/O2P atoms in `adh026` from the NDB is not consistent with the convention. Thus the RMSD values for full-atom model corresponds those after correction with a utility program `o1p_o2p`.

By matching `Atomic_?.pdb` with base sequence, it is also possible to generate a nucleic acid structure with mixed A- and B-DNA conformations, as in `pde0128.pdb`.

## 5 Some technical details

In our experience, following step-by-step working examples is the best way to understand any nontrivial algorithm. To make things right, every technical detail counts. Over the years, we have received many requests asking for exactly how 3DNA parameters are calculated. Here, we provide details of some key 3DNA components so that interested users could understand them better and possibly apply the same techniques to other related situations.

As an example, we will use the first dinucleotide step GG/CC from A-DNA adh026 (with base sequence GGGCGCCC), included in 3DNA distribution, which shows Slide and Roll more clearly.

### 5.1 Least-squares fitting procedures

3DNA starts with a least-squares procedure to fit a standard base with an embedded reference frame to an observed base structure. It implements a closed-form solution of absolute orientation using unit quaternions first introduced by Horn (1987). This method can be applied when one or both of the structures are perfectly planar. This section is based on the following URL:

[http://rutchem.rutgers.edu/~olson/jmb/ls\\_fit.html](http://rutchem.rutgers.edu/~olson/jmb/ls_fit.html)

- Standard reference frame (Olson *et al.*, 2001)

Using base G as an example, its xyz coordinates in standard reference frame in PDB format are as follows (check BASEPARS directory for other cases):

ATOM	1	C1'	G A	1	-2.477	5.399	0.000
ATOM	2	N9	G A	1	-1.289	4.551	0.000
ATOM	3	C8	G A	1	0.023	4.962	0.000
ATOM	4	N7	G A	1	0.870	3.969	0.000
ATOM	5	C5	G A	1	0.071	2.833	0.000
ATOM	6	C6	G A	1	0.424	1.460	0.000
ATOM	7	O6	G A	1	1.554	0.955	0.000
ATOM	8	N1	G A	1	-0.700	0.641	0.000
ATOM	9	C2	G A	1	-1.999	1.087	0.000
ATOM	10	N2	G A	1	-2.949	0.139	-0.001
ATOM	11	N3	G A	1	-2.342	2.364	0.001
ATOM	12	C4	G A	1	-1.265	3.177	0.000

- Least-squares fitting procedure

Least-squares fitting in 3DNA uses only (available) ring atoms: nine for purines (' N9 ' ; ' C8 ' ; ' N7 ' ; ' C5 ' ; ' C6 ' ; ' N1 ' ; ' C2 ' ; ' N3 ' ; ' C4 '), and six for pyrimidines (' N1 ' ; ' C2 ' ; ' N3 ' ; ' C4 ' ; ' C5 ' ; ' C6 ').

Using the first G residue in chain A of adh026, denoted thereafter as A\_G1, the corresponding 9 ring atoms in standard (sX, sY, sZ) and experimental (eX, eY, eZ) structures are as follows:

		sX	sY	sZ	eX	eY	eZ
1	N9	-1.289	4.551	0.000	11.417	-2.904	-4.880
2	C8	0.023	4.962	0.000	10.759	-1.995	-5.662
3	N7	0.870	3.969	0.000	11.469	-0.913	-5.867
4	C5	0.071	2.833	0.000	12.638	-1.108	-5.156
5	C6	0.424	1.460	0.000	13.759	-0.273	-5.036
6	N1	-0.700	0.641	0.000	14.767	-0.848	-4.249
7	C2	-1.999	1.087	0.000	14.663	-2.116	-3.719
8	N3	-2.342	2.364	0.001	13.625	-2.934	-3.830
9	C4	-1.265	3.177	0.000	12.625	-2.328	-4.545
-----							
s_ave:		-0.6897	2.7827	0.0001	e_ave:	12.8580	-1.7132 -4.7716

Where s\_ave and e\_ave are the geometric centers of the nine ring atoms in the standard and experimental bases respectively.

We collect the two sets of coordinates in the  $9 \times 3$  matrices S and E corresponding respectively to the standard and experimental bases. We then construct  $3 \times 3$  covariance matrix (C) between S and E using the following formula:

$$C = \frac{1}{N-1} [S' E - \frac{1}{N} S' i i' E]$$

$$=$$

-0.6849	0.8393	-0.7418
-2.1919	-0.5683	-0.7888
0.0001	-0.0002	0.0001

Here N, the number of atoms in each base, is 9, and i is an  $N \times 1$  column vector consisting of only ones. S' and i' are the transpose of matrix S and column vector i respectively.

From the nine elements of C, we subsequently generate the  $4 \times 4$  real symmetric matrix M using the expression:

$$M = \begin{vmatrix} c_{11}+c_{22}+c_{33} & c_{23}-c_{32} & c_{31}-c_{13} & c_{12}-c_{21} \\ c_{23}-c_{32} & c_{11}-c_{22}-c_{33} & c_{12}+c_{21} & c_{31}+c_{13} \\ c_{31}-c_{13} & c_{12}+c_{21} & -c_{11}+c_{22}-c_{33} & c_{23}+c_{32} \\ c_{12}-c_{21} & c_{31}+c_{13} & c_{23}+c_{32} & -c_{11}-c_{22}+c_{33} \end{vmatrix}$$

$$=$$

-1.2530	-0.7886	0.7419	3.0312
-0.7886	-0.1167	-1.3526	-0.7417
0.7419	-1.3526	0.1165	-0.7889

$$\begin{matrix} 3.0312 & -0.7417 & -0.7889 & 1.2533 \end{matrix}$$

The largest eigenvalue of matrix M is 3.5896, and its corresponding unit eigenvector ( $q_i$ ,  $i = 0 \dots 3$ ) is:

$$[ \ q_0 \quad q_1 \quad q_2 \quad q_3 \ ] = [ \ 0.5460 \quad -0.2921 \quad 0.0524 \quad 0.7835 \ ]$$

The rotation matrix R is deduced from the  $q_i$  as:

$$\begin{aligned} R &= \begin{pmatrix} q_0q_0+q_1q_1-q_2q_2-q_3q_3 & 2(q_1q_2-q_0q_3) & 2(q_1q_3+q_0q_2) \\ 2(q_2q_1+q_0q_3) & q_0q_0-q_1q_1+q_2q_2-q_3q_3 & 2(q_2q_3-q_0q_1) \\ 2(q_3q_1-q_0q_2) & 2(q_3q_2+q_0q_1) & q_0q_0-q_1q_1-q_2q_2+q_3q_3 \end{pmatrix} \\ &= \begin{pmatrix} -0.2331 & -0.8862 & -0.4004 \\ 0.8249 & -0.3983 & 0.4012 \\ -0.5150 & -0.2368 & 0.8238 \end{pmatrix} \end{aligned}$$

Following coordinate transformation with matrix R, the translation vector for the least-squares fit between standard and experimental bases is defined as:

$$o = e_{ave} - s_{ave} R' = [15.1632 \quad -0.0362 \quad -4.4678]$$

Here o and R are the origin and orientation, i.e., the reference frame, of the experimental base. Moreover, R is guaranteed to be orthogonal.

The least-squares fitted coordinates (F) of the standard base atoms on the experimental structure are then given by:

$$\begin{aligned} F &= S R' + o \\ &= \begin{pmatrix} 11.4307 & -2.9119 & -4.8818 \\ 10.7607 & -1.9933 & -5.6548 \\ 11.4432 & -0.8991 & -5.8558 \\ 12.6361 & -1.1058 & -5.1753 \\ 13.7706 & -0.2678 & -5.0319 \\ 14.7583 & -0.8689 & -4.2591 \\ 14.6659 & -2.1180 & -3.6958 \\ 13.6138 & -2.9091 & -3.8208 \\ 12.6427 & -2.3449 & -4.5688 \end{pmatrix} \end{aligned}$$

Here S is the ( $N \times 3$ ) matrix of the original coordinates of the standard base.

The difference matrix (D) between F and E, the ( $N \times 3$ ) matrix of the original coordinates of the experimental base, and the root-mean-square (RMS) deviation between the two structures are found to be:

```

D = E - F
=
-0.0137    0.0079    0.0018
-0.0017   -0.0017   -0.0072
 0.0258   -0.0139   -0.0112
 0.0019   -0.0022    0.0193
-0.0116   -0.0052   -0.0041
 0.0087    0.0209    0.0101
-0.0029    0.0020   -0.0232
 0.0112   -0.0249   -0.0092
-0.0177    0.0169    0.0238

```

```
RMS deviation = sqrt(sum(d^2)/N) = 0.0236
```

## 5.2 Base reference frames

Following the above least-squares procedure, we get the following for A\_G1:

```

RMSD = 0.0236
o = [15.1632   -0.0362   -4.4678]
R = [
-0.2331   -0.8862   -0.4004
 0.8249   -0.3983    0.4012
-0.5150   -0.2368    0.8238]

```

where the first column corresponds to x-axis, the 2nd to y-axis, and the 3rd to z-axis respectively.

Similarly for B\_C8 (i.e., C8 on chain B) which pairs with A\_G1, we have the following:

```

RMSD = 0.0229
o = [14.9124    0.2803   -4.7498]
R = [
-0.2339    0.9100    0.3422 ==> -0.2339   -0.9100   -0.3422
 0.7496    0.3930   -0.5326 ==>  0.7496   -0.3930    0.5326
-0.6191    0.1320   -0.7741 ==> -0.6191   -0.1320    0.7741]

```

Since their dot product is negative (-0.9884), the z-axis of B\_C8 is anti-parallel to that of A\_G1. We reverse the y- and z-axes of B\_C8 (by a 180-degree rotation around the x-axis) to make its z-axis parallel to that of A\_G1, as shown on the right side of R following the arrows.

For A\_G2, we have:

```

RMSD = 0.0171
o = [14.8757    2.9250   -2.4635]
R = [
-0.6807   -0.6274   -0.3781

```

```

0.3893    -0.7471    0.5388
-0.6205    0.2195    0.7528]

```

For B\_C7, which pairs with A\_G2, we have:

```
RMSD = 0.0213
```

```
o = [14.4982    3.0313   -2.3001]
```

```
R = [
```

```

-0.5797    0.6905    0.4326 ==>  -0.5797    -0.6905    -0.4326
 0.3207    0.6814   -0.6579 ==>   0.3207    -0.6814     0.6579
-0.7491   -0.2426   -0.6165 ==>  -0.7491     0.2426     0.6165]

```

### 5.3 General rotation matrix

The general rotation matrix  $\mathbf{R}_{\hat{\mathbf{u}}}(\varphi)$  (Eq. [1]) describes a rotation of magnitude  $\varphi$  about an arbitrary unit vector  $\hat{\mathbf{u}} = u_1\hat{\mathbf{i}} + u_2\hat{\mathbf{j}} + u_3\hat{\mathbf{k}}$ , where  $\hat{\mathbf{i}}, \hat{\mathbf{j}}, \hat{\mathbf{k}}$  are unit vectors along the axes of the local Cartesian frame:

$$\mathbf{R}_{\hat{\mathbf{u}}}(\varphi) = \begin{bmatrix} \cos \varphi + (1 - \cos \varphi)u_1^2 & (1 - \cos \varphi)u_1u_2 - u_3 \sin \varphi & (1 - \cos \varphi)u_1u_3 + u_2 \sin \varphi \\ (1 - \cos \varphi)u_1u_2 + u_3 \sin \varphi & \cos \varphi + (1 - \cos \varphi)u_2^2 & (1 - \cos \varphi)u_2u_3 - u_1 \sin \varphi \\ (1 - \cos \varphi)u_1u_3 - u_2 \sin \varphi & (1 - \cos \varphi)u_2u_3 + u_1 \sin \varphi & \cos \varphi + (1 - \cos \varphi)u_3^2 \end{bmatrix}, \quad (1)$$

### 5.4 Base-pair parameters

Given the above base reference frames for A\_G1, B\_C8 and A\_G2, B\_C7, we can calculate the base-pair parameters (Shear, Stretch, Stagger, Buckle, Propeller, and Opening). The procedure is exactly the same as for step parameters detailed below. Here we just give the parameters for the two base-pairs, and users who are interested in knowing the details should work them through and should get exactly the same numbers.

	Shear	Stretch	Stagger	Buckle	Propeller	Opening
A_G1-B_C8	-0.4683	-0.1516	-0.0156	-5.4713	-6.7936	-2.8660
A_G2-B_C7	-0.1643	-0.2112	-0.3299	-4.6532	-9.8008	2.7357

The “middle frame” used in calculating the bp parameters becomes the bp reference frame. For base pair A\_G1-B\_C8, we have:

```
o1 = [15.0378    0.1221   -4.6088]
```

```
R1 = [
```

```

-0.2323   -0.8985   -0.3724
 0.7889   -0.3980    0.4682
-0.5689   -0.1851    0.8013]

```

For base pair A\_G2-B\_C7, we have:



```

o2 = [14.6869      2.9781     -2.3818]
R2 = [
  -0.6319     -0.6594     -0.4072
    0.3583     -0.7144      0.6010
  -0.6873      0.2339      0.6877]

```

These two reference frames are used in the next two sections to calculate step and helical parameters.

## 5.5 Step parameters

Given the 2 bp reference frames above, the following procedures are used to calculate step parameters (Shift, Slide, Rise, Tilt, Roll and Twist). It follows CEHS definition, as detailed in SCHNAaP.

1. Hinge axis is the cross product between z1 (3rd column of R1) and z2 (3rd column of R2):

```

[-0.3724      0.4682      0.8013] x [-0.4072      0.6010      0.6877]
= [-0.1596     -0.0702     -0.0332]. When normalized, it becomes:
[-0.8993     -0.3953     -0.1870]

```

Geometrically, hinge axis is the intersection line between the two base pair planes.

2. The RollTilt angle ( $\Gamma$ , i.e., net bending angle) is the magnitude between z1 and z2, which is given by their dot product:  $\Gamma = 10.2221^\circ$ .
3. Now we rotate R2 by  $(-0.5 * \Gamma = -5.1111)$  degrees around the hinge axis (see Eq. [1]):

```

R_hinge(-5.1111) = [
  0.9992     -0.0152      0.0359
  0.0181      0.9966     -0.0798
 -0.0345      0.0804      0.9962]
R2' = R_hinge(-5.1111) * R2 = [
 -0.6616     -0.6396     -0.3914
  0.4005     -0.7426      0.5368
 -0.6340      0.1984      0.7475]

```

Similarly, we rotate R1 by  $(+0.5 * \Gamma)$  degrees along the hinge axis:

```

R1' = R_hinge(+5.1111) * R1 = [
 -0.1982     -0.8986     -0.3914
  0.7441     -0.3978      0.5368
 -0.6381     -0.1848      0.7475]

```

By definition, the z-axes of R1' and R2' (the third column) are the same, i.e., after symmetric rotations, we have perfectly aligned the z-axes of the two bps.

4. The x-, y- and z-axes of the “middle frame” are simply the average between those of  $R1'$  and  $R2'$ , and by definition, they are orthogonal. The origin of the bp is the geometric average of  $o1$  and  $o2$ . For the above case, we have the “middle frame” as follows:

$$R_m = \begin{bmatrix} -0.4490 & -0.8033 & -0.3914 \\ 0.5977 & -0.5955 & 0.5368 \\ -0.6642 & 0.0071 & 0.7475 \end{bmatrix}$$

$$o_m = [14.8624 \quad 1.5501 \quad -3.4953]$$

5. The translational parameters (Shift, Slide, and Rise) are simply the projections of the vector linking from  $o1$  to  $o2$  onto the x-, y-, and z-axes of the “middle frame”:

$$\begin{aligned} [\text{Shift} \quad \text{Slide} \quad \text{Rise}] &= (o2 - o1) * R_m \\ &= [-0.3509 \quad 2.8561 \quad 2.2270] * R_m \\ &= [0.3853 \quad -1.4033 \quad 3.3349] \end{aligned}$$

6. Twist is the angle from  $y1$  to  $y2$  (or from  $x1$  to  $x2$  of the “rotated”  $R1'$  and  $R2'$  matrices respectively). Its sign is defined with reference to the “middle frame” z-axis (the 3rd column of  $R_m$ ), following right-handed rule for positive Twist angle. More specifically, the magnitude of the angle between  $y1$  and  $y2$  is:  $33.5296^\circ$ . The sign is determined by:

$$\begin{aligned} (y1' \times y2') \cdot z &= \\ [-0.2162 \quad 0.2965 \quad 0.4129] \cdot [-0.3914 \quad 0.5368 \quad 0.7475] &= 0.5524 \end{aligned}$$

So Twist is positive.

7. The phase angle ( $\phi$ ) is the angle from the hinge-axis to the “middle frame” y-axis (2nd column of  $R_m$ ). By definition, hinge-axis lies in the xy-plane of the “middle-frame” since it is perpendicular to the z-axis of the “middle-frame”. The phase angle also has a sign associate with it, determined in the same way as Twist shown above. In this case,  $\phi = +16.9598^\circ$ .

8. Roll is defined as:

$$\Gamma * \cos(\phi) = 10.2221 * \cos(16.9598 * \pi/180) = 9.7776^\circ.$$

Similarly, Tilt is defined as:

$$\Gamma * \sin(\phi) = 10.2221 * \sin(16.9598 * \pi/180) = 2.9818^\circ.$$

9. Overall, the six step parameters are:

Shift	Slide	Rise	Tilt	Roll	Twist
0.3853	-1.4033	3.3349	2.9818	9.7776	33.5296

## 5.6 Local helical parameters

The geometric approach described below gives exactly the same numerical values as those from the RNA (Babcock *et al.*, 1994). The pivot point issue does not apply here: it is only used in defining the base pair reference frame. Given the reference frames of the two base pairs, the procedure used in 3DNA to calculate the local helical parameters (x-displacement, y-displacement, helical rise, inclination, tip, and helical rise) is analogous to the one detailed above for step parameters, by using a tip-inclination combination.

Please note that to define a local helical axis, we need **two** base-pair reference frames. 3DNA finds the single-helical axis (which is actually  $dx \times dy$ ) that brings 1 to coincide with 2 by a Helical Twist angle. The position which this helix passes through is defined by Chasles' theorem as detailed in Figures 12 & 13 of Babcock *et al.* (1994). The calculation of x-displacement, y-displacement, tip and inclination is then exactly as described in SCHNAaP (Lu *et al.*, 1997a).

To make the above point clear, let's use A1-A2-A3 triplet as an example. First, A1-A2 define a local helical axis and a set of local base-pair helical parameters are calculated. In 3DNA, these parameters are defined in a symmetric manner that bp A1:T1 and bp A2:T2 have exactly the same values. Similarly, step A2-A3 defines another set of local base-pair helical parameters. Thus bp A2:T2 has two sets of helical parameters associated with it depending on its context, i.e., either with bp A1:T1 or with bp A3:T3. Moreover, the local Helical Rise and Helical Twist are directly related to a dinucleotide step. These are the reasons that "Local base-pair helical parameters" as given in 3DNA refer to base-pair steps. As a matter of fact, the schematic diagrams illustrating the local helical parameters (x-displacement, y-displacement, inclination and tip) as given in 3DNA website and this user's manual were based on two perfectly overlapped base pair blocks.

Using the first GG/CC dinucleotide step in adh026 as an example, the detailed procedure is as follows:

1. The local helical axis is defined by the cross product of  $(x_2 - x_1)$  and  $(y_2 - y_1)$ :

$$\begin{bmatrix} -0.3997 & -0.4307 & -0.1183 \end{bmatrix} \times \begin{bmatrix} 0.2391 & -0.3164 & 0.4190 \end{bmatrix} = \begin{bmatrix} -0.2179 & 0.1392 & 0.2294 \end{bmatrix}, \text{ which when normalized, gives: } \\ h = \begin{bmatrix} -0.6303 & 0.4026 & 0.6638 \end{bmatrix}$$

2. Local helical frame of base-pair 1:

- TipInclination angle ( $\Psi$ , in magnitude) is the angle between unit vectors  $h$  and  $z_1$ , and is  $17.2279^\circ$ .
- Hinge axis is defined by a cross product from  $h$  to  $z_1$ , normalized to give:  $\begin{bmatrix} 0.0399 & 0.8707 & -0.4902 \end{bmatrix}$ . It lies in the xy-plane of local helical frame given below.
- Rotate  $R_1$  through the above hinge axis by negative TipInclination angle (i.e.,  $-17.2279^\circ$ ) will align the resultant z-axis with  $h$ , which gives us the local helical reference frame:

$$\begin{aligned}
H1 &= R\_hinge(-17.2279) * R1 = [ \\
&\quad -0.1880 \quad -0.7532 \quad -0.6303 \\
&\quad 0.7504 \quad -0.5242 \quad 0.4026 \\
&\quad -0.6337 \quad -0.3973 \quad 0.6638]
\end{aligned}$$

3. Similarly, the local helical frame for base pair 2:

- TipInclination angle ( $\Psi$ , in magnitude) is the angle between unit vectors  $h$  and  $z2$ , and is  $17.2279^\circ$ . Note that by (symmetric) definition, it is the same as for base pair 1.
- Hinge axis is defined by a cross product from  $h$  to  $z2$ , normalized to give:  $[-0.4121 \ 0.5511 \ -0.7256]$
- Rotate  $R2$  through the above hinge axis by negative TipInclination angle (i.e.,  $-17.2279^\circ$ ) will align the resultant  $z$ -axis with  $h$ , which gives us the local helical reference frame:

$$\begin{aligned}
H2 &= R\_hinge(-17.2279) * R2 = [ \\
&\quad -0.5861 \quad -0.5091 \quad -0.6303 \\
&\quad 0.3139 \quad -0.8599 \quad 0.4026 \\
&\quad -0.7470 \quad 0.0381 \quad 0.6638]
\end{aligned}$$

4. The “middle helical frame” is the average of  $H1$  and  $H2$ , and by definition, it is orthogonal.

$$\begin{aligned}
Hm &= [ \\
&\quad -0.4058 \quad -0.6618 \quad -0.6303 \\
&\quad 0.5580 \quad -0.7256 \quad 0.4026 \\
&\quad -0.7238 \quad -0.1883 \quad 0.6638]
\end{aligned}$$

5. Helical twist is the angle from  $y(H1)$  to  $y(H2)$  (or from  $x(H1)$  to  $x(H2)$ ) of the local helical reference frames defined above, with reference to the local helical axis  $h$  for sign determination:  $35.0103^\circ$ .

6. Helical rise is the projection of the vector linking  $\circ 1$  to  $\circ 2$  onto the local helical axis  $h$ :  $2.8493 \text{ \AA}$ .

7. Use **base pair 1** to calculate tip and inclination (base pair 2 gives the same result):

- Phase angle  $\psi$  is defined from hinge axis ( $[0.0399 \ 0.8707 \ -0.4902]$ ) to positive tip axis (2nd column of  $H1$ ,  $[-0.7532 \ -0.5242 \ -0.3973]$ ) with reference the local helical axis  $h$  ( $[-0.6303 \ 0.4026 \ 0.6638]$ ) for sign control:  $106.9598^\circ$ .
- Tip is defined as:  $\Psi * \cos(\psi) = 17.2279 * \cos(106.9598 * \pi/180) = -5.0254^\circ$
- Inclination is defined as:  $\Psi * \sin(\psi) = 17.2279 * \sin(106.9598 * \pi/180) = 16.4787^\circ$

8. Get the origin of local helical frame of base-pair 1. It is based on Chasles' theorem as used in RNA (Babcock *et al.*, 1994, see Figure 13, page 141 for a detailed illustration). The procedure implemented in 3DNA is as follows:

- Vector AB is defined by:

$$\begin{aligned} \text{AB} &= [\text{o2} - \text{o1}] - \text{Helical\_Rise} * \text{h} = \\ &[-0.3509 \ 2.8561 \ 2.2270] - 2.8493 * [-0.6303 \ 0.4026 \ 0.6638] \\ &= [1.4451 \ 1.7089 \ 0.3357] \end{aligned}$$

- Vector AD is defined by an angle of

$$90 - 0.5 * \text{Helical\_Twist} = 72.4948$$

from vector AB with reference to h:

$$\text{AD} = \text{R\_h}(72.4948) * \text{AB}' = [-0.5182 \ 1.6306 \ -1.4812]'$$

$\text{R\_h}(72.4948)$  is the rotation matrix along h (Eq. [1]) by an angle of  $72.4948^\circ$ .

$\text{AB}'$  changes row vector AB to a column vector to be compatible with the

3-by-3 rotation matrix. When normalized, the AD vector is:  $[-0.2290 \ 0.7205 \ -0.6545]$

- The magnitude of AD is defined by:

$$\begin{aligned} \text{AD\_magnitude} &= \\ 0.5 * \text{AB\_magnitude} / \sin(0.5 * \text{Helical\_Twist} * \pi / 180) &= \\ = 0.5 * 2.2631 / 0.3008 &= 3.7618 \end{aligned}$$

- We then get “D”, the position vector where the local helical axis passes through. It is the origin of local helical frame of base pair 1.

$$\begin{aligned} \text{o1\_h} &= \text{o1} + \text{AD\_magnitude} * \text{AD} = \\ [15.0378 \ 0.1221 \ -4.6088] &+ 3.7618 * [-0.2290 \ 0.7205 \ -0.6545] \\ = [14.1764 \ 2.8326 \ -7.0709] \end{aligned}$$

The origin of the local helical frame of base pair 2 is given by:

$$\text{o2\_h} = \text{o1\_h} + \text{Helical\_Rise} * \text{h} = [12.3805 \ 3.9798 \ -5.1797]$$

The origin of the “middle helical frame” is the average of  $\text{o1\_h}$  and  $\text{o2\_h}$ :

$$[13.2785 \ 3.4062 \ -6.1253]$$

9. X-displacement and y-displacement are defined as the projections of the vector from  $\text{o1\_h}$  to  $\text{o1}$  onto the x- and y-axis of local helical frame of base pair 1 ( $\text{H1}$ ):

$$\begin{aligned} [\text{o1} - \text{o1\_h}] * \text{H1}(:, 1:2) &= \\ ([15.0378 \ 0.1221 \ -4.6088] - [14.1764 \ 2.8326 \ -7.0709]) * \text{H1}(:, 1:2) &= \\ = [-3.7562 \ -0.2063] \end{aligned}$$

So x-displacement =  $-3.7562 \text{ \AA}$  and y-displacement =  $-0.2063 \text{ \AA}$ .

Using base pair 2 gives *exactly the same values* due to symmetric definition.

10. Overall, the six local helical parameters are:

X-disp	Y-disp	Rise	Incl.	Tip	Twist
-3.7562	-0.2063	2.8493	16.4787	-5.0254	35.0103

## 5.7 Rebuilding based on local step parameters

The local step parameters defined in 3DNA are rigorous and thus reversible. Given a set of step parameters, the relative position and orientation of the two base pairs can be exactly reproduced by a rebuilding procedure as detailed in SCHNAaP/SCHNArP (Lu *et al.*, 1997a; Lu *et al.*, 1997b). Here we provide step-by-step working example so users can understand the algorithm better.

Using the six step parameters of the first GG/CC step in adh026 as calculated above:

Shift	Slide	Rise	Tilt	Roll	Twist
0.3853	-1.4033	3.3349	2.9818	9.7776	33.5296




Figure 7: Roll-tilt combination

1. We first Twist base-pairs 2 and 1 about the “middle frame”  $z$ -axis (which at this stage is coincident with the  $z$ -axes of both base-pair reference frames) through  $+\Omega/2$  and  $-\Omega/2$  respectively:

$$T2 = R_z(+\Omega/2), \quad T1 = R_z(-\Omega/2) \quad (2)$$

Note that to be consistent with previous literatures, the symbol used for Twist here is still  $\Omega$  (not  $\omega$ ), which is  $33.5296^\circ$ . The rotation matrix  $R_z$  corresponds to the Eq. [1] with unit vector  $[0 \ 0 \ 1]$ . Similarly,  $R_x$  corresponds to  $[1 \ 0 \ 0]$  and  $R_y$  corresponds to  $[0 \ 1 \ 0]$ . Here, the “middle frame” is the “global” reference frame and corresponds to an identity matrix.

2. The RollTilt angle ( $\Gamma$ ) is given by:

$$\Gamma = \sqrt{\rho^2 + \tau^2} = \sqrt{9.7776^2 + 2.9818^2} = 10.2221^\circ \quad (3)$$

3. The Roll-Tilt axis is determined by the values of Roll and Tilt, and it lies in the  $x$ - $y$  plane of the “middle frame”. For the above case, it is  $[0.2917 \ 0.9565 \ 0]$ , when normalized.

The angle from the Roll-Tilt axis ( $[0.2917 \ 0.9565 \ 0]$ ) to the “middle frame” y-axis (Roll) ( $[0 \ 1 \ 0]$ ) is called  $\phi$ . Its sign is determined with reference to the “middle frame” z-axis ( $[0 \ 0 \ 1]$ ), with values in range from  $-180^\circ$  to  $180^\circ$ . In this case, it is  $16.9598^\circ$ .

4. We now rotate T2 and T1 through  $+\Gamma/2$  and  $-\Gamma/2$  respectively about the Roll-Tilt axis to give the final orientation of the two base-pairs with respect to the “middle frame”. Analytically, the rotation about the Roll-Tilt axis can be expressed in terms of three rotations with reference to “middle frame” axes: First a rotation of angle  $\phi$  around z-axis to align Roll-Tilt axis with y-axis, then a rotation of angle  $\pm\Gamma/2$  around y-axis, and finally a rotation of angle  $-\phi$  around z-axis to bring the Roll-Tilt axis back. Overall, the required rotations can be written as:

$$\begin{aligned} T2^m &= [\mathbf{R}_z(-\phi) \mathbf{R}_y(+\Gamma/2) \mathbf{R}_z(+\phi)] T2 \\ &= \mathbf{R}_z(-\phi) \mathbf{R}_y(+\Gamma/2) \mathbf{R}_z(\phi + \Omega/2) \end{aligned} \quad (4)$$

$$\begin{aligned} T1^m &= [\mathbf{R}_z(-\phi) \mathbf{R}_y(-\Gamma/2) \mathbf{R}_z(+\phi)] T1 \\ &= \mathbf{R}_z(-\phi) \mathbf{R}_y(-\Gamma/2) \mathbf{R}_z(\phi - \Omega/2) \end{aligned} \quad (5)$$

5. To get the correct positions of the origins of the reference frame of base-pair 2 and 1, we apply the translation vectors  $(+D_x/2, +D_y/2, +D_z/2)$  and  $(-D_x/2, -D_y/2, -D_z/2)$  respectively. Both translations are defined with reference to the “middle frame”. Here  $D_x, D_y, D_z$  represent Shift, Slide and Rise.
6. The set of equations described above for the synthesis of a dinucleotide step defines the positions and orientations of the two base-pairs in a dinucleotide step with respect to the “middle frame”. However, by means of a set of simple matrix transformations, we can change the reference frame from the “middle frame” to the reference frame of the base-pair 1. The orientation and position of base-pair 2 and the “middle frame” with respect to base-pair 1 are now given by the following equations:

$$\mathbf{R}_m = [\mathbf{T}1^m]^{-1} = \mathbf{R}_z(\Omega/2 - \phi) \mathbf{R}_y(\Gamma/2) \mathbf{R}_z(\phi) \quad (6)$$

$$\mathbf{R}2 = [\mathbf{T}1^m]^{-1} \mathbf{T}2^m = \mathbf{R}_z(\Omega/2 - \phi) \mathbf{R}_y(\Gamma) \mathbf{R}_z(\Omega/2 + \phi) \quad (7)$$

$$\mathbf{o}2 = [D_x \ D_y \ D_z] \mathbf{R}_m' \quad (8)$$

Where  $\mathbf{R}_m'$  means the transpose of the “middle frame” expressed with reference to base pair 1.

Using the above set of step parameters, we have the following:

$$\begin{aligned} \mathbf{R}_m &= [ \\ &\quad 0.9537 \quad -0.2873 \quad 0.0891 \\ &\quad 0.2885 \quad 0.9575 \quad -0.0003 \\ &\quad -0.0852 \quad 0.0260 \quad 0.9960 ] \end{aligned}$$

$$\begin{aligned} \mathbf{R}2 &= [ \\ &\quad 0.8204 \quad -0.5436 \quad 0.1775 \\ &\quad 0.5524 \quad 0.8336 \quad -0.0006 \\ &\quad -0.1476 \quad 0.0985 \quad 0.9841 ] \end{aligned}$$

$$\mathbf{o}2 = [0.3853 \quad -1.4033 \quad 3.3349] * \mathbf{R}_m' = [1.0677 \quad -1.2336 \quad 3.2524]$$

Expressed in terms of the experimental coordinate reference frame of base pair 1, we get the reference frame for base pair 2 which are the same as shown in section 5.4: “Base-pair parameters”

$$\begin{aligned} R1\_exp = [ \\ & -0.2323 \quad -0.8985 \quad -0.3724 \\ & 0.7889 \quad -0.3980 \quad 0.4682 \\ & -0.5689 \quad -0.1851 \quad 0.8013 ] \end{aligned}$$

$$\begin{aligned} R2\_exp = R1\_exp * R2 = [ \\ & -0.6319 \quad -0.6594 \quad -0.4072 \\ & 0.3583 \quad -0.7144 \quad 0.6010 \\ & -0.6873 \quad 0.2339 \quad 0.6877 ] \end{aligned}$$

$$o1\_exp = [15.0378 \quad 0.1221 \quad -4.6088]$$

$$o2\_exp = o1\_exp + o2 * R1\_exp' = [14.6869 \quad 2.9781 \quad -2.3818]$$

## 5.8 Rebuilding based on local helical parameters

The local helical parameters defined in 3DNA are rigorous and thus reversible. Given a set of helical parameters, the relative position and orientation of the two base pairs can be exactly reproduced by a rebuilding procedure. Here we provide step-by-step working example so users can understand the algorithm better.

Using the six helical parameters of the first GG/CC step in adh026 as calculated above:

X-disp	Y-disp	Rise	Incl.	Tip	Twist
-3.7562	-0.2063	2.8493	16.4787	-5.0254	35.0103

There are four frames for a dinucleotide step consisting of base-pairs 1 and 2: the base-pair frames  $R_1$  and  $R_2$  and their corresponding helical frames  $R_{1h}$  and  $R_{2h}$ .  $R_{1h}$  and  $R_1$  are related by x-displacement, y-displacement, inclination and tip, so are  $R_{2h}$  with  $R_2$ .  $R_{1h}$  and  $R_{2h}$  are related by *helical* rise and *helical* twist.

### 1. $R_{1h}$ as reference ( $1h$ in superscript)

Starting with  $R_{1h}$  as the reference for the dinucleotide step concerned, then  $R_{1h}^{1h}$  is the identity matrix  $I$ :

$$R_{1h}^{1h} = I \quad (9)$$

$R_{2h}^{1h}$  is got by the helical twist:

$$R_{2h}^{1h} = R_z(\Omega_h) \quad (10)$$

$R_1^{1h}$  is got by a combined rotation of magnitude  $\Psi (= \sqrt{\eta^2 + \theta^2})$  along tip-inclination axis (Fig. 8 (a)). Thus  $\Psi$  is the exact angle between base-pair normal and the local






Figure 8: Combination of Tip and Inclination, and calculation of helical Twist.

helical helix. Let the angle from tip-inclination axis to tip axis be  $\psi$ , the rotation can be expressed as a rotation about  $z$ -axis by angle  $\psi$ , followed by a rotation about  $y$ -axis by angle  $\Psi$ , and then a rotation about  $z$ -axis by angle  $-\psi$  to bring the axis back:

$$\mathbf{R}_1^{1h} = \mathbf{R}_z(-\psi) \mathbf{R}_y(\Psi) \mathbf{R}_z(\psi) \quad (11)$$

As shown in Fig. 8 (b),  $\mathbf{R}_2^{1h}$  is got by a helical twist, and then a rotation about the tip-inclination axis of  $\mathbf{R}_{2h}^{1h}$  (which is related to the tip  $y$ -axis of  $\mathbf{R}_{1h}$  by the helical twist  $\Omega_h$ ):

$$\mathbf{R}_2^{1h} = [\mathbf{R}_z(\Omega_h - \psi) \mathbf{R}_y(\Psi) \mathbf{R}_z(\psi - \Omega_h)] \mathbf{R}_z(\Omega_h) = \mathbf{R}_z(\Omega_h - \psi) \mathbf{R}_y(\Psi) \mathbf{R}_z(\psi) \quad (12)$$

## 2. $\mathbf{R}_1$ as reference

With Eqs. 9 to 12, we can do some simple matrix transformations to make  $\mathbf{R}_1$  as the reference:

$$\mathbf{R}_1 = \mathbf{I} \quad (13)$$

$$\mathbf{R}_{1h} = [\mathbf{R}_1^{1h}]^{-1} = \mathbf{R}_z(-\psi) \mathbf{R}_y(-\Psi) \mathbf{R}_z(\psi) \quad (14)$$

$$\mathbf{R}_{2h} = [\mathbf{R}_1^{1h}]^{-1} \mathbf{R}_2^{1h} = \mathbf{R}_z(-\psi) \mathbf{R}_y(-\Psi) \mathbf{R}_z(\psi + \Omega_h) \quad (15)$$

$$\begin{aligned} \mathbf{R}_2 &= [\mathbf{R}_1^{1h}]^{-1} \mathbf{R}_2^{1h} \\ &= \mathbf{R}_z(-\psi) \mathbf{R}_y(-\Psi) \mathbf{R}_z(\psi) \mathbf{R}_z(\Omega_h - \psi) \mathbf{R}_y(\Psi) \mathbf{R}_z(\psi) \\ &= \mathbf{R}_z(-\psi) \mathbf{R}_y(-\Psi) \mathbf{R}_z(\Omega_h) \mathbf{R}_y(\Psi) \mathbf{R}_z(\psi) \end{aligned} \quad (16)$$

Equ. 16 describes the orientation of base-pair 2 with reference to 1.

## 3. Position vector of base-pair 2 with reference to 1

- From base-pair 1 origin to base-pair 1 helical origin:  $-[dx \ dy] [x_{1h} \ y_{1h}]'$
- From base-pair 1 helical origin to 2 helical origin:  $dz \cdot z'_{1h} = dz \cdot z'_{2h}$

- From base-pair 2 helical origin to base-pair 2 origin:  $+ [dx \ dy] [x_{2h} \ y_{2h}]'$

Combining the above three items, we have the position (base-pair origin) of 2 relative to 1 as follows:

$$\mathbf{o}_2 = [dx \ dy] \{ [x_{2h} \ y_{2h}]' - [x_{1h} \ y_{1h}]' \} + dz \cdot z'_{1h} \quad (17)$$

Note that  $x_{1h}, y_{1h}, z_{1h}, x_{2h}, y_{2h}, z_{2h}$  are  $3 \times 1$  column vectors.  $'$  means the transpose of a vector to change it to a row vector.

4. Using the example where inclination =  $16.4787^\circ$  and tip =  $-5.0254^\circ$ , we have  $\Psi = \sqrt{\eta^2 + \theta^2} = 17.2279^\circ$ . The tip-inclination axis lies in the x-y plane of local helical frame of base pair 1:  $[16.4787 \ -5.0254 \ 0]$ ; which when normalized gives:  $[0.9565 \ -0.2917 \ 0]$ . The angle from the tip-inclination axis ( $[0.9565 \ -0.2917 \ 0]$ ) to the base pair 1 helical y-axis (tip) ( $[0 \ 1 \ 0]$ ) with reference to the helical z-axis ( $[0 \ 0 \ 1]$ ) is  $\psi = 106.9598^\circ$ .

Using equations 15 to 17, we having the following:

$$\begin{aligned} R1h = [ & \\ & 0.9962 \quad -0.0125 \quad 0.0864 \\ & -0.0125 \quad 0.9590 \quad 0.2833 \\ & -0.0864 \quad -0.2833 \quad 0.9551] \end{aligned}$$

$$\begin{aligned} R2h = [ & \\ & 0.8087 \quad -0.5818 \quad 0.0864 \\ & 0.5399 \quad 0.7926 \quad 0.2833 \\ & -0.2333 \quad -0.1825 \quad 0.9551] \end{aligned}$$

$$\begin{aligned} R2 = [ & \\ & 0.8204 \quad -0.5436 \quad 0.1775 \\ & 0.5524 \quad 0.8336 \quad -0.0006 \\ & -0.1476 \quad 0.0985 \quad 0.9841] \end{aligned}$$

$$\mathbf{o}_2 = [1.0677 \quad -1.2336 \quad 3.2524]$$

Compared with the numbers based on step parameters, it is clear that Eqs. 16 and 7, Eqs. 17 and 8 are equivalent.

## 5.9 Relation between local helical and step parameters

To refer the orientation and position of one base-pair relative to the other, 6 parameters (3 rotations and 3 translations) are required. One set of such parameters is (Shift, Slide, Rise, Tilt, Roll and Twist), and the other set is (X-displacement, Y-displacement, Helical Rise, Inclination, Tip and Helical Twist).

Obviously these two sets should be completely reversible/dependent: from any one set you can get the other, rigorously. You can verify this point using `step_hel`, a utility program in 3DNA. Graphically this is best illustrated by the Calladine-Drew A to B transition model by introducing uniform Roll and Slide values at each dinucleotide step. The key point is that by introducing Roll, you also get Inclination, and with Slide, you get X-displacement.

The `rebuild` program in 3DNA can construct a DNA structure using either set of these parameters. Examples of such input files (e.g., `bp_step.par` and `bp_helical.par`) can be generated by `analyze` (`Examples/Analyze_Rebuild` directory.)

We have two sets of simple equations:

$$\theta/\eta = -\tau/\rho$$

$$2 \cos \Omega_h = \cos \Omega (1 + \cos \Gamma) - (1 - \cos \Gamma)$$

## 6 Citation

Xiang-Jun Lu & Wilma K. Olson (2003). “3DNA: a software package for the analysis, rebuilding and visualization of three-dimensional nucleic acid structures.” *Nucl. Acids. Res.* **31**(17), 5108-5121.

## 7 Acknowledgments

The development of 3DNA benefit greatly from extensive testing and enthusiastic feedbacks of an increasing user community. In particular, we would like to thank Andrew Colasanti, Haim Rozenberg, A. R. Srinivasan, Patrick Furrer, Karolin Luger and Zukang Feng for validating the program, and Zippi Shakked for constructive suggestions.

This work has been generously supported by the U. S. Public Health Service under research grant GM20861. Computations were carried out at the Rutgers University Center for Computational Chemistry and through the facilities of the Nucleic Acid Database project (NSF grant DBI 9510703).

## 8 Appendix: 3DNA output for Tc3 transposon DNA

Detailed structural parameters for the analysis of Tc3 transposon DNA (data file pde0128.out)

```
*****
*****
```

3DNA (v1.5, Nov. 2002) by Xiang-Jun Lu at Wilma K. Olson's Lab.

```
*****
```

1. The list of the parameters given below correspond to the 5' to 3' direction of strand I and 3' to 5' direction of strand II.

2. All angular parameters, except for the phase angle of sugar pseudo-rotation, are measured in degrees in the range of [-180, +180], and all displacements are measured in Angstrom units.

```
*****
```

File name: pde0128.pdb

Date and time: Tue Nov 12 00:26:08 2002

Number of base-pairs: 19

Number of atoms: 1287

```
*****
```

```
HEADER      PROTEIN/DNA                                07-JUL-97    1TC3
TITLE       TRANSPOSASE TC3A1-65 FROM CAENORHABDITIS ELEGANS
COMPND      MOL_ID: 1;
COMPND      2 MOLECULE: TC3 TRANSPOSASE;
COMPND      3 CHAIN: C;
COMPND      4 FRAGMENT: SPECIFIC DNA BINDING DOMAIN, RESIDUES 2 - 52;
COMPND      5 ENGINEERED: YES;
COMPND      6 MUTATION: C-TERMINAL 6-HIS TAG;
COMPND      7 MOL_ID: 2;
COMPND      8 MOLECULE: DNA
COMPND      9 (5'-D(*AP*GP*GP*GP*GP*GP*GP*TP*CP*CP*TP*AP*TP*AP*GP*A
COMPND     10 P*AP*CP*TP*T)-3');
COMPND     11 CHAIN: A;
COMPND     12 ENGINEERED: YES;
COMPND     13 MOL_ID: 3;
COMPND     14 MOLECULE: DNA
COMPND     15 (5'-D(*AP*GP*TP*TP*CP*TP*AP*TP*AP*GP*GP*AP*CP*CP*CP*CP*C
COMPND     16 P*CP*CP*T)-3');
COMPND     17 CHAIN: B;
COMPND     18 ENGINEERED: YES
SOURCE      MOL_ID: 1;
SOURCE      2 ORGANISM_SCIENTIFIC: CAENORHABDITIS ELEGANS;
SOURCE      3 STRAIN: BERGERAC;
SOURCE      4 VARIANT: TR679;
SOURCE      5 ORGANELLE: NUCLEUS;
SOURCE      6 GENE: TC3A;
SOURCE      7 EXPRESSION_SYSTEM: ESCHERICHIA COLI;
SOURCE      8 EXPRESSION_SYSTEM_STRAIN: BL21 (DE3) PLYSS;
SOURCE      9 EXPRESSION_SYSTEM_CELLULAR_LOCATION: CYTOPLASM;
SOURCE     10 EXPRESSION_SYSTEM_VECTOR_TYPE: PET3C;
SOURCE     11 EXPRESSION_SYSTEM_PLASMID: PRP1200;
SOURCE     12 EXPRESSION_SYSTEM_GENE: TC3A N1-65;
SOURCE     13 MOL ID: 2;
```

SOURCE 14 SYNTHETIC: YES;  
 SOURCE 15 MOL\_ID: 3;  
 SOURCE 16 SYNTHETIC: YES  
 KEYWDS TRANSPOSASE, DNA BINDING, HELIX-TURN-HELIX, TC1/MARINER  
 KEYWDS 2 FAMILY, COMPLEX (TRANSPOSASE/DNA), PROTEIN/DNA  
 EXPDTA X-RAY DIFFRACTION  
 AUTHOR G.VAN POUDEROYEN,R.F.KETTING,A.PERRAKIS,R.H.A.PLASTERK,  
 AUTHOR 2 T.K.SIXMA  
 REVSTAT 1 12-NOV-97 0  
 JRNL AUTH G.VAN POUDEROYEN,R.F.KETTING,A.PERRAKIS,  
 JRNL AUTH 2 R.H.A.PLASTERK,T.K.SIXMA  
 JRNL TITL CRYSTAL STRUCTURE OF THE SPECIFIC DNA-BINDING  
 JRNL TITL 2 DOMAIN OF TC3 TRANSPOSASE OF C. ELEGANS IN COMPLEX  
 JRNL TITL 3 WITH TRANSPOSON DNA  
 JRNL REF EMBO J. V. 16 6044 1997  
 JRNL REFN ASTM EMJODG UK ISSN 0261-4189 0897  
 HELIX 1 1 ASP C 209 LEU C 220 1  
 HELIX 2 2 LEU C 225 ILE C 232 1  
 HELIX 3 3 ARG C 236 LYS C 244 1

\*\*\*\*\*

RMSD of the bases (----- for WC bp, + for isolated bp, x for helix change)

	Strand I	Strand II	Helix
1	(0.025) A:...2_: [...G]G-----C[...C]:.119_:B (0.016)		
2	(0.028) A:...3_: [...G]G-----C[...C]:.118_:B (0.019)		
3	(0.022) A:...4_: [...G]G-----C[...C]:.117_:B (0.014)		
4	(0.017) A:...5_: [...G]G-----C[...C]:.116_:B (0.017)		
5	(0.027) A:...6_: [...G]G-----C[...C]:.115_:B (0.020)		
6	(0.020) A:...7_: [...G]G-----C[...C]:.114_:B (0.014)		
7	(0.034) A:...8_: [...G]G-----C[...C]:.113_:B (0.019)		
8	(0.019) A:...9_: [...T]T-----A[...A]:.112_:B (0.020)		
9	(0.009) A:...10_: [...C]C-----G[...G]:.111_:B (0.017)		
10	(0.010) A:...11_: [...C]C-----G[...G]:.110_:B (0.034)		
11	(0.023) A:...12_: [...T]T-----A[...A]:.109_:B (0.016)		
12	(0.023) A:...13_: [...A]A-----T[...T]:.108_:B (0.021)		
13	(0.018) A:...14_: [...T]T-----A[...A]:.107_:B (0.024)		
14	(0.026) A:...15_: [...A]A-----T[...T]:.106_:B (0.018)		
15	(0.017) A:...16_: [...G]G-----C[...C]:.105_:B (0.025)		
16	(0.033) A:...17_: [...A]A-----T[...T]:.104_:B (0.019)		
17	(0.031) A:...18_: [...A]A-----T[...T]:.103_:B (0.017)		
18	(0.012) A:...19_: [...C]C-----G[...G]:.102_:B (0.026)		
19	(0.021) A:...20_: [...T]T-----A[...A]:.101_:B (0.023)		

\*\*\*\*\*

Detailed H-bond information: atom-name pair and length [ON]

1	G-----C	[3]	O6 - N4	2.53	N1 - N3	2.85	N2 - O2	3.00
2	G-----C	[3]	O6 - N4	2.61	N1 - N3	2.80	N2 - O2	2.77
3	G-----C	[3]	O6 - N4	2.86	N1 - N3	2.80	N2 - O2	2.62
4	G-----C	[3]	O6 - N4	2.72	N1 - N3	2.64	N2 - O2	2.40
5	G-----C	[3]	O6 - N4	2.89	N1 - N3	2.76	N2 - O2	2.59
6	G-----C	[3]	O6 - N4	2.64	N1 - N3	2.84	N2 - O2	2.87
7	G-----C	[3]	O6 - N4	2.57	N1 - N3	2.67	N2 - O2	2.83
8	T-----A	[2]	N3 - N1	2.51	O4 - N6	2.90		
9	C-----G	[3]	O2 - N2	2.38	N3 - N1	2.64	N4 - O6	2.83

```


10 C-----G [3] O2 - N2 2.66 N3 - N1 2.74 N4 - O6 2.78
11 T-----A [2] N3 - N1 2.74 O4 - N6 2.77
12 A-----T [2] N6 - O4 2.67 N1 - N3 2.61
13 T-----A [2] N3 - N1 2.60 O4 - N6 2.60
14 A-----T [2] N6 - O4 3.52 N1 - N3 2.86
15 G-----C [3] O6 - N4 3.43 N1 - N3 2.94 N2 - O2 2.60
16 A-----T [2] N6 - O4 3.13 N1 - N3 2.91
17 A-----T [2] N6 - O4 2.91 N1 - N3 2.63
18 C-----G [3] O2 - N2 2.47 N3 - N1 2.56 N4 - O6 2.57
19 T-----A [2] N3 - N1 2.65 O4 - N6 2.66

```

\*\*\*\*\*

Overlap area in Angstrom<sup>2</sup> between polygons defined by atoms on successive bases. Polygons projected in the mean plane of the designed base-pair step.

Values in parentheses measure the overlap of base ring atoms only. Those outside parentheses include exocyclic atoms on the ring. Intra- and inter-strand overlap is designated according to the following diagram:



step	i1-i2	i1-j2	j1-i2	j1-j2	sum
1 GG/CC	4.01( 2.20)	0.00( 0.00)	0.03( 0.00)	0.81( 0.00)	4.85( 2.20)
2 GG/CC	3.97( 2.70)	0.00( 0.00)	0.19( 0.00)	0.00( 0.00)	4.16( 2.70)
3 GG/CC	3.18( 1.65)	0.00( 0.00)	1.16( 0.00)	0.00( 0.00)	4.34( 1.65)
4 GG/CC	3.36( 1.96)	0.00( 0.00)	0.56( 0.00)	0.00( 0.00)	3.93( 1.96)
5 GG/CC	4.26( 2.90)	0.00( 0.00)	0.43( 0.00)	0.00( 0.00)	4.69( 2.90)
6 GG/CC	3.30( 1.83)	0.00( 0.00)	0.61( 0.00)	0.00( 0.00)	3.91( 1.83)
7 GT/AC	6.08( 1.72)	0.00( 0.00)	0.00( 0.00)	4.79( 3.35)	10.87( 5.06)
8 TC/GA	5.10( 0.61)	0.00( 0.00)	0.00( 0.00)	4.69( 1.63)	9.78( 2.25)
9 CC/GG	1.18( 0.00)	0.00( 0.00)	0.00( 0.00)	8.23( 5.73)	9.41( 5.73)
10 CT/AG	7.29( 1.31)	0.00( 0.00)	0.00( 0.00)	2.86( 1.70)	10.14( 3.01)
11 TA/TA	2.56( 0.00)	0.00( 0.00)	0.00( 0.00)	1.53( 0.00)	4.08( 0.00)
12 AT/AT	5.73( 1.71)	0.00( 0.00)	0.00( 0.00)	6.31( 2.41)	12.05( 4.12)
13 TA/TA	1.76( 0.00)	0.00( 0.00)	0.00( 0.00)	2.27( 0.00)	4.03( 0.00)
14 AG/CT	6.78( 4.02)	0.00( 0.00)	0.00( 0.00)	4.40( 0.27)	11.18( 4.29)
15 GA/TC	3.83( 0.87)	0.00( 0.00)	0.00( 0.00)	4.69( 0.53)	8.52( 1.40)
16 AA/TT	4.06( 2.70)	0.00( 0.00)	0.00( 0.00)	5.91( 0.32)	9.97( 3.02)
17 AC/GT	5.32( 4.25)	0.00( 0.00)	0.00( 0.00)	7.39( 2.57)	12.70( 6.81)
18 CT/AG	4.16( 0.08)	0.00( 0.00)	0.18( 0.00)	3.04( 2.60)	7.39( 2.68)

\*\*\*\*\*

Origin (Ox, Oy, Oz) and mean normal vector (Nx, Ny, Nz) of each base-pair in the coordinate system of the given structure

bp	Ox	Oy	Oz	Nx	Ny	Nz
1 G-C	-0.32	155.89	29.45	0.62	-0.76	-0.20
2 G-C	2.94	154.00	28.78	0.70	-0.69	-0.15

3	G-C	5.94	152.64	29.63	0.62	-0.79	-0.01
4	G-C	8.66	150.82	31.66	0.55	-0.83	0.07
5	G-C	10.24	148.16	33.94	0.42	-0.90	0.15
6	G-C	10.19	144.79	35.97	0.30	-0.94	0.17
7	G-C	9.09	141.17	37.41	0.14	-0.99	0.07
8	T-A	9.15	137.99	37.55	0.15	-0.99	-0.03
9	C-G	10.37	134.67	37.58	0.12	-0.99	-0.08
10	C-G	9.49	130.73	38.76	0.14	-0.99	0.04
11	T-A	10.52	127.69	38.52	0.26	-0.97	-0.03
12	A-T	10.13	124.46	40.11	0.15	-0.99	0.01
13	T-A	10.83	121.09	39.68	0.18	-0.98	-0.05
14	A-T	9.50	117.55	38.32	0.19	-0.97	-0.17
15	G-C	8.98	114.42	36.82	-0.05	-1.00	-0.08
16	A-T	9.55	111.47	36.25	-0.05	-0.98	-0.17
17	A-T	9.56	108.56	35.71	0.04	-0.97	-0.24
18	C-G	8.48	104.79	34.55	-0.05	-0.97	-0.25
19	T-A	7.31	101.48	33.18	-0.03	-0.94	-0.33

\*\*\*\*\*

#### Local base-pair parameters

bp	Shear	Stretch	Stagger	Buckle	Propeller	Opening
1 G-C	-0.18	-0.27	0.58	8.57	-9.77	-8.26
2 G-C	0.16	-0.26	0.13	-5.72	-10.04	-3.52
3 G-C	-0.10	-0.25	0.42	-1.54	-7.63	-0.67
4 G-C	0.02	-0.33	0.08	-3.56	-6.03	1.74
5 G-C	-0.21	-0.24	-0.13	-3.78	-0.69	3.13
6 G-C	0.04	-0.22	-0.35	-8.83	-11.08	-3.03
7 G-C	-0.32	-0.40	-0.38	-16.06	-17.27	-3.53
8 T-A	0.43	-0.40	-0.05	-11.97	-14.80	12.03
9 C-G	0.52	-0.43	-0.34	-10.52	4.23	3.87
10 C-G	0.27	-0.33	0.52	-15.53	-4.19	-2.52
11 T-A	-0.18	-0.23	-0.13	-9.98	-5.91	-1.56
12 A-T	-0.57	-0.49	0.47	0.67	-7.27	-0.77
13 T-A	-0.34	-0.35	0.33	-3.99	-14.34	-5.83
14 A-T	-0.69	-0.02	-0.04	-2.98	-0.45	12.91
15 G-C	-0.51	-0.07	-0.61	3.47	-9.27	12.05
16 A-T	-0.66	-0.15	-0.12	12.32	-13.66	3.93
17 A-T	-0.41	-0.38	0.73	21.10	-23.76	-1.56
18 C-G	0.40	-0.55	0.33	-5.21	-10.24	-3.28
19 T-A	0.56	-0.49	0.46	-21.17	-8.34	-4.72

~~~~~

|      |       |       |      |       |       |      |
|------|-------|-------|------|-------|-------|------|
| ave. | -0.09 | -0.31 | 0.10 | -3.93 | -8.97 | 0.55 |
| s.d. | 0.40  | 0.14  | 0.38 | 10.22 | 6.42  | 6.13 |

\*\*\*\*\*

#### Local base-pair step parameters

| step    | Shift | Slide | Rise | Tilt  | Roll  | Twist |
|---------|-------|-------|------|-------|-------|-------|
| 1 GG/CC | 0.19  | -1.13 | 3.65 | 5.58  | 3.88  | 39.73 |
| 2 GG/CC | 0.43  | -1.67 | 2.94 | -3.14 | 10.32 | 29.99 |
| 3 GG/CC | 0.17  | -2.24 | 3.13 | 1.31  | 6.72  | 31.31 |
| 4 GG/CC | 0.08  | -1.92 | 3.33 | 3.67  | 8.74  | 27.62 |
| 5 GG/CC | 0.50  | -1.91 | 3.41 | 4.23  | 5.60  | 31.56 |
| 6 GG/CC | 0.18  | -2.14 | 3.43 | 4.68  | 10.29 | 27.30 |
| 7 GT/AC | -0.07 | -0.42 | 3.15 | -0.69 | 5.91  | 35.56 |
| 8 TC/GA | 0.25  | 0.76  | 3.46 | 3.31  | 0.88  | 37.83 |



|       |       |       |       |      |       |       |       |
|-------|-------|-------|-------|------|-------|-------|-------|
| 9     | CC/GG | -1.73 | 0.69  | 3.77 | -7.18 | -1.10 | 31.39 |
| 10    | CT/AG | 0.35  | -0.32 | 3.19 | 5.36  | 6.16  | 26.98 |
| 11    | TA/TA | 0.41  | 1.89  | 3.06 | -6.05 | -2.74 | 45.60 |
| 12    | AT/AT | -0.34 | -0.18 | 3.44 | 1.73  | -3.42 | 31.69 |
| 13    | TA/TA | 0.29  | 2.18  | 3.37 | -2.75 | -6.21 | 49.54 |
| 14    | AG/CT | 0.33  | 1.29  | 3.25 | -0.15 | 15.17 | 23.41 |
| 15    | GA/TC | -0.68 | 0.30  | 2.96 | -4.96 | 0.68  | 35.87 |
| 16    | AA/TT | -0.06 | -0.03 | 2.97 | -6.65 | -0.95 | 34.68 |
| 17    | AC/GT | 0.37  | -1.03 | 3.93 | 4.89  | 1.28  | 38.27 |
| 18    | CT/AG | -0.19 | -1.07 | 3.61 | 1.91  | 4.78  | 37.16 |
| ~~~~~ |       |       |       |      |       |       |       |
|       | ave.  | 0.03  | -0.38 | 3.34 | 0.28  | 3.67  | 34.19 |
|       | s.d.  | 0.53  | 1.37  | 0.28 | 4.40  | 5.57  | 6.62  |

\*\*\*\*\*

Local base-pair helical parameters

| step  | X-disp | Y-disp | h-Rise | Incl. | Tip   | h-Twist |       |
|-------|--------|--------|--------|-------|-------|---------|-------|
| 1     | GG/CC  | -2.13  | 0.43   | 3.53  | 5.66  | -8.14   | 40.28 |
| 2     | GG/CC  | -4.56  | -1.25  | 2.20  | 19.17 | 5.83    | 31.83 |
| 3     | GG/CC  | -5.13  | -0.10  | 2.61  | 12.26 | -2.38   | 32.03 |
| 4     | GG/CC  | -5.58  | 0.59   | 2.60  | 17.66 | -7.41   | 29.17 |
| 5     | GG/CC  | -4.42  | -0.14  | 3.07  | 10.15 | -7.65   | 32.31 |
| 6     | GG/CC  | -6.26  | 0.59   | 2.47  | 20.70 | -9.43   | 29.51 |
| 7     | GT/AC  | -1.48  | 0.03   | 3.05  | 9.60  | 1.12    | 36.04 |
| 8     | TC/GA  | 1.05   | 0.06   | 3.48  | 1.35  | -5.09   | 37.98 |
| 9     | CC/GG  | 1.48   | 1.57   | 4.03  | -2.00 | 13.05   | 32.20 |
| 10    | CT/AG  | -2.07  | 0.50   | 3.05  | 12.84 | -11.16  | 28.17 |
| 11    | TA/TA  | 2.64   | -1.00  | 2.88  | -3.51 | 7.75    | 46.05 |
| 12    | AT/AT  | 0.32   | 0.95   | 3.42  | -6.23 | -3.15   | 31.91 |
| 13    | TA/TA  | 3.03   | -0.54  | 3.07  | -7.37 | 3.26    | 49.97 |
| 14    | AG/CT  | -1.36  | -0.72  | 3.43  | 33.28 | 0.33    | 27.84 |
| 15    | GA/TC  | 0.40   | 0.46   | 3.03  | 1.09  | 8.01    | 36.20 |
| 16    | AA/TT  | 0.09   | -0.80  | 2.93  | -1.57 | 11.03   | 35.30 |
| 17    | AC/GT  | -1.75  | 0.16   | 3.92  | 1.95  | -7.42   | 38.59 |
| 18    | CT/AG  | -2.34  | 0.56   | 3.44  | 7.45  | -2.98   | 37.50 |
| ~~~~~ |        |        |        |       |       |         |       |
|       | ave.   | -1.56  | 0.07   | 3.12  | 7.36  | -0.80   | 35.16 |
|       | s.d.   | 2.81   | 0.73   | 0.48  | 10.73 | 7.47    | 5.99  |

\*\*\*\*\*

Structure classification:

This is a right-handed nucleic acid structure

\*\*\*\*\*

lambda: virtual angle between C1'-YN1 or C1'-RN9 glycosidic bonds and the  
base-pair C1'-C1' line

C1'-C1': distance between C1' atoms for each base-pair  
RN9-YN1: distance between RN9-YN1 atoms for each base-pair  
RC8-YC6: distance between RC8-YC6 atoms for each base-pair

| bp | lambda(I) | lambda(II) | C1'-C1' | RN9-YN1 | RC8-YC6 |     |
|----|-----------|------------|---------|---------|---------|-----|
| 1  | G-C       | 51.0       | 58.7    | 10.7    | 9.0     | 9.8 |
| 2  | G-C       | 58.2       | 48.2    | 10.6    | 8.9     | 9.7 |
| 3  | G-C       | 62.1       | 54.4    | 10.3    | 8.8     | 9.8 |

|    |     |      |      |      |     |     |
|----|-----|------|------|------|-----|-----|
| 4  | G-C | 51.7 | 52.9 | 10.5 | 8.7 | 9.7 |
| 5  | G-C | 51.6 | 55.9 | 10.4 | 8.7 | 9.8 |
| 6  | G-C | 54.2 | 60.3 | 10.5 | 8.9 | 9.8 |
| 7  | G-C | 62.5 | 57.0 | 10.2 | 8.7 | 9.5 |
| 8  | T-A | 57.3 | 53.0 | 9.9  | 8.3 | 9.5 |
| 9  | C-G | 52.6 | 54.7 | 10.3 | 8.6 | 9.6 |
| 10 | C-G | 49.3 | 52.7 | 10.6 | 8.8 | 9.6 |
| 11 | T-A | 56.0 | 57.1 | 10.4 | 8.8 | 9.7 |
| 12 | A-T | 54.9 | 59.9 | 10.1 | 8.6 | 9.5 |
| 13 | T-A | 55.0 | 50.7 | 10.5 | 8.8 | 9.7 |
| 14 | A-T | 49.2 | 62.3 | 10.4 | 8.7 | 9.9 |
| 15 | G-C | 59.0 | 60.5 | 10.2 | 8.8 | 9.9 |
| 16 | A-T | 53.9 | 62.2 | 10.3 | 8.8 | 9.8 |
| 17 | A-T | 45.7 | 51.3 | 10.5 | 8.5 | 9.5 |
| 18 | C-G | 57.3 | 57.8 | 10.2 | 8.7 | 9.5 |
| 19 | T-A | 46.6 | 45.9 | 10.4 | 8.5 | 9.3 |

\*\*\*\*\*

Classification of each dinucleotide step in a right-handed nucleic acid structure: A-like; B-like; TA-like; intermediate of A and B, or other cases

| step     | Xp    | Yp   | Zp    | XpH   | YpH  | ZpH   | Form |
|----------|-------|------|-------|-------|------|-------|------|
| 1 GG/CC  | -2.03 | 8.22 | 1.96  | -4.14 | 8.00 | 2.77  | A    |
| 2 GG/CC  | -2.00 | 8.35 | 2.44  | -6.36 | 7.15 | 4.91  | A    |
| 3 GG/CC  | -1.96 | 8.21 | 2.78  | -6.89 | 7.47 | 4.41  | A    |
| 4 GG/CC  | -1.61 | 8.45 | 2.50  | -6.98 | 7.34 | 4.87  | A    |
| 5 GG/CC  | -1.44 | 8.13 | 2.57  | -5.65 | 7.57 | 3.94  | A    |
| 6 GG/CC  | -2.30 | 8.54 | 1.72  | -8.31 | 7.41 | 4.67  | A    |
| 7 GT/AC  | -3.10 | 8.76 | 0.52  | -4.51 | 8.57 | 1.89  |      |
| 8 TC/GA  | -3.37 | 8.71 | -0.37 | -2.26 | 8.72 | -0.19 | B    |
| 9 CC/GG  | -2.98 | 8.85 | -0.08 | -1.58 | 8.84 | -0.56 | B    |
| 10 CT/AG | -3.21 | 8.87 | -0.51 | -4.98 | 8.79 | 1.29  | B    |
| 11 TA/TA | -2.15 | 8.08 | -0.04 | 0.28  | 8.06 | -0.45 | B    |
| 12 AT/AT | -3.22 | 8.85 | -0.34 | -2.89 | 8.76 | -1.29 | B    |
| 13 TA/TA | -2.31 | 8.32 | -0.37 | 0.44  | 8.22 | -1.32 | B    |
| 14 AG/CT | -3.69 | 8.49 | -0.96 | -5.02 | 7.68 | 3.76  | B    |
| 15 GA/TC | -3.87 | 8.83 | -1.10 | -3.34 | 8.85 | -0.88 | B    |
| 16 AA/TT | -3.52 | 8.81 | 0.03  | -3.19 | 8.81 | -0.12 | B    |
| 17 AC/GT | -3.62 | 9.14 | 0.44  | -5.27 | 9.12 | 0.70  | B    |
| 18 CT/AG | -2.81 | 9.09 | 0.83  | -5.00 | 8.91 | 1.98  |      |

\*\*\*\*\*

Minor and major groove widths: direct P-P distances and refined P-P distances which take into account the directions of the sugar-phosphate backbones

(Subtract 5.8 Angstrom from the values to take account of the vdw radii of the phosphate groups, and for comparison with FreeHelix and Curves.)

Ref: M. A. El Hassan and C. R. Calladine (1998). ``Two Distinct Modes of Protein-induced Bending in DNA.'' J. Mol. Biol., v282, pp331-343.

|   | Minor Groove |         | Major Groove |         |
|---|--------------|---------|--------------|---------|
|   | P-P          | Refined | P-P          | Refined |
| 1 | GG/CC        | ---     | ---          | ---     |
| 2 | GG/CC        | ---     | ---          | ---     |

|    |       |      |      |      |      |
|----|-------|------|------|------|------|
| 3  | GG/CC | 16.3 | ---  | 17.1 | ---  |
| 4  | GG/CC | 17.0 | 16.0 | 20.4 | 18.6 |
| 5  | GG/CC | 16.0 | 15.3 | 20.8 | 18.5 |
| 6  | GG/CC | 14.7 | 13.8 | 18.6 | 18.0 |
| 7  | GT/AC | 13.9 | 13.2 | 17.9 | 17.9 |
| 8  | TC/GA | 12.8 | 12.5 | 16.4 | 16.2 |
| 9  | CC/GG | 13.0 | 13.0 | 15.9 | 15.9 |
| 10 | CT/AG | 13.1 | 13.1 | 17.3 | 17.2 |
| 11 | TA/TA | 11.6 | 11.6 | 15.8 | 15.5 |
| 12 | AT/AT | 10.5 | 10.5 | 17.6 | 16.9 |
| 13 | TA/TA | 11.8 | 11.8 | 17.2 | 16.6 |
| 14 | AG/CT | 14.0 | 14.0 | 16.8 | 16.7 |
| 15 | GA/TC | 13.7 | 13.6 | 16.1 | 15.7 |
| 16 | AA/TT | 12.2 | ---  | 19.2 | ---  |
| 17 | AC/GT | ---  | ---  | ---  | ---  |
| 18 | CT/AG | ---  | ---  | ---  | ---  |

\*\*\*\*\*

Global linear helical axis defined by equivalent C1' and RN9/YN1 atom pairs  
Deviation from regular linear helix: 2.85(0.76)

\*\*\*\*\*

Main chain and chi torsion angles:

Note: alpha: O3'(i-1)-P-O5'-C5'  
beta: P-O5'-C5'-C4'  
gamma: O5'-C5'-C4'-C3'  
delta: C5'-C4'-C3'-O3'  
epsilon: C4'-C3'-O3'-P(i+1)  
zeta: C3'-O3'-P(i+1)-O5'(i+1)

chi for pyrimidines(Y): O4'-C1'-N1-C2

chi for purines(R): O4'-C1'-N9-C4

Strand I

| base | alpha  | beta   | gamma  | delta | epsilon | zeta   | chi    |
|------|--------|--------|--------|-------|---------|--------|--------|
| 1 G  | ---    | -170.2 | 73.1   | 82.0  | -152.8  | -70.0  | -172.9 |
| 2 G  | -79.5  | -178.4 | 49.9   | 90.6  | -149.4  | -67.0  | -162.6 |
| 3 G  | -78.2  | 169.8  | 57.7   | 73.5  | -163.3  | -64.3  | -172.9 |
| 4 G  | -92.9  | -177.5 | 71.5   | 76.5  | -140.6  | -93.3  | -176.2 |
| 5 G  | -59.6  | 158.4  | 55.7   | 67.7  | -155.6  | -61.5  | -168.2 |
| 6 G  | -71.0  | 161.4  | 62.0   | 73.1  | -156.0  | -88.8  | -177.2 |
| 7 G  | 123.5  | -165.1 | -137.7 | 125.7 | -147.4  | -82.7  | -164.8 |
| 8 T  | -176.7 | -142.6 | 149.5  | 101.5 | -148.3  | -61.2  | -153.5 |
| 9 C  | -58.8  | -174.1 | 21.1   | 159.4 | -110.7  | 171.2  | -79.4  |
| 10 C | -92.0  | 143.7  | 68.4   | 154.1 | -156.8  | -107.9 | -122.7 |
| 11 T | -35.9  | -174.3 | -6.0   | 169.1 | -117.1  | 167.1  | -69.6  |
| 12 A | -93.0  | 148.2  | 66.1   | 161.4 | -120.5  | -151.0 | -109.0 |
| 13 T | -33.2  | 120.8  | 38.5   | 116.9 | -123.5  | -172.6 | -123.3 |
| 14 A | -57.0  | 149.4  | 48.3   | 158.9 | -164.6  | -84.7  | -94.1  |
| 15 G | -82.2  | -176.7 | 40.0   | 154.1 | -96.9   | 159.0  | -69.5  |
| 16 A | -75.0  | 135.8  | 50.2   | 121.8 | -147.3  | -154.9 | -112.8 |
| 17 A | 6.5    | 98.0   | 45.3   | 80.0  | 177.5   | -81.0  | -136.9 |
| 18 C | 105.5  | -173.8 | -165.4 | 81.5  | 83.6    | 54.5   | -153.6 |
| 19 T | -142.8 | 170.5  | 85.0   | 80.9  | ---     | ---    | -167.3 |

## Strand II

| base | alpha | beta   | gamma  | delta | epsilon | zeta   | chi    |
|------|-------|--------|--------|-------|---------|--------|--------|
| 1 C  | -72.4 | -176.5 | 67.2   | 128.1 | ---     | ---    | -129.0 |
| 2 C  | -43.5 | 172.3  | 43.4   | 148.7 | -173.4  | -112.6 | -122.5 |
| 3 C  | 142.3 | -163.6 | -168.5 | 97.5  | -136.0  | -81.8  | -170.0 |
| 4 C  | -60.4 | 174.5  | 48.5   | 83.2  | -179.6  | -85.3  | -166.2 |
| 5 C  | -43.3 | 172.4  | 34.5   | 79.4  | -151.6  | -66.2  | -155.8 |
| 6 C  | 107.3 | -161.6 | -159.1 | 89.0  | -141.1  | -88.0  | -165.4 |
| 7 C  | 161.3 | -116.0 | 148.8  | 88.1  | -155.2  | -103.4 | -151.3 |
| 8 A  | -78.6 | 126.0  | 59.1   | 86.8  | 175.4   | -81.2  | -135.3 |
| 9 G  | -69.2 | 179.3  | 44.6   | 141.5 | -109.5  | -178.9 | -92.7  |
| 10 G | -99.0 | -157.7 | 76.8   | 111.6 | -174.7  | -74.4  | -108.6 |
| 11 A | -64.6 | 168.0  | 47.6   | 139.1 | 153.4   | -72.2  | -88.7  |
| 12 T | -82.1 | -174.6 | 63.1   | 146.1 | -132.6  | -163.4 | -112.7 |
| 13 A | -70.6 | 130.0  | 55.8   | 140.4 | 175.2   | -94.7  | -97.2  |
| 14 T | -82.3 | -167.0 | 69.2   | 156.5 | -102.6  | 171.9  | -88.1  |
| 15 C | -92.1 | 153.6  | 77.9   | 78.5  | -170.8  | -83.1  | -129.7 |
| 16 T | 38.9  | 90.6   | 38.3   | 104.6 | -156.8  | -86.5  | -119.6 |
| 17 T | 154.2 | -128.5 | 165.8  | 80.1  | -145.5  | -126.3 | -179.1 |
| 18 G | 98.2  | -172.9 | -121.4 | 136.6 | 176.7   | -104.8 | -131.4 |
| 19 A | ---   | ---    | -157.3 | 94.7  | -173.6  | -100.2 | -137.4 |

\*\*\*\*\*

## Sugar conformational parameters:

Note: v0: C4'-O4'-C1'-C2'

v1: O4'-C1'-C2'-C3'

v2: C1'-C2'-C3'-C4'

v3: C2'-C3'-C4'-O4'

v4: C3'-C4'-O4'-C1'

tm: amplitude of pseudorotation of the sugar ring

P: phase angle of pseudorotation of the sugar ring

## Strand I

| base | v0    | v1    | v2    | v3    | v4    | tm   | P     | Puckering |
|------|-------|-------|-------|-------|-------|------|-------|-----------|
| 1 G  | 1.7   | -26.8 | 40.6  | -40.2 | 24.6  | 42.3 | 16.2  | C3'-endo  |
| 2 G  | 11.7  | -33.7 | 42.2  | -36.2 | 15.7  | 42.2 | 2.8   | C3'-endo  |
| 3 G  | -5.5  | -25.6 | 45.2  | -50.0 | 35.1  | 49.9 | 25.1  | C3'-endo  |
| 4 G  | -22.3 | -11.0 | 38.7  | -52.3 | 46.7  | 52.7 | 42.8  | C4'-exo   |
| 5 G  | -5.4  | -28.9 | 50.6  | -55.1 | 38.0  | 55.4 | 24.1  | C3'-endo  |
| 6 G  | -4.0  | -27.0 | 47.5  | -51.1 | 34.3  | 51.6 | 23.1  | C3'-endo  |
| 7 G  | 39.7  | -32.4 | 14.6  | 7.4   | -28.7 | 38.0 | 292.5 | C1'-endo  |
| 8 T  | -21.3 | 7.8   | 7.4   | -19.8 | 25.5  | 25.3 | 73.1  | O4'-endo  |
| 9 C  | -22.0 | 44.1  | -47.8 | 36.3  | -9.4  | 48.3 | 172.1 | C2'-endo  |
| 10 C | -32.4 | 50.5  | -49.1 | 31.8  | 0.3   | 51.9 | 161.2 | C2'-endo  |
| 11 T | -18.0 | 43.1  | -48.9 | 41.5  | -15.1 | 48.9 | 178.3 | C2'-endo  |
| 12 A | -15.5 | 35.7  | -41.9 | 34.2  | -11.8 | 41.9 | 177.7 | C2'-endo  |
| 13 T | -51.6 | 50.3  | -29.7 | 0.8   | 31.5  | 52.3 | 124.6 | C1'-exo   |
| 14 A | -7.3  | 30.1  | -40.2 | 36.9  | -19.3 | 40.6 | 188.6 | C3'-exo   |
| 15 G | -23.0 | 41.8  | -42.1 | 30.7  | -5.1  | 43.2 | 167.4 | C2'-endo  |
| 16 A | -36.4 | 38.4  | -26.2 | 6.3   | 18.6  | 38.6 | 132.8 | C1'-exo   |
| 17 A | -55.8 | 34.1  | -0.4  | -32.8 | 55.7  | 58.0 | 90.3  | O4'-endo  |

|      |       |       |      |       |      |      |      |          |
|------|-------|-------|------|-------|------|------|------|----------|
| 18 C | -6.0  | -19.6 | 36.1 | -40.6 | 29.0 | 40.4 | 26.8 | C3'-endo |
| 19 T | -41.2 | 10.7  | 21.0 | -45.1 | 52.9 | 53.0 | 66.6 | C4'-exo  |

# Strand II

| base | v0    | v1    | v2    | v3    | v4    | tm   | P     | Puckering |
|------|-------|-------|-------|-------|-------|------|-------|-----------|
| 1 C  | -21.4 | 29.0  | -26.2 | 14.7  | 4.0   | 29.2 | 153.8 | C2'-endo  |
| 2 C  | -35.3 | 47.2  | -41.3 | 22.0  | 8.1   | 46.9 | 151.6 | C2'-endo  |
| 3 C  | 8.7   | -24.7 | 30.5  | -26.9 | 11.4  | 30.5 | 3.0   | C3'-endo  |
| 4 C  | -18.5 | -9.5  | 31.4  | -43.1 | 39.0  | 43.1 | 43.3  | C4'-exo   |
| 5 C  | -1.5  | -21.6 | 36.9  | -38.7 | 24.3  | 39.4 | 20.7  | C3'-endo  |
| 6 C  | 4.2   | -27.5 | 39.4  | -38.2 | 21.3  | 40.4 | 12.9  | C3'-endo  |
| 7 C  | -44.0 | 21.3  | 6.9   | -32.2 | 47.4  | 47.6 | 81.7  | O4'-endo  |
| 8 A  | -18.2 | -4.7  | 24.3  | -35.3 | 33.2  | 36.0 | 47.6  | C4'-exo   |
| 9 G  | -39.6 | 49.4  | -40.7 | 19.0  | 12.6  | 48.7 | 146.6 | C2'-endo  |
| 10 G | -33.8 | 28.7  | -14.6 | -4.5  | 23.9  | 32.9 | 116.3 | C1'-exo   |
| 11 A | 13.5  | 4.9   | -19.9 | 28.4  | -27.3 | 28.9 | 226.4 | C4'-endo  |
| 12 T | -39.1 | 48.1  | -38.2 | 17.2  | 13.5  | 46.9 | 144.6 | C2'-endo  |
| 13 A | -27.6 | 39.7  | -37.0 | 22.0  | 3.3   | 40.2 | 156.9 | C2'-endo  |
| 14 T | -10.5 | 30.5  | -37.5 | 33.0  | -14.4 | 37.5 | 183.2 | C3'-exo   |
| 15 C | -29.8 | 3.6   | 22.2  | -40.0 | 44.0  | 44.2 | 59.8  | C4'-exo   |
| 16 T | -37.5 | 34.3  | -18.7 | -2.4  | 25.0  | 37.3 | 120.2 | C1'-exo   |
| 17 T | -28.6 | 1.7   | 24.4  | -41.8 | 43.4  | 44.7 | 57.0  | C4'-exo   |
| 18 G | -3.0  | 15.4  | -21.2 | 20.1  | -11.0 | 21.6 | 190.9 | C3'-exo   |
| 19 A | -44.0 | 27.3  | -1.9  | -23.3 | 41.5  | 44.3 | 92.5  | O4'-endo  |

\*\*\*\*\*

Same strand P--P and C1'--C1' virtual bond distances

| Strand I |      |          | Strand II |      |          |
|----------|------|----------|-----------|------|----------|
| base     | P--P | C1'--C1' | base      | P--P | C1'--C1' |
| 1 G/G    | 5.8  | 5.6      | 1 C/C     | 6.6  | 5.3      |
| 2 G/G    | 5.7  | 5.3      | 2 C/C     | 6.3  | 5.6      |
| 3 G/G    | 5.8  | 5.4      | 3 C/C     | 6.3  | 5.5      |
| 4 G/G    | 5.4  | 5.7      | 4 C/C     | 5.4  | 5.4      |
| 5 G/G    | 5.9  | 5.1      | 5 C/C     | 6.6  | 5.6      |
| 6 G/G    | 5.8  | 5.7      | 6 C/C     | 6.7  | 5.4      |
| 7 G/T    | 6.7  | 5.3      | 7 C/A     | 6.7  | 4.6      |
| 8 T/C    | 6.9  | 5.2      | 8 A/G     | 6.1  | 4.8      |
| 9 C/C    | 6.4  | 5.4      | 9 G/G     | 6.7  | 4.4      |
| 10 C/T   | 7.2  | 4.9      | 10 G/A    | 7.0  | 4.7      |
| 11 T/A   | 6.2  | 5.0      | 11 A/T    | 6.5  | 5.9      |
| 12 A/T   | 7.1  | 4.3      | 12 T/A    | 7.0  | 4.5      |
| 13 T/A   | 6.3  | 5.7      | 13 A/T    | 6.7  | 5.1      |
| 14 A/G   | 7.1  | 4.4      | 14 T/C    | 6.6  | 5.6      |
| 15 G/A   | 6.6  | 4.9      | 15 C/T    | 6.3  | 4.7      |
| 16 A/A   | 6.7  | 4.4      | 16 T/T    | 6.6  | 5.2      |
| 17 A/C   | 6.5  | 4.9      | 17 T/G    | 7.1  | 4.5      |
| 18 C/T   | 6.7  | 4.9      | 18 G/A    | ---  | 5.0      |

\*\*\*\*\*

Helix radius (radial displacement of P, O4', and C1' atoms in local helix frame of each dimer)

| Strand I |   |     | Strand II |     |     |
|----------|---|-----|-----------|-----|-----|
| step     | P | O4' | P         | O4' | C1' |

|    |       |      |      |      |      |      |     |
|----|-------|------|------|------|------|------|-----|
| 1  | GG/CC | 8.9  | 8.1  | 7.2  | 9.1  | 7.2  | 6.5 |
| 2  | GG/CC | 8.0  | 8.3  | 7.8  | 11.1 | 10.1 | 9.4 |
| 3  | GG/CC | 9.9  | 9.6  | 9.1  | 10.4 | 9.6  | 9.1 |
| 4  | GG/CC | 10.7 | 10.4 | 10.0 | 9.5  | 9.4  | 9.0 |
| 5  | GG/CC | 9.4  | 9.1  | 8.6  | 9.5  | 9.2  | 8.6 |
| 6  | GG/CC | 10.9 | 10.9 | 10.4 | 11.4 | 9.8  | 9.4 |
| 7  | GT/AC | 9.4  | 7.4  | 6.5  | 10.0 | 7.3  | 6.5 |
| 8  | TC/GA | 9.3  | 6.0  | 5.3  | 8.7  | 6.1  | 5.3 |
| 9  | CC/GG | 10.3 | 7.3  | 6.9  | 7.8  | 4.3  | 3.8 |
| 10 | CT/AG | 10.8 | 7.7  | 7.3  | 9.4  | 6.9  | 6.1 |
| 11 | TA/TA | 7.2  | 4.3  | 4.1  | 9.0  | 6.6  | 6.1 |
| 12 | AT/AT | 10.3 | 6.9  | 6.4  | 8.1  | 5.0  | 4.6 |
| 13 | TA/TA | 7.4  | 5.0  | 4.6  | 9.0  | 6.0  | 5.8 |
| 14 | AG/CT | 9.5  | 6.8  | 6.0  | 9.0  | 7.6  | 6.6 |
| 15 | GA/TC | 10.0 | 6.8  | 6.2  | 9.0  | 5.7  | 5.1 |
| 16 | AA/TT | 9.0  | 6.0  | 5.1  | 9.8  | 7.0  | 6.2 |
| 17 | AC/GT | 11.0 | 7.5  | 6.8  | 10.1 | 7.3  | 6.4 |
| 18 | CT/AG | 10.5 | 7.9  | 7.2  | 10.0 | 7.2  | 6.6 |

\*\*\*\*\*

Position (Px, Py, Pz) and local helical axis vector (Hx, Hy, Hz)  
for each dinucleotide step

|    | bp    | Px    | Py     | Pz    | Hx    | Hy    | Hz    |
|----|-------|-------|--------|-------|-------|-------|-------|
| 1  | GG/CC | 1.94  | 155.38 | 31.00 | 0.62  | -0.78 | -0.03 |
| 2  | GG/CC | 2.47  | 151.13 | 32.67 | 0.43  | -0.85 | -0.30 |
| 3  | GG/CC | 3.81  | 149.58 | 33.38 | 0.45  | -0.88 | -0.11 |
| 4  | GG/CC | 4.52  | 147.45 | 33.83 | 0.35  | -0.92 | -0.17 |
| 5  | GG/CC | 6.63  | 145.33 | 32.97 | 0.33  | -0.94 | -0.05 |
| 6  | GG/CC | 5.82  | 142.28 | 32.01 | 0.41  | -0.89 | -0.20 |
| 7  | GT/AC | 9.07  | 139.52 | 36.07 | 0.30  | -0.95 | 0.03  |
| 8  | TC/GA | 9.17  | 136.12 | 38.33 | 0.20  | -0.97 | -0.11 |
| 9  | CC/GG | 8.08  | 132.84 | 37.25 | -0.09 | -1.00 | 0.03  |
| 10 | CT/AG | 11.95 | 129.88 | 38.43 | 0.34  | -0.91 | 0.25  |
| 11 | TA/TA | 7.96  | 125.84 | 38.27 | 0.16  | -0.98 | -0.14 |
| 12 | AT/AT | 11.32 | 123.00 | 39.49 | 0.27  | -0.96 | 0.02  |
| 13 | TA/TA | 11.22 | 119.89 | 36.48 | 0.31  | -0.95 | -0.09 |
| 14 | AG/CT | 7.77  | 116.25 | 37.52 | -0.12 | -0.78 | -0.61 |
| 15 | GA/TC | 9.57  | 112.91 | 37.03 | 0.08  | -0.99 | -0.12 |
| 16 | AA/TT | 10.14 | 110.11 | 35.51 | 0.11  | -0.99 | -0.06 |
| 17 | AC/GT | 8.61  | 107.25 | 33.63 | 0.01  | -0.93 | -0.36 |
| 18 | CT/AG | 8.50  | 103.79 | 31.77 | 0.09  | -0.96 | -0.27 |

## References

- Babcock, M. S., Pednault, E. P. D. & Olson, W. K. (1994). Nucleic acid structure analysis: Mathematics for local cartesian and helical structure parameters that are truly comparable between structures, *J. Mol. Biol.* **237**, 125–156.
- Calladine, C. R. & Drew, H. R. (1997). *Understanding DNA; The Molecule & How It Works*, 2nd edn, Academic Press, London.
- Chandrasekaran, R. & Arnott, S. (1989). The structures of DNA and RNA helices in oriented fibers, in W. Saenger (ed.), *Landolt-Börnstein Numerical Data and Functional Relationships in Science and Technology*, Vol. VII/1b, Springer-Verlag, pp. 31–170.
- Clowney, L., Jain, S. C., Srinivasan, A. R., Westbrook, J., Olson, W. K. & Berman, H. M. (1996). Geometric parameters in nucleic acids: Nitrogenous bases, *J. Am. Chem. Soc.* **118**, 509–518.
- Dickerson, R. E. (1998). DNA bending: The prevalence of kinkiness and the virtues of normality, *Nucl. Acids Res.* **26**, 1906–1926.
- Dickerson, R. E., Bansal, M., Calladine, C. R., Diekmann, S., Hunter, W. N., Kennard, O., Lavery, R., Nelson, H. C. M., Olson, W. K., Saenger, W., Shakked, Z., Sklenar, H., Soumpasis, D. M., Tung, C.-S., von Kitzing, E., Wang, A. H.-J. & Zhurkin, V. B. (1989). Definitions and nomenclature of nucleic acid structure parameters, *J. Mol. Biol.* **205**, 787–791.
- El Hassan, M. A. & Calladine, C. R. (1995). The assessment of the geometry of dinucleotide steps in double-helical DNA: A new local calculation scheme, *J. Mol. Biol.* **251**, 648–664.
- El Hassan, M. A. & Calladine, C. R. (1998). Two distinct modes of protein-induced bending in DNA, *J. Mol. Biol.* **282**, 331–343.
- Guzikevich-Guerstein, G. & Shakked, Z. (1996). A novel form of the DNA double helix imposed on the TATA-box by the TATA-binding protein, *Nature Struct. Biol.* **3**, 32–37.
- Horn, B. K. P. (1987). Closed-form solution of absolute orientation using unit quaternions, *J. Opt. Soc. Am. A* **4**(4), 629–642.
- Lavery, R. & Sklenar, H. (1989). Defining the structure of irregular nucleic acids: Conventions and principles, *J. Biomol. Struct. Dynam.* **6**, 655–667.
- Leslie, A. G. W., Arnott, S., Chandrasekaran, R., Birdsall, D. L. & Ratliff, R. L. (1980). Left-handed DNA helices, *Nature* **283**, 743–746.
- Lu, X. J., Babcock, M. S. & Olson, W. K. (1999). Mathematical overview of nucleic acid analysis programs, *J. Biomol. Struct. Dynam.* **16**, 833–843.
- Lu, X. J., El Hassan, M. A. & Hunter, C. A. (1997a). Structure and conformation of helical nucleic acids: Analysis program (SCHNAaP), *J. Mol. Biol.* **273**, 668–680.

- Lu, X. J., El Hassan, M. A. & Hunter, C. A. (1997b). Structure and conformation of helical nucleic acids: Rebuilding program (SCHNArP), *J. Mol. Biol.* **273**, 681–691.
- Lu, X. J. & Olson, W. K. (1999). Resolving the discrepancies among nucleic acid conformational analyses, *J. Mol. Biol.* **285**, 1563–1575.
- Lu, X. J., Shakked, Z. & Olson, W. K. (2000). A-dna conformational motifs in ligand-bound double helices, *J. Mol. Biol.* **300**, 819–840.
- Merritt, E. A. & Bacon, D. J. (1997). Raster3d: Photorealistic molecular graphics, *Methods in Enzymology* **277**, 505–524.
- Olson, W. K., Bansal, M., Burley, S. K., Dickerson, R. E., Gerstein, M., Harvey, S. C., Heinemann, X.-J. L. U., Neidle, S., Shakked, Z., Sklenar, H., Suzuki, M., Tung, C.-S., Westhof, E., Wolberger, C. & Berman, H. M. (2001). A standard reference frame for the description of nucleic acid base-pair geometry, *J. Mol. Biol.* **313**, 229–237.
- Shui, X., McFail-Isom, L., Hu, G. G. & Williams, L. D. (1998). The B-DNA dodecamer at high resolution reveals a spine of water on sodium, **37**, 8341–8355.
- van Pouderooyan, G., Ketting, G., Perrakis, R. F. & Sixma, T. K. (1997). Crystal structure of the specific DNA-binding domain of Tc3 transposase of *c. elegans* in complex with transposon DNA, *EMBO J.* **16**, 6044–6054.