
Supplementary Online Content

Sun X, Deng Y, Zhang Y, et al; for the BASIS Investigators. Balloon angioplasty vs aggressive medical management for symptomatic intracranial artery stenosis: the BASIS randomized clinical trial. JAMA.

Published online September 5, 2024. doi:10.1001/jama.2024.12829

Corrected online November 27, 2024.

This supplemental material has been provided by the authors to give readers additional information about their work.

Supplementary Appendix

Contents

List of participating sites and investigators in the BASIS trial	Page 2
Study Committee Members of the BASIS trial	Page 4
Study design and treatment allocation	Page 5
Distribution of 31 participating centers in 14 provinces	Page 6
On-site and online training arrangement of the BASIS trial	Page 7
Research Progress of BASIS	Page 8
Supplementary Figures S1-S6	Page 9-12
Figure S1. Flow chart of clinical-event adjudication committee (CEC) assessing and confirming endpoint events	Page 9
Figure S2. Enrollment progress of the BASIS trial by month	Page 10
Figure S3. Number of patients enrolled by center	Page 11
Figure S4. Distribution of 90-day mRS	Page 12
Figure S5. Distribution of 1-year mRS	Page 12
Supplementary Tables S1-S8	
Table S1. Demographic and baseline clinical characteristics	Page 13-26
Table S2. Measures of risk factors at baseline, 3-months and 1-year follow-up	Page 13
Table S3. The proportion of patients receiving antiplatelet and statin medicine at baseline, 3-months and 1-year follow-up	Page 15
Table S4. Details of balloon angioplasty	Page 16
Table S5. The number of patients and reason for revascularisation beyond 30 days through 1 year in the ITT population of the BASIS trial	Page 17
Table S6. Post-hoc analysis of center-effect adjustment	Page 18
Table S7. Post-hoc analysis of treatment interaction on the primary outcome regarding different sites	Page 19
Table S8. Study outcomes in the per-protocol population	Page 20
Table S9. Study outcomes in the as-treated population	Page 21
Table S10. Post-hoc analysis of the comparison of any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion or any ischemic stroke from the qualifying artery beyond 30 days through 1 year after enrollment between BA and AMM groups	Page 22
Table S11. Number of patients with other serious adverse events (by system organ class) up to 1-year visit	Page 23
Table S12. The proportion of patients meeting the proposed target of vascular risk factor management at 1-year follow-up	Page 24
	Page 25

➤ **List of participating sites and investigators in the BASIS trial**


BASIS site	Investigator
Beijing Tiantan Hospital, Capital Medical University	Zhongrong Miao
Beijing Tiantan Hospital, Capital Medical University	Yilong Wang
The Affiliated Hospital of Qingdao University	Yong Zhang
Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School	Yun Xu
Zhengzhou University People's Hospital	Tianxiao Li
The Second Affiliated Hospital of Guangzhou Medical University	Yunxiang Ji
Beijing Luhe Hospital, Capital Medical University	Xiaokun Geng
The First Hospital of Jilin University	Shouchun Wang
Qilu Hospital of Shandong University	Wei Wu
Hunan Provincial People's Hospital	Xiaoping Gao
The First Affiliated Hospital, Sun Yat-sen University	Shihui Xing
The Affiliated Hospital of Guizhou Medical University	Hua Yang
Shandong Provincial Hospital	Yifeng Du
General Hospital of The Northern Theater of The Chinese People's Liberation Army	Huisheng Chen
Guangdong Sanjiu Brain Hospital	Peiming Wang
Shenzhen Hospital of Southern Medical University	Yajie Liu
West China Hospital Sichuan University	Hongbo Zheng
Wuhan First Hospital	Wenhua Liu
The First Affiliated Hospital of Anhui Medical University	Weimin Yang
Jiangxi Provincial People's Hospital	Wenfeng Cao
Liaocheng People's Hospital	Guisheng Jiang
The First Affiliated Hospital of Xi'an Jiaotong University	Jianfeng Han
The First People's Hospital of Changzhou	Ya Peng
Chinese PLA General Hospital	Jun Wang
Beijing Chao-Yang Hospital, Capital Medical University	Yang Wang
Beijing Fengtai You Anmen Hospital	Shiyong Zhang
Dongfang Hospital Beijing University of Chinese Medicine	Gesheng Wang

BASIS site	Investigator
Beijing ShunYi Hospital	Liyu Wang
The Second Affiliated Hospital Of NanJing Medical University	Jin Wu
Guangdong Provincial Hospital of Traditional Chinese Medicine	Guifu Li
The Affiliated Hospital of Northwest University	Mingze Chang

➤ **Study Committee Members of the BASIS trial**


Cooperation team	Unit or Hospital name	Investigator Name
Sponsor	Beijing Tiantan Hospital, Capital Medical University	NA
CRO	Beijing Jingchengtong Medicine Technology Co., Ltd.	NA
Statistical analysis	Department of Biostatistics, Peking University Clinical Research Institute	NA
Imaging core lab	China National Clinical Research Center for Neurological Diseases, Tiantan Neuroimaging Center of Excellence	Jing Jing
	China National Clinical Research Center for Neurological Diseases, Tiantan Neuroimaging Center of Excellence	Nan Qi
CEC team	Tangdu Hospital of Air Force Medical University	Zhenwei Zhao
	Beijing Tiantan Hospital, Capital Medical University	Kehui Dong
	The First Affiliated Hospital of Harbin Medical University	Huaizhang Shi
DSMB	People's Liberation Army Air Force General Hospital	Jin Shi
	Strategic Support Force Specialty Medical Center	Yiling Cai
	China National Clinical Research Center for Neurological Diseases	Yuesong Pan
AE Committee	Beijing Tiantan Hospital, Capital Medical University	Xuan Sun
	Beijing Tiantan Hospital, Capital Medical University	Yiming Deng
	Beijing Tiantan Hospital, Capital Medical University	Ning Ma
Steering Committee	Beijing Tiantan Hospital, Capital Medical University	Zhongrong Miao
	Beijing Tiantan Hospital, Capital Medical University	Yilong Wang
	Beijing Tiantan Hospital, Capital Medical University	Yongjun Wang
	Department of Neurology, University of California Irvine, Irvine, California, USA	Wengui Yu
	Beijing Tiantan Hospital, Capital Medical University	Xingquan Zhao
	Neurovascular Imaging Research Core and Department of Neurology, University of California, USA	David S Liebeskind
	Department of Neurosurgery, University Hospitals Cleveland Medical Center/ Case Western Reserve University School of Medicine, Ohio, USA	Sepideh Amin-Hanjani

➤ Study design and treatment allocation



ICAS denotes intracranial atherosclerosis stenosis. DSA denotes digital subtraction angiography. TIA denotes transient ischemic attack. ICA denotes internal carotid artery. MCA denotes middle cerebral artery. VA denotes vertebral artery. mRS denotes modified Rankin Scale. AMM denotes aggressive medical management.

➤ **Distribution of 31 participating centers in 14 provinces**



➤ On-site and online training arrangement of the BASIS trial



➤ Research Progress of BASIS



First investigators meeting DSMB

2017/11/24



Leading Center Kick off meeting

2018/6/19



First Clinical Technique Review Meeting

2018/11/08



Regular quality control meeting

2020/10/15



Leading Center Ethical approval



First patient enrolled



First DSMB review meeting



Core lab meeting



First CEC review meeting

2021/05/30

Last patient enroll in

2021/11/20

2022/04/02

2022/09/07



Fourth CEC review meeting

2023/04/10

Complete all data cleaning, core laboratory analysis, and independent interpretation by CEC

2023/05/02

2023/06

Second CEC review meeting




Third CEC review meeting




All patients completed 1-year primary endpoint follow-up

Supplementary Results

- **Figure S1. Flow chart of clinical-event adjudication committee (CEC) assessing and confirming endpoint events**



➤ **Figure S2. Enrollment Progress of the BASIS trial by month**



➤ **Figure S3. Number of patients enrolled by each participating center**




Figure S4. Distribution of the modified Rankin Scale at 90 days

BA denotes balloon angioplasty. AMM denotes aggressive medical management.





Figure S5. Distribution of modified Rankin Scale at 1 year

BA denotes balloon angioplasty. AMM denotes aggressive medical management



➤ **Table S1. Demographic and baseline clinical characteristics of patients**

Characteristics	Balloon angioplasty group (n=249)	Aggressive medical management group (n=252)
Ethnicity, n (%)		
Han	245 (98.4)	249 (98.8)
Korean	0 (0)	1 (0.4)
Manchus	2 (0.8)	1 (0.4)
Mongol	2 (0.8)	0 (0)
Yi	0 (0.0)	1 (0.4)
Medical History, n (%)		
Coronary heart disease	29 (11.7)	28 (11.1)
ischemic stroke	50 (20.1)	55 (21.8)
ASITN/SIR, n (%)	1(1,1)	1 (1,1)
0-2	222 (89.2)	219 (86.9)
3-4	27 (10.8)	33 (13.1)
Lesion length, mm, median (IQR)	6.1 (5.0,8.0)	6.0 (4.7,7.8)
Reference-vessel diameter, mm	2.3 (2.0,2.8)	2.4 (2.1,2.9)
Minimal luminal diameter, mm	0.5 (0.3,0.6)	0.5 (0.4,0.7)
ischemic stroke, n (%)	215 (86.4)	208 (82.5)
Border zone infarct	84 (39.1)	85(40.9)
Non-zone infarct	131(61.9)	123(59.1)
eTICI before the procedure, n (%)		
0	0 (0)	0 (0)
1	3 (1.2)	3 (1.2)
2a	12 (4.8)	17 (6.8)
2b50	80 (32.1)	78 (31.0)
2b67	112 (45.0)	121 (48.0)
2c	39 (15.7)	31 (12.3)
3	3 (1.2)	2 (0.8)
Received medical therapy at qualifying event n (%)		
Statin	123 (49.4)	126 (50.0)
ACE inhibitor or angiotensin II receptor	29 (11.7)	19 (7.5)
Beta-Blocker	4 (1.6)	4 (1.6)
Calcium antagonist	78 (31.3)	43 (17.1)
Diuretic	1 (0.4)	1 (0.4)
Sulfonylureas	5 (2.0)	8 (3.2)
Biguanides	10 (4.0)	17 (6.8)
Glycoside inhibitor	16 (6.4)	23 (9.1)
Insulin	47 (19.0)	25 (10.0)

Data are n (%), or median (IQR). IQR denotes interquartile range. eTICI denotes expanded Thrombolysis in Cerebral Infarction. BA denotes balloon angioplasty. AMM denotes aggressive medical management.

➤ **Table S2. Measures of risk factors at baseline, 3-months and 1-year follow-up**

Characteristics	Balloon angioplasty group (n=249)			Aggressive medical management group (n=252)		
	Baseline	3 months	1 year	Baseline	3 months	1 year
No. Patients evaluated	249	245	248	252	245	246
BMI, kg/m ² , median (IQR)	25.4 (23.7,27.5)	25.4 (23.7,27.5)	25.2 (23.7,27.3)	25.4 (23.5,27.7)	25.4 (23.6,27.7)	25.6 (23.5,27.7)
No. Patients evaluated	249	248	248	252	252	246
Current smoking, n (%)	60 (24.1)	11 (4.4)	22 (8.9)	66 (26.2)	15 (6.0)	16 (6.5)
No. Patients evaluated	249	249	NA	252	252	NA
No. Patients evaluated	248	243	247	249	251	246
Activity level per week, h, median (IQR)	3.5 (2.5,7.0)	4.0 (3.0,7.0)	4.0 (2.5,7.0)	7.0 (3.5,7.0)	5.0 (3.5,7.0)	5.0 (3.0,7.0)
No. Patients evaluated	249	239	247	252	233	243
Blood pressure, mm Hg, median (IQR)						
Systolic	140.0 (130.0,154.0)	135.0 (130.0,138.0)	130.0 (125.0,139.0)	140.0 (130.0, 151.0)	135.0 (130.0,140.0)	130.0 (125.0,139.0)
Diastolic	83.0 (77.0,91.0)	80.0 (75.0,85.0)	80.0 (70.0,82.0)	84.0 (77.0, 92.0)	80.0 (75.0,85.0)	80.0 (75.0,85.0)
No. Patients evaluated	233	128	140	230	143	121
Cholesterol, mg/dl, median (IQR)	136.1 (115.2,163.2)	137.3(121.0,155.3)	134.8(115.6,150.4)	139.2 (119.9,161.3)	141.9 (119.5,157.4)	134.2(115.6,158.9)
No. Patients evaluated	232	119	140	226	137	120
High-density lipoprotein, mg/dl	41.8(35.2,49.1)	44.5 (38.3,51.4)	45.2 (38.3,53.9)	39.8 (34.8,48.0)	43.7 (37.9,50.7)	42.0 (34.4,48.5)
No. Patients evaluated	233	121	139	226	141	120
Low-density lipoprotein, mg/dl	72.7 (57.6,93.6)	73.1 (59.2,89.7)	70.8 (55.7,86.6)	76.0 (61.5,94.7)	75.4 (61.1,89.3)	69.0 (56.3,88.8)
No. Patients evaluated	150	88	65	132	107	66
Glycosylated hemoglobin, %	6.1 (5.6,7.0)	6.0 (5.7,6.7)	6.2 (5.8,7.4)	6.5 (5.8,7.8)	6.1 (5.6,6.9)	6.1 (5.6,7.1)

Data are n (%), or median (IQR). BMI denotes body-mass index. BA denotes balloon angioplasty. AMM denotes aggressive medical management. IQR denotes interquartile range.

➤ **Table S3. The proportion of patients receiving antiplatelet and statin medicine at baseline, 3-months and 1-year follow up**

Characteristics	Balloon angioplasty group (n=249)			Aggressive medical management group (n=252)		
	Baseline	3 months	1 year	Baseline	3 months	1 year
Receiving aspirin, n (%)	249(100)	234(93.9)	197(79.1)	252(100)	236(93.7)	185(73.4)
Receiving ticagrelor or clopidogrel, n (%)	249(100)	238(95.6)	NA	252(100)	238(94.4)	NA
Receiving statin, n (%)	249(100)	237(95.2)	199(79.9)	252(100)	237(94.0)	191(75.8)

Data are n (%). BA denotes balloon angioplasty. AMM denotes aggressive medical management.

➤ **Table S4. Details of balloon angioplasty**

Characteristics	Balloon angioplasty group ^a (n=249)
Balloon angioplasty duration, s	30.0 (20.0,60.0)
Balloon angioplasty time, no. (%)	
1	182 (75.5)
2	48 (19.9)
3	11 (4.6)
Balloon pressure, atm, no. (%)	6.0 (4.5,7.0)
Residual stenosis degree after the procedure, no. (%)	
0-50%	214 (88.8)
50%-70%	19 (7.9)
70%-100%	8 (3.3)
eTICI after the procedure, no. (%)	
0	0 (0.0)
1	2 (0.8)
2a	1 (0.4)
2b50	9 (3.7)
2b67	20 (8.3)
2c	93 (38.6)
3	116 (48.1)
Rescue stenting, no. (%)	39 (16.2)
Intra-procedural complications, no. (%)	42 (17.4)
Vasospasm	3 (1.2)
Arterial dissection	35 (14.5)
Pseudoaneurysm	0 (0.0)
Arterial occlusion	1 (0.4)
Arterial perforation	0 (0.0)
Arterial rupture	0 (0.0)
hemorrhage	1 (0.4)
Thrombosis	4 (1.7)
Restenosis of the qualifying artery within 1 y ^{b,c} , no. (%)	24 (15.7)
TIA or stroke clearly related to restenosis ^c	3/153 (2.0)

Data are n (%), or median (IQR). IQR denotes interquartile range. eTICI denotes expanded Thrombolysis in the Cerebral Infarction. BA denotes balloon angioplasty.

^a 241 of 249 patients underwent BA. 8 patients didn't undergo BA due to crossover to the AMM group (7 patients) and direct balloon-expanding stenting (1 patient).

^bRestenosis of the qualifying artery within 1 year was based on subsequent neurovascular imaging, stenosis degree >70% or increased by 30%.

^c153 patients had intact follow-up cerebrovascular imaging within 1 year to assess the restenosis of the qualifying artery.

➤ **Table S5 The number of patients and reason for revascularization beyond 30 days through 1 year in the primary population of the BASIS trial**

Reasons for revascularization	Balloon angioplasty group	Aggressive medical management group	Number
Acute revascularization (acute qualifying artery occlusion accompanied by neurological deficit)	1	1	2
Elective revascularization (Hard TIA: culprit artery stenosis that causes recurrent transient ischemic attack lasting longer than 10 minutes or new disabling neurological symptoms)	2	10	12
Elective revascularization (ischemic stroke caused by the culprit artery stenosis)	0	10	10
Total	3	21	24

TIA denotes transient ischemic attack.

➤ **Table S6 Post hoc analysis of center-effect adjustment**

Outcome	Balloon angioplasty group (n=249)	Aggressive medical management group (n=252)	Measure of Effect	Treatment Effect (95% CI) *	P value
Primary outcome, no. (%)					
Any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion, or any ischemic stroke or revascularization of the qualifying artery beyond 30 days through 1 year after enrollment	9 (3.9)	34 (13.5)	Hazard ratio	0.32 (0.16 to 0.62)	.001

Data are n (%). CI denotes confidence interval. IQR denotes interquartile range.

* Treatment effects are reported as hazard ratio with the 95% CI by adjusting for groups and centers.

We divided the participating centers into Beijing Tiantan Hospital and other centers when adjusting the center effect.

➤ **Table S7 Post-hoc analysis of treatment interaction on the primary outcome regarding different sites**

Site	No. of patients	Balloon angioplasty group (N=233)	Aggressive medical management group (N =238)	Measure of Effect	Treatment Effect (95% CI)	P for interaction
Beijing Tiantan Hospital	256	4(2.9)	19(16.1)	Hazard ratio	0.17(0.06 to 0.51)	.10
Other centers	245	7(6.3)	15(11.2)	Hazard ratio	0.56(0.23 to 1.36)	

Data are n (%). CI denotes confidence interval. IQR denotes interquartile range.

➤ **Table S8. Study outcomes in the per-protocol population (PPS)**

Outcome	Balloon angioplasty group (n=249)	Aggressive medical management group (n=252)	Measure of Effect	Treatment Effect (95% CI)	p value
Primary outcome, no. (%)					
Any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion, or any ischemic stroke or revascularization of the qualifying artery beyond 30 days through 1 year after enrollment ^a	9 (3.9)	33 (13.9)	Hazard ratio	0.27 (0.13 to 0.56)	<.001
Any stroke or all-cause death within 30 days after enrollment	6 (2.6)	4 (1.7)	Hazard ratio	1.55 (0.44 to 5.49)	
Any ischemic stroke from the qualifying artery beyond 30 days through 1 year after enrollment	1 (0.4)	18 (7.6)	Hazard ratio	0.06 (0.01 to 0.41)	
Revascularization of the qualifying artery beyond 30 days through 1 year after enrollment ^a	3 (1.3)	20 (8.4)	Hazard ratio	0.15 (0.04 to 0.50)	

Data are n (%). CI denotes confidence interval.

^a Fulfills one of the following requirements to be considered for revascularization of the qualifying artery: (1) Acute revascularization: acute qualifying artery occlusion accompanied by neurological deficit, requiring intravenous thrombolysis, intraarterial thrombolysis, mechanical thrombectomy, balloon/stent angioplasty, or intracranial-extracranial bypass; (2) Elective revascularization: neurologic symptom-driven revascularization, including BA, stent implantation, or intracranial-extracranial bypass.

➤ **Table S9. Study outcomes in the as-treated population (ATS)**

Outcome	Balloon angioplasty group (n=249)	Aggressive medical management group (n=252)	Measure of Effect	Treatment Effect (95% CI)	p value
Primary outcome, no. (%)					
Any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion, or any ischemic stroke or revascularization of the qualifying artery beyond 30 days through 1 year after enrollment ^a	11 (4.5)	34 (13.4)	Hazard ratio	0.32 (0.16 to 0.64)	.001
Any stroke or all-cause death within 30 days after enrollment	8 (3.3)	4 (1.6)	Hazard ratio	2.09 (0.63 to 6.92)	
Any ischemic stroke from the qualifying artery beyond 30 days through 1 year after enrollment	1 (0.4)	19 (7.5)	Hazard ratio	0.05 (0.01 to 0.39)	
Revascularization from the qualifying artery beyond 30 days through 1 year after enrollment ^a	3 (1.2)	21 (8.3)	Hazard ratio	0.14 (0.04 to 0.47)	

Data are n (%). CI denotes confidence interval.

^a Fulfills one of the following requirements to be considered for revascularization of the qualifying artery: (1) Acute revascularization: acute qualifying artery occlusion accompanied by neurological deficit, requiring intravenous thrombolysis, intraarterial thrombolysis, mechanical thrombectomy, balloon/stent angioplasty, or intracranial-extracranial bypass; (2) Elective revascularization: neurologic symptom-driven revascularization, including BA, stent implantation, or intracranial-extracranial bypass.

➤ **Table S10. Post-hoc analysis of the comparison of any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion or any ischemic stroke from the qualifying artery beyond 30 days through 1 year after enrollment between BA and AMM groups**

Outcome	Balloon angioplasty group	Aggressive medical management group	Measure of Effect	Treatment Effect (95% CI)	p value
Intention-to-treat population, no. (%)	N=249	N=252			
Any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion, or any ischemic stroke from the qualifying artery beyond 30 days through 1 year after enrollment	9 (3.6)	23 (9.1)	Hazard ratio	0.39 (0.18-0.85)	.01
Per-protocol population, no. (%)	N=233	N=238			
Any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion, or any ischemic stroke from the qualifying artery beyond 30 days through 1 year after enrollment	7 (3.0)	22 (9.2)	Hazard ratio	0.32 (0.14-0.75)	.01
As-treated population, no. (%)	N=247	N=254			
Any stroke or all-cause death within 30 days after enrollment or after BA of the qualifying lesion, or any ischemic stroke from the qualifying artery beyond 30 days through 1 year after enrollment	9 (3.6)	23 (9.1)	Hazard ratio	0.40 (0.19-0.86)	.02

Data are n (%). CI denotes confidence interval.

➤ **Table S11. Number of patients with other serious adverse events (by system organ class) up to 1-year visit**

System organ class, n (%)	Balloon angioplasty group (n=249)	Aggressive medical management group (n=252)	P value
Overall	12 (4.8)	14 (5.6)	0.84 ^c
Nervous system disorders	7 (2.8)	12 (4.8)	0.35 ^c
Symptomatic intracranial hemorrhage ^a	3 (1.2)	1 (0.4)	0.37 ^d
Asymptomatic intracranial hemorrhage	3 (1.2)	0 (0.0)	0.12 ^d
Any intracranial hemorrhage	6 (2.4)	1 (0.4)	0.07 ^d
Disabling stroke ^b	6 (2.4)	18 (7.1)	0.02 ^c
Vascular and lymphatic system disorder	2 (0.8)	1 (0.4)	0.62 ^d
Metabolic and nutritional disease	1 (0.4)	1 (0.4)	1.00 ^d
Infection	0 (0.0)	2 (0.8)	0.50 ^d
Various surgical and medical operations	2 (0.8)	0 (0.0)	0.25 ^d
Respiratory, thoracic and mediastinal disorder	1 (0.4)	1 (0.4)	1.00 ^d
Gastrointestinal disorder	1 (0.4)	1 (0.4)	1.00 ^d
Injury and poisoning	0 (0.0)	1 (0.4)	1.00 ^d
Benign, malignant, and unexplained tumor (including cystic and polypoid)	1 (0.4)	0 (0.0)	0.50 ^d
Reproductive system and breast disease	1 (0.4)	0 (0.0)	0.50 ^d

^a Symptomatic intracranial hemorrhage was defined as subarachnoid, parenchymal, or intraventricular hemorrhage identified on brain MRI or CT, which led to new neurologic symptoms (consciousness level change, headache, or focal neurologic symptoms), lasting over 24 hours or a seizure.

^b Disabling stroke was defined as modified Rankin Scale ≥ 2 at 1 year.

^c Tested by Chi-square test.

^d Tested by Fisher's exact test.

➤ **Table S12. The proportion of patients meeting the proposed target of vascular risk factor management at 1-year follow-up**

Risk Factors	Balloon angioplasty group (n=249)	Aggressive medical management group (n=252)
No. Patients evaluated	139	120
LDL-c level is < 1.8mmol/L, no. (%)	66(47.5)	61(50.8)
No. Patients evaluated	141	122
Triglycerides < 200mg/dL, no. (%)	127(90.1)	104(85.3)
No. Patients evaluated	65	66
HbA1c < 7.0%, no. (%)	46(70.8)	48(72.7)
No. Patients evaluated	166	160
≤140/90 mmHg (without DM), no. (%)	119(71.7)	119(74.4)
No. Patients evaluated	81	83
≤130/80 mmHg (DM), no. (%)	17(21.0)	24(28.9)