

Supplementary Online Content

Flint AJ, Meyers BS, Rothschild AJ, et al. Effect of continuing olanzapine vs placebo on relapse among patients with psychotic depression in remission: the STOP-PD II randomized clinical trial. *JAMA*. doi:10.1001/jama.2019.10517

eFigure 1. Mixed Model Estimated Least Square Means of Weight in Randomized Groups in the STOP-PD II Clinical Trial

eFigure 2. Mixed Model Estimated Least Square Means of Waist Circumference in Randomized Groups in the STOP-PD II Clinical Trial

eFigure 3. Mixed Model Estimated Least Square Means of Total Cholesterol in Randomized Groups in the STOP-PD II Clinical Trial

eFigure 4. Mixed Model Estimated Least Square Means of LDL Cholesterol in Randomized Groups in the STOP-PD II Clinical Trial

eFigure 5. Mixed Model Estimated Least Square Means of HDL Cholesterol in Randomized Groups in the STOP-PD II Clinical Trial

eFigure 6. Mixed Model Estimated Least Square Means of Triglycerides (for Pattern Mixture Model) in Randomized Groups in the STOP-PD II Clinical Trial


eFigure 7. Mixed Model Estimated Least Square Means of Glucose in Randomized Groups in the STOP-PD II Clinical Trial

eFigure 8. Mixed Model Estimated Least Square Means of HbA1c in Randomized Groups in the STOP-PD II Clinical Trial


eFigure 9. Mixed Model Estimated Least Square Means of Simpson Angus Scale Total Score in Randomized Groups in the STOP-PD II Clinical Trial

This supplementary material has been provided by the authors to give readers additional information about their work.


eFigure 1. Mixed Model Estimated Least Square Means of Weight in Randomized Groups in the STOP-PD II Clinical Trial




eFigure 2. Mixed Model Estimated Least Square Means of Waist Circumference in Randomized Groups in the STOP-PD II Clinical Trial




eFigure 3. Mixed Model Estimated Least Square Means of Total Cholesterol in Randomized Groups in the STOP-PD II Clinical Trial




eFigure 4. Mixed Model Estimated Least Square Means of LDL Cholesterol in Randomized Groups in the STOP-PD II Clinical Trial




eFigure 5. Mixed Model Estimated Least Square Means of HDL Cholesterol in Randomized Groups in the STOP-PD II Clinical Trial




eFigure 6. Mixed Model Estimated Least Square Means of Triglycerides (for Pattern Mixture Model) in Randomized Groups in the STOP-PD II Clinical Trial



eFigure 7. Mixed Model Estimated Least Square Means of Glucose in Randomized Groups in the STOP-PD II Clinical Trial



eFigure8. Mixed Model Estimated Least Square Means of HbA1c in Randomized Groups in the STOP-PD II Clinical Trial



eFigure9. Mixed Model Estimated Least Square Means of Simpson Angus Scale Total Score in Randomized Groups in the STOP-PD II Clinical Trial

