

Supplemental eTable 2 (cont). Baseline dietary variables and their changes by randomized treatment group. Foods and food groups.

	MULTIPLE IMPUTATION: ALL RANDOMIZED PARTICIPANTS			
Food and food groups	Intervention group	Control group	Between-group difference	
	N=3,272	N=3,311		p value
Low-fat dairy (g/week)				
Baseline, median (IQR)	658 (0, 1,775)	687 (0, 1,775)		
6-month change	326 (279 to 373)	68 (23 to 113)	258 (193 to 323)	<0.001
12-month change	267 (217 to 318)	-4 (-50 to 42)	271 (206 to 336)	<0.001
Whole-fat dairy (g/week)				
Baseline, median (IQR)	0 (0, 375)	0 (0, 375)		
6-month change	-174 (-196 to -151)	-110 (-132 to -88)	-63 (-95 to -32)	<0.001
12-month change	-160 (-184 to -135)	-101 (-123 to -78)	-59 (-91 to -27)	<0.001
Total alcohol intake (g/week)				
Baseline, median (IQR)	32 (5, 98)	36 (5, 103)		
6-month change	-13 (-16 to -10)	-6 (-9 to -3)	-7 (-11 to -3)	0.001
12-month change	-12 (-15 to -9)	-4 (-7 to -1)	-8 (-12 to -3)	<0.001
Red wine (g/week)				
Baseline, median (IQR)	33 (0, 29)	4 (0, 29)		
6-month change	1 (-1 to 3)	2 (0 to 4)	-1 (-4 to 1)	0.36
12-month change	2 (0 to 4)	3 (1 to 5)	-1 (-4 to 1)	0.37

IQR, Interquartile range

Baseline data are median (IQR)

6-month and 12-month change are means (95% CI) calculated using mixed effect models taking into account site and intra-cluster correlations (couples) as random factors.