

Supplemental Online Content

Laslett LL, Scheepers LEJM, Antony B, et al. Krill oil for knee osteoarthritis: a randomized clinical trial. *JAMA*. Published online May 22, 2024.
doi:10.1001/jama.2024.6063

eTable 1. Analysis of primary outcome (knee pain (VAS) at 24 weeks), adjustment for age to address missingness

eTable 2. Participant adherence over the study period

eTable 3. Descriptive statistics for Omega-3 Index (%)

eTable 4. Change in secondary endpoints between krill oil and placebo groups, mean (95% CI) at the intermediate time points

eTable 5. Change in Whole-Organ Magnetic Resonance Imaging Score (WORMS) effusion score, 24 weeks

eTable 6. Descriptive statistics for analgesic use at all time points

eTable 7. Summary of adverse events during the study per treatment group (n)

eTable 8. Summary of serious adverse events during the study per treatment group (relationship with treatment)

eFigure. Three-dimensional rendering of effusion-synovitis of a participant at (a) baseline and (b) 6 months

This supplemental material has been provided by the authors to give readers additional information about their work.

eTable 1. Analysis of primary outcome (knee pain (VAS) at 24 weeks), adjustment for age to address missingness

	Change at 6 months mean (95% CI) KRILL OIL	Change at 6 months mean (95% CI) PLACEBO	Absolute between group difference	<i>P</i>
Knee VAS (original data)	-19.93 (-24.67 to -15.20)	-20.21 (-24.87 to -15.54)	-0.27 (-6.92 to 6.38)	0.94
Knee VAS (+ age)	-19.95 (-24.68 to -15.21)	-20.21 (-24.88 to -15.55)	-0.26 (-6.91 to 6.38)	0.94
Knee VAS (Multiple imputation)	-19.94 (-24.56 to -15.32)	-20.29 (-24.9 to -15.68)	-0.35 (-6.79 to 6.09)	0.92

eTable 2. Participant adherence over the study period

	Krill oil (n = 130)	Placebo (n = 132)
Total days of follow-up, mean (SD)	190 (44.5)	184 (34.5)
Follow-up time exceeds 24 weeks, n (%)	55 (42.3)	46 (34.8)
Adherence, consumption of at least 80% of the study treatment*		
At 0 – 12 weeks follow-up, n (%)	91 (100)	89 (100)
At 12 – 24 weeks follow-up, n (%)	82 (96.5)	82 (90.1)
Overall, 0 – 24 weeks follow-up (n = 167), n (%)	82 (98.8)	81 (96.4)
* Adherence only calculated for those with available pill count data and who completed the trial (n = 165 [75%]).		

eTable 3. Descriptive statistics for Omega-3 Index (%)

	Krill oil (n = 130)			Placebo (n = 132)		
	Baseline (n = 116)	12 weeks (n = 100)	24 weeks (n = 96)	Baseline (n = 111)	12 weeks (n = 93)	24 weeks (n = 91)
Mean (SD)	6.5 (1.61)	7.8 (1.51)	8.0 (1.65)	6.5 (1.53)	6.5 (1.28)	6.6 (1.30)
Mean (SD) Δ	-	1.25 (1.42) (n = 98)	1.47 (1.50) (n = 94)	-	0.10 (1.17) (n = 92)	0.14 (1.31) (n = 90)

eTable 4. Change in secondary endpoints between krill oil and placebo groups, mean (95% CI) at all time points (4, 8, 12, 20, 24 weeks)

Element		Krill oil (n = 130)					Placebo (n = 132)				
	Time	n	Final, mean (95% CI) ^d	Baseline	Change mean (95% CI)	n	Final, mean (95% CI) ^d	Baseline	Change mean (95% CI)	Between-group difference in change, mean (95% CI)	P value
Knee pain, VAS ^a	4	117	39.7 (35.3 to 44.1)	47.8 (44.3 to 51.3)	-8.4 (-11.7 to -5.2)	123	40.6 (36.8 to 44.5)	50.2 (46.7 to 53.7)	-8.9 (-12.0 to -5.7)	-0.4 (-4.9 to 4.1)	0.85
	8	106	33.6 (29.3 to 37.9)	47.8 (44.3 to 51.3)	-13.4 (-16.8 to -9.9)	116	35.5 (31.2 to 39.9)	50.2 (46.7 to 53.7)	-14.2 (-17.5 to -10.8)	-0.8 (-5.6 to 4.0)	0.75
	12	109	31.5 (27.0 to 36.0)	47.8 (44.3 to 51.3)	-16.0 (-19.6 to -12.3)	111	33.1 (28.4 to 37.9)	50.2 (46.7 to 53.7)	-16.0 (-19.6 to -12.4)	-0.1 (-5.2 to 5.1)	0.98
	16	103	29.3 (24.7 to 33.9)	47.8 (44.3 to 51.3)	-18.4 (-22.4 to -14.4)	107	31.5 (26.9 to 36.1)	50.2 (46.7 to 53.7)	-18.1 (-22.0 to -14.2)	0.3 (-5.3 to 5.9)	0.91
	20	106	26.5 (21.8 to 31.1)	47.8 (44.3 to 51.3)	-20.1 (-24.5 to -15.7)	109	29.3 (24.7 to 34.0)	50.2 (46.7 to 53.7)	-20.6 (-25.0 to -16.3)	-0.5 (-6.7 to 5.6)	0.87
	24	106	26.5 (22.0 to 31.1)	47.8 (44.3 to 51.3)	-19.9 (-24.7 to -15.2)	109	29.3 (24.4 to 34.2)	50.2 (46.7 to 53.7)	-20.2 (-24.9 to -15.5)	-0.3 (-6.9 to 6.4)	0.94
WOMAC ^b											
Total pain score	4	120	155 (136 to 174)	186 (170 to 203)	-33 (-45 to -20)	124	163 (145 to 180)	210 (192 to 227)	-38 (-50 to -25)	-5 (-23 to 13)	0.59
	8	107	134 (117 to 152)	186 (170 to 203)	-50 (-64 to -37)	117	149 (129 to 170)	210 (192 to 227)	-53 (-67 to -40)	-3 (-22 to 16)	0.77
	12	110	133 (113 to 152)	186 (170 to 203)	-54 (-69 to -40)	113	145 (124 to 166)	210 (192 to 227)	-57 (-72 to -43)	-3 (-24 to 18)	0.76
	16	104	131 (109 to 153)	186 (170 to 203)	-67 (-83 to -51)	109	131 (109 to 153)	210 (192 to 227)	-68 (-84 to -52)	-1 (-24 to 22)	0.92
	20	106	110 (90 to 129)	186 (170 to 203)	-72 (-90 to -54)	111	133 (110 to 157)	210 (192 to 227)	-67 (-85 to -49)	5 (-21 to 30)	0.72
	24	106	110 (89 to 130)	186 (170 to 203)	-87 (-110 to -63)	111	130 (107 to 152)	210 (192 to 227)	-83 (-106 to -59)	3 (-24 to 31)	0.81
Weight bearing pain	4	120	100 (88 to 112)	127 (116 to 138)	-84 (-122 to -46)	124	108 (97 to 119)	141 (130 to 151)	-103 (-141 to -65)	3 (-10 to 16)	0.66
	8	107	88 (77 to 99)	127 (116 to 138)	-37 (-46 to -27)	117	101 (88 to 114)	141 (130 to 151)	-36 (-45 to -26)	1 (-12 to 14)	0.88
	12	110	89 (76 to 101)	127 (116 to 138)	-38 (-47 to -29)	113	98 (84 to 111)	141 (130 to 151)	-38 (-47 to -29)	0.0 (-13 to 13)	0.99
	16	104	81 (69 to 93)	127 (116 to 138)	-46 (-55 to -37)	109	89 (75 to 103)	141 (130 to 151)	-46 (-56 to -37)	0.0 (-14 to 13)	0.97
	20	106	73 (60 to 86)	127 (116 to 138)	-51 (-60 to -41)	111	89 (75 to 104)	141 (130 to 151)	-47 (-57 to -38)	4 (-10 to 17)	0.60
	24	106	74 (61 to 87)	127 (116 to 138)	-51 (-61 to -42)	111	86 (72 to 101)	141 (130 to 151)	-48 (-58 to -39)	3 (-10 to 16)	0.70
Non weight bearing pain	4	120	55 (45 to 64)	60 (52 to 68)	-6 (-11 to 0)	124	54 (47 to 62)	69 (61 to 77)	-10 (-16 to -5)	-5 (-13 to 3)	0.25
	8	107	46 (38 to 54)	60 (52 to 68)	-14 (-20 to -8)	117	49 (40 to 57)	69 (61 to 77)	-17 (-23 to -11)	-3 (-12 to 6)	0.52
	12	110	44 (35 to 53)	60 (52 to 68)	-17 (-23 to -10)	113	47 (38 to 56)	69 (61 to 77)	-19 (-25 to -12)	-2 (-11 to 7)	0.64
	16	104	38 (31 to 46)	60 (52 to 68)	-22 (-29 to -14)	109	42 (34 to 51)	69 (61 to 77)	-22 (-29 to -15)	0 (-10 to 10)	0.94
	20	106	37 (29 to 45)	60 (52 to 68)	-22 (-30 to -14)	111	44 (35 to 53)	69 (61 to 77)	-21 (-28 to -13)	1 (-10 to 12)	0.82
	24	106	36 (27 to 44)	60 (52 to 68)	-23 (-32 to -15)	111	43 (35 to 52)	69 (61 to 77)	-21 (-29 to -13)	2 (-10 to 14)	0.70
Function	4	120	493 (429 to 557)	578 (524 to 633)	-84 (-122 to -46)	121	503 (444 to 561)	618 (560 to 676)	-103 (-141 to -65)	-19 (-73 to 35)	0.48
	8	105	416 (357 to 475)	578 (524 to 633)	-145 (-186 to -103)	115	473 (406 to 540)	618 (560 to 676)	-149 (-189 to -109)	-4 (-62 to 53)	0.88
	12	107	411 (350 to 473)	578 (524 to 633)	-169 (-214 to -125)	112	457 (387 to 526)	618 (560 to 676)	-159 (-203 to -116)	10 (-52 to 72)	0.75
	16	103	371 (310 to 431)	578 (524 to 633)	-201 (-250 to -152)	108	436 (363 to 509)	618 (560 to 676)	-179 (-227 to -132)	22 (-47 to 90)	0.53
	20	105	342 (279 to 405)	578 (524 to 633)	-219 (-273 to -166)	110	437 (361 to 513)	618 (560 to 676)	-186 (-240 to -133)	33 (-43 to 109)	0.39
	24	105	319 (255 to 383)	578 (524 to 633)	-240 (-299 to -182)	110	419 (345 to 494)	618 (560 to 676)	-189 (-247 to -132)	51 (-31 to 133)	0.22
Hand pain, VAS ^c	4	107	21.1 (16.4 to 25.9)	21.7 (16.9 to 26.4)	-2.0 (-5.2 to 1.2)	115	20.5 (16.1 to 25)	24.4 (19.3 to 29.6)	-3.6 (-6.7 to -0.4)	-1.6 (-6.1 to 2.9)	0.49
	8	97	15.2 (11.0 to 19.4)	21.7 (16.9 to 26.4)	-7.1 (-10.5 to -3.7)	103	22.8 (17.5 to 28.1)	24.4 (19.3 to 29.6)	-3.3 (-6.6 to 0.1)	3.8 (-0.9 to 8.6)	0.12
	12	100	17.7 (13.4 to 22.1)	21.7 (16.9 to 26.4)	-4.6 (-8.2 to -1.1)	106	19.2 (14.4 to 24.0)	24.4 (19.3 to 29.6)	-4.7 (-8.2 to -1.2)	-0.1 (-5.0 to 4.9)	0.98
	16	95	16.5 (12.4 to 20.5)	21.7 (16.9 to 26.4)	-5.6 (-9.4 to -1.8)	98	19.2 (14.1 to 24.3)	24.4 (19.3 to 29.6)	-5.2 (-8.9 to -1.4)	0.4 (-4.9 to 5.7)	0.89

Back pain, VAS ^d	20	99	14.6 (11.1 to 18.1)	21.7 (16.9 to 26.4)	-6.9 (-11 to -2.8)	100	19.6 (14.6 to 24.6)	24.4 (19.3 to 29.6)	-5.2 (-9.3 to -1.2)	1.7 (-4.1 to 7.5)	0.57
	24	99	14.4 (10.6 to 18.1)	21.7 (16.9 to 26.4)	-7.6 (-11.9 to -3.3)	100	17.6 (12.8 to 22.3)	24.4 (19.3 to 29.6)	-6.3 (-10.6 to -2.1)	1.3 (-4.8 to 7.3)	0.69
	4	108	24.8 (20.0 to 29.7)	26.7 (22.0 to 31.4)	-2.8 (-6.0 to 0.4)	115	25.8 (21 to 30.6)	33.0 (27.5 to 38.5)	-4.2 (-7.4 to -1.1)	-1.4 (-5.9 to 3.0)	0.53
	8	98	20.0 (15.4 to 24.6)	26.7 (22.0 to 31.4)	-6.2 (-9.5 to -2.8)	104	28.2 (22.9 to 33.5)	33.0 (27.5 to 38.5)	-2.3 (-5.6 to 1.0)	3.9 (-0.9 to 8.6)	0.11
	12	101	19.9 (15.6 to 24.1)	26.7 (22.0 to 31.4)	-7.4 (-10.9 to -3.9)	107	25.6 (20.4 to 30.7)	33.0 (27.5 to 38.5)	-3.3 (-6.7 to 0.2)	4.1 (-0.8 to 9.1)	0.10
Lower leg strength (N) ^d	16	95	18.3 (14.3 to 22.2)	26.7 (22.0 to 31.4)	-9.4 (-13.2 to -5.7)	99	25.1 (20.1 to 30.2)	33.0 (27.5 to 38.5)	-5.1 (-8.8 to -1.4)	4.3 (-0.9 to 9.6)	0.11
	20	99	20.0 (15.5 to 24.6)	26.7 (22.0 to 31.4)	-6.5 (-10.5 to -2.6)	101	23.3 (18.2 to 28.4)	33.0 (27.5 to 38.5)	-6.9 (-10.8 to -2.9)	-0.3 (-6.0 to 5.3)	0.91
	24	99	18.4 (14.2 to 22.7)	26.7 (22.0 to 31.4)	-7.6 (-12.2 to -3.0)	101	22.8 (18 to 27.5)	33.0 (27.5 to 38.5)	-8.9 (-13.5 to -4.4)	-1.3 (-7.9 to 5.2)	0.69
	12	101	72.6 (65.1 to 80.1)	66.5 (59.8 to 73.2)	-2.8 (-6.0 to 0.4)	107	70.2 (62.3 to 78.1)	65.9 (59.4 to 72.3)	-4.2 (-7.4 to -1.1)	-1.4 (-5.9 to 3.0)	0.53
	24	100	72.8 (65.7 to 79.8)	66.5 (59.8 to 73.2)	6.7 (2.6 to 10.7)	107	71.7 (64.3 to 79.1)	65.9 (59.4 to 72.3)	4.4 (0.5 to 8.4)	-2.2 (-7.9 to 3.4)	0.44
Quality of life (AQoL-6D) ^f	12	108	0.8 (0.8 to 0.8)	0.8 (0.7 to 0.8)	0.03 (0.01 to 0.04)	111	0.8 (0.8 to 0.8)	0.7 (0.7 to 0.8)	0.03 (0.01 to 0.04)	-0.0003 (-0.02 to 0.02)	0.98
	24	104	0.8 (0.8 to 0.9)	0.8 (0.7 to 0.8)	0.04 (0.02 to 0.06)	108	0.8 (0.8 to 0.8)	0.7 (0.7 to 0.8)	0.03 (0.01 to 0.05)	-0.01 (-0.04 to 0.01)	0.38
	4	120			17%	123			20%	0.85 (0.50 to 1.46)	0.57
	8	108			26%	117			30%	0.87 (0.57 to 1.32)	0.51
	12	110			35%	113			36%	0.98 (0.69 to 1.39)	0.90
	16	104			37%	108			39%	0.94 (0.66 to 1.33)	0.72
	20	101			43%	100			40%	1.06 (0.77 to 1.48)	0.71
OMERACT-OARSI responder criteria ^g	24	107			47%	110			41%	1.14 (0.84 to 1.55)	0.39
Metabolic factors											
High sensitivity C-reactive protein	12	91	1.2 (0.7 to 2.4)	1.4 (0.7 to 2.9)	-0.53 (-1.32 to 0.26)	92	1.6 (0.9 to 3.1)	1.9 (1.0 to 3.8)	0.24 (-0.55 to 1.03)	0.07 (-1.19 to 1.33)	0.92
	24	82	1.3 (0.8 to 2.3)	1.4 (0.7 to 2.9)	-0.76 (-1.6 to 0.08)	84	1.9 (1.1 to 3.3)	1.9 (1.0 to 3.8)	-0.12 (-0.97 to 0.73)	0.64 (-0.56 to 1.84)	0.30
Triglycerides	12	111	1.3 (1.2 to 1.5)	1.4 (1.2 to 1.6)	-0.11 (-0.23 to 0.02)	114	1.5 (1.3 to 1.6)	1.3 (1.2 to 1.4)	0.14 (0.01 to 0.26)	0.24 (0.07 to 0.42)	0.01
	24	98	1.3 (1.2 to 1.4)	1.4 (1.2 to 1.6)	-0.08 (-0.21 to 0.05)	102	1.4 (1.3 to 1.6)	1.3 (1.2 to 1.4)	0.07 (-0.06 to 0.20)	0.15 (-0.04 to 0.33)	0.11
HDL	12	111	1.6 (1.5 to 1.7)	1.6 (1.5 to 1.6)	0.04 (0.0 to 0.08)	113	1.6 (1.5 to 1.6)	1.5 (1.5 to 1.6)	0.03 (-0.01 to 0.07)	-0.01 (-0.07 to 0.05)	0.76
Cholesterol	24	97	1.6 (1.5 to 1.6)	1.6 (1.5 to 1.6)	0.02 (-0.03 to 0.06)	101	1.5 (1.4 to 1.6)	1.5 (1.5 to 1.6)	-0.01 (-0.06 to 0.03)	-0.03 (-0.09 to 0.03)	0.32
LDL Cholesterol	12	110	3.1 (2.9 to 3.3)	3.0 (2.9 to 3.2)	0.1 (-0.01 to 0.2)	112	3.1 (2.9 to 3.3)	3.2 (3.0 to 3.3)	-0.05 (-0.15 to 0.06)	-0.14 (-0.29 to 0.005)	0.06
	24	96	3.0 (2.8 to 3.2)	3.0 (2.9 to 3.2)	-0.01 (-0.12 to 0.1)	101	3.2 (3.0 to 3.4)	3.2 (3.0 to 3.3)	0.0005 (-0.11 to 0.11)	0.01 (-0.14 to 0.17)	0.90
Fasting glucose	12	112	5.4 (5.3 to 5.6)	5.3 (5.2 to 5.5)	0.09 (-0.07 to 0.24)	112	5.5 (5.2 to 5.7)	5.3 (5.1 to 5.5)	0.15 (-0.01 to 0.31)	0.07 (-1.19 to 1.33)	0.92
	24	100	5.4 (5.2 to 5.6)	5.3 (5.2 to 5.5)	0.07 (-0.12 to 0.26)	100	5.4 (5.2 to 5.6)	5.3 (5.1 to 5.5)	0.11 (-0.08 to 0.29)	0.04 (-0.23 to 0.30)	0.79

Abbreviations: AQoL-6D: Assessment of Quality of Life – 6D; hs-CRP: high-sensitivity C-reactive protein; HDL: high-density lipoprotein; ICOAP: LDL: low-density lipoprotein; OMERACT-OARSI: Outcome Measures in Rheumatology – Osteoarthritis Research Society International; WOMAC: Western Ontario and McMaster University Index.

No values were imputed for missing values. To address the impact of missing data, we assessed variables related to missingness. In the primary outcome analyses, age was the only variable associated with missingness. Therefore, as a sensitivity analysis, we adjusted for the impact of age and the missingness induced by age. No significant difference in the results was observed (Supplement eTable 1).

Analyses are adjusted for baseline if p values for difference in Table 1 are ≤0.1, that is, high sensitivity-CRP, back pain, WOMAC total pain score and weight bearing pain score.

^a Knee pain (VAS), 0 to 100, with higher scores indicating worse pain;

^b Composite score total pain (WOMAC), 0 to 500, with higher scores indicating worse pain, weight bearing pain (0-300), non-weight bearing pain (0-200); composite score physical function (WOMAC), 0 – 1700, with higher scores indicating worse physical function;

^c hand pain (VAS), 0 to 100, with higher scores indicating worse pain;

^d back pain (VAS), 0 to 100, with higher scores indicating worse pain;

^e Lower limb strength (Newtons), 0 to 250, with higher scores indicating greater force or strength of the index lower limb;

^f AQL-6D, -0.04 to 1.0; a utility score that ranges from 1.00 (full health) to 0.00 (death-equivalent health states) to -0.04 (health states worse than death);

^g Response to treatment according to OMERACT-OARSI responder criteria,²⁵ where responder is defined as either (i) $\geq 50\%$ improvement and an absolute change ≥ 20 points in pain (using the VAS) or function (using WOMAC function score); or (ii) improvement of $\geq 20\%$ with an absolute change ≥ 10 mm in at least two of the following three categories, at all time points (4, 8, 12, 16, 20, and 24 weeks; MCID unknown);

^h Values are shown as mean (SD) and corresponding 95% confidence intervals, except for effusion-synovitis volume and high sensitivity C-reactive protein (hs-CRP) which are shown as median (IQR)

^k Results are relative risks (RR) with 95% CI, derived from log binomial regression. Responder index, calculated at the final follow-up visit, is a measure of change from baseline.

eTable 5. Change in Whole-Organ Magnetic Resonance Imaging Score (WORMS) effusion score, 24 weeks

	Krill oil	Placebo
Smaller - 2 units	2 (1.9%)	2 (1.9%)
Smaller - 1 unit	10 (12%)	16 (12%)
No change	80 (72%)	75 (72%)
Larger - 1 unit	12 (12%)	13 (12%)
Larger - 2 units	3 (2.8%)	3 (2.8%)
Total	107 (100%)	109 (100%)

eTable 6. Descriptive statistics for analgesic use at all time points

Weeks	Krill oil		Placebo	
	Increase, n	Decrease, n	Increase, n	Decrease, n
4	--	--	--	--
8	--	--	--	--
12	1	1	3	--
16	--	--	--	--
20		--	--	--
24	--	--	1	--
The change in analgesic use was defined as any modification in the use of nonsteroidal anti-inflammatory drugs (NSAIDs) compared to the previous visit.				

eTable 7. Summary of adverse events during the study per treatment group (n)

Adverse Event¹	Krill oil (n = 130)	Placebo (n = 132)
Musculoskeletal and connective tissue disorders		
Ankle fracture		1
Arthritis		1
Bursitis		1
Extra skeletal ossification		1
Gout flare	1	
Inguinal herniorrhaphy		1
Joint clicking		1
Joint pain		4
Leg cramps	1	
Ligament sprain		2
Meniscus injury	1	
Muscle spasms	1	
Muscle strain		1
Osteoarthritis	1	
Osteodynia		1
Osteoporosis	1	
Pain in back	3	1
Pain in elbow		1
Pain in extremity	1	5
Pain in fingers	1	1
Pain in foot	1	1
Pain in hip	3	2
Pain in jaw	1	
Pain in knee	10	9
Pain in legs	1	
Pain in shoulder	1	
Pain in wrist	1	
Radius fracture		1
Rheumatoid arthritis		1
Sacral pain		1
Spinal osteoarthritis		1
Stress fracture	1	
Synovial rupture	1	1
Tendonitis		2
Tennis elbow	1	
Trochanteric bursitis		1
Respiratory, thoracic and mediastinal		
Asthma		1
Atypical pneumonia		1
Cough		2
Epistaxis		1
Lower respiratory tract infection	1	1
Nasopharyngitis	1	2

Obstruction bronchial	1	
Pharyngitis	1	
Pneumonia		1
Pulmonary mass	1	
Sarcoidosis		1
Shortness of breath		1
Upper respiratory tract infection	4	4
Viral infections NEC		1
Gastrointestinal discomfort and disorders		
Abdominal discomfort	3	
Blood in stool	1	
Constipation	1	
Diarrhoea	1	2
Diverticulitis		1
Gastroenteritis	1	
Gastroesophageal reflux disease	3	1
Hiatus hernia		1
Loose stool	1	
Nausea or vomiting		1
Pruritus	1	
Toothache	1	
Investigations		
Blood parathyroid hormone abnormal		1
Blood test	1	
Bowel screening		1
Colonoscopy		3
Endoscopy	1	
Knee arthroscopy	2	
Liver function test abnormal	1	
MRI of knee	1	
Protein in urine present	1	
Suprapubic pain	1	
Weight increase	2	3
Surgical and medical procedures		
Blepharoplasty		1
Bunion operation	2	
Cataract extraction	1	1
Cholecystectomy (elective)		1
Hip joint injection		1
Inguinal hernia repair		1
Knee arthroplasty	1	
Mammoplasty		1
Polyp removal		1
Prostate biopsy	1	
Spinal fusion		1
Injury, poisoning and procedural complications		
Ankle sprain	2	

Fall injury	2	1
Food poisoning		1
Injury	1	
Knee injury		2
Ligament sprain		1
Scaphoid bone fracture	1	
Sport injury		1
Skin and subcutaneous tissue disorders		
Onycholysis		1
Petechiae	1	
Pruritus	3	1
Rash macular	1	
Wound		1
General system disorders		
Chest pain		1
Fatigue	2	
Flu like symptoms		1
Peripheral swelling	1	
Nervous system disorders		
Carpal tunnel syndrome		1
Headache	1	2
Neuropathic pain	1	
Infections and infestations		
Helicobacter pylori infection	1	
Influenza		1
Shingles	1	
Viral infection		1
Vascular disorders		
Blood pressure high	1	
Haemorrhoids	1	
Hypertension		1
Increased tendency to bruise		1
Cardiac disorders		
Cardiac arrhythmias		1
Heart enlarged		1
Palpitations	1	
Eye disorders		
Corneal ulcer		1
Eye pain	1	
Macular hole		1
Renal and urinary disorders		
Urinary tract infection	1	1
Urine colour abnormal	1	
Neurological disorders		
Hypoesthesia		1
Vertigo		1

Metabolism and nutrition disorders		
Elevated cholesterol	1	1
Blood and lymphatic system disorders		
Transfusion dependent anaemia		1
Reproductive system and breast disorders		
Breast lump	1	
Psychiatric disorders		
Depression	1	
Statistical analysis of differences in risk between groups was not undertaken because of low numbers of adverse events reported		

eTable 8. Summary of serious adverse events during the study per treatment group (relationship with treatment)

Serious Adverse Event	Placebo (n = 132)				Krill oil (n = 130)		
	Unknown*	Mild	Moderate	Severe	Mild	Moderate	Severe
Appendicitis				1 (Unlikely)			
Basal cell carcinoma		1 (Possible)					
Bronchiolitis	1 (NA)						
Cholecystectomy						1 (Possible)	
Colectomy							1 (Possible)
Colonic polyp					1 (Possible)		
Malignant melanoma			1 (Possible)	1 (Possible)	1 (Possible)	1 (Possible)	
Mastoidectomy							1 (Possible)
Open reduction of fracture with internal fixation							1 (Unlikely)
Skin lesion removal	1 (NA)						1 (Possible)
Transfusion dependent anaemia							1 (Probable)
* Severity and relationship with treatment is not recorded. Intensity of the adverse event are classified as: (i) Mild: Symptom(s) barely noticeable to the patient or does not make the patient uncomfortable. The adverse events does not influence performance or functioning. Prescription drugs are not ordinarily needed for relief of symptom(s); (ii) Moderate: Symptom(s) of a sufficient severity to make the patient uncomfortable. Performance of daily activities is influenced. Treatment of symptom(s) may be needed; or (iii) Severe: Symptom(s) of a sufficient severity to cause the patient severe discomfort. Severity may cause cessation of treatment with the drug. Treatment for symptom(s) may be given. Statistical analysis of differences in risk between groups was not undertaken because of low numbers of adverse events reported.							

eFigure. Three-dimensional rendering of effusion-synovitis of a participant at (left) baseline and (right) 24 weeks

