

Assignment 3: Data Exploration

Elsie Liu, Section #3

OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Exploration.

Directions

1. Change “Student Name, Section #” on line 3 (above) with your name and section number.
2. Work through the steps, **creating code and output** that fulfill each instruction.
3. Be sure to **answer the questions** in this assignment document.
4. When you have completed the assignment, **Knit** the text and code into a single PDF file.
5. After Knitting, submit the completed exercise (PDF file) to the dropbox in Sakai. Add your last name into the file name (e.g., “FirstLast_A03_DataExploration.Rmd”) prior to submission.

The completed exercise is due on <>.

Set up your R session

1. load necessary packages (tidyverse), upload: Neonics: ECOTOX neonicotinoid dataset (ECOTOX_Neonicotinoids_Insects_raw.csv) Litter: Niwot Ridge NEON dataset for litter and woody debris: (NEON_NIWO_Litter_massdata_2018-08_raw.csv)

```
library(tidyverse)
library(ggplot2)
Neonics <- read.csv("../Data/Raw/ECOTOX_Neonicotinoids_Insects_raw.csv", stringsAsFactors = TRUE)
Litter <- read.csv("../Data/Raw/NEON_NIWO_Litter_massdata_2018-08_raw.csv", stringsAsFactors = TRUE)
```

Learn about your system

2. The neonicotinoid dataset was collected from the Environmental Protection Agency’s ECOTOX Knowledgebase, a database for ecotoxicology research. Neonicotinoids are a class of insecticides used widely in agriculture. The dataset that has been pulled includes all studies published on insects. Why might we be interested in the ecotoxicology of neonicotinoids on insects? Feel free to do a brief internet search if you feel you need more background information.

Answer: Neonicotinoid is a kind of active ingredient for insecticides and have been applied to control pests in a variety of crops; but at the same time, they may also have effect on non-target organisms such as pollinators (e.g. bees).

3. The Niwot Ridge litter and woody debris dataset was collected from the National Ecological Observatory Network, which collectively includes 81 aquatic and terrestrial sites across 20 ecoclimatic domains. 32 of these sites sample forest litter and woody debris, and we will focus on the Niwot Ridge long-term ecological research (LTER) station in Colorado. Why might we be interested in studying litter and woody debris that falls to the ground in forests? Feel free to do a brief internet search if you feel you need more background information.

Answer: The woody debris and litter could benefit forests' nutrient cycling, form habitat for some reptile species, or even become shelters for fish and amphibians while at the same time they could be an important fuel source for wildfires.

4. How is litter and woody debris sampled as part of the NEON network? Read the NEON_Litterfall_UserGuide.pdf document to learn more. List three pieces of salient information about the sampling methods here:

Answer:

1. sampling in 20 40m x 40m plots (for sites with low-statured vegetation over the tower airsheds, litter sampled in 4 40m x 40m tower plots plus 26 20m x 20m plots;
2. plot edges are separated by a distance 150% of one edge of the plot;
3. temporal design: ground traps are sampled once per year.

Obtain basic summaries of your data (Neonics)

5. What are the dimensions of the dataset?

```
dim(Neonics)
```

```
## [1] 4623 30
```

```
dim(Litter)
```

```
## [1] 188 19
```

6. Using the `summary` function on the "Effect" column, determine the most common effects that are studied. Why might these effects specifically be of interest?

```
summary(Neonics$Effect)
```

```
##      Accumulation      Avoidance      Behavior      Biochemistry
##           12           102           360           11
##      Cell(s)      Development      Enzyme(s) Feeding behavior
##           9           136           62           255
##      Genetics      Growth      Histology      Hormone(s)
##          82           38           5           1
## Immunological      Intoxication      Morphology      Mortality
##          16           12           22           1493
##      Physiology      Population      Reproduction
##           7           1803           197
```

Answer: "Population" is the most common effect in the "Effect" column. Researchers might be interested in how one (or several) effect could influence insects' behavior separately (or together).

7. Using the `summary` function, determine the six most commonly studied species in the dataset (common name). What do these species have in common, and why might they be of interest over other insects? Feel free to do a brief internet search for more information if needed.

```
Neonics.summ <- as.data.frame(summary(Neonics$Species.Common.Name))
colnames(Neonics.summ) <- "count"
Neonics.summ$species <- rownames(Neonics.summ)
head(Neonics.summ[order(Neonics.summ$count, decreasing = T),],n=7)
```

```
##      count      species
## (Other)      670      (Other)
## Honey Bee    667      Honey Bee
## Parasitic Wasp 285      Parasitic Wasp
## Buff Tailed Bumblebee 183 Buff Tailed Bumblebee
## Carniolan Honey Bee 152 Carniolan Honey Bee
## Bumble Bee   140      Bumble Bee
```

```
## Italian Honeybee      113      Italian Honeybee
```

Answer: Honey Bee, Parasitic Wasp, Buff Tailed Bumblebee, Carniolan Honey Bee, Bumble Bee, Italian Honeybee are the most commonly studied species. These insect species are pollinators and play important roles in the ecosystem. Once these species have been influenced by the pesticide, some primary producer that rely on them would lose their way of reproduction.

8. Concentrations are always a numeric value. What is the class of `Conc.1..Author.` in the dataset, and why is it not numeric?

```
class(Neonics$Conc.1..Author.)
```

```
## [1] "factor"
```

```
summary(Neonics$Conc.1..Author.)
```


```
##      0.37/      10/      NR/      NR      1      1023      0.40/      2/
##      208      127      108      94      82      80      69      63
##      10      0.053/      100      50/      0.5/      0.03      0.05/      0.45
##      62      59      56      51      45      44      43      43
##      0.1/      0.45/      1.0/      2.27/      50      0.125      500/      0.5
##      42      40      40      40      36      33      33      32
##      0.048/      0.15/      1/      48      25.0/      12/      0.027      2.4
##      30      30      30      30      28      27      26      26
##      0.2/      0.56/      100/      3      0.01/      1000/      3/      0.336
##      25      24      23      23      22      22      22      21
##      1.5/      0.05      1.5      2.60/      20.0/      6      6.80/      62.5/
##      21      20      20      20      20      20      20      20
##      0.005      0.4/      0.18/      0.3/      1000      40      0.00355/      0.1
##      18      18      17      17      17      17      16      16
##      0.4      150/      300      80/      0.053      0.24      0.28      125/
##      16      16      16      16      15      15      15      15
##      9      0.0001      0.0004/      0.084/      0.15      0.6      12.5/      144.0/
##      15      14      14      14      14      14      14      14
##      350/      40.0/      48/      56      84/      0.17/      125      14
##      14      14      14      14      14      13      13      13
##      16      17      0.047/      0.25/      0.28/      1.28/      1.81/      112
##      13      13      12      12      12      12      12      12
##      150      2.5/      25      60/      75/      0.02/      0.025/      0.29
##      12      12      12      12      12      11      11      11
##      37.5/      4/      5      (Other)
##      11      11      11      1817
```

Answer: According to EPA's ECOTOX Web site, "Conc.1..Author." stands for "Author Reported and Standardized Concentration". The class of this field is factor but not numeric because there are some non-numeric strings like "NR/" or "numbers+"/.

Explore your data graphically (Neonics)


9. Using `geom_freqpoly`, generate a plot of the number of studies conducted by publication year.

```
ggplot(Neonics) +
  geom_freqpoly(aes(x = Publication.Year), bins = 50, color = "navyblue")
```



10. Reproduce the same graph but now add a color aesthetic so that different Test.Location are displayed as different colors.

```
ggplot(Neonics) +  
  geom_freqpoly(aes(x = Publication.Year,color = Test.Location), bins = 50)
```




Interpret this graph. What are the most common test locations, and do they differ over time?

Answer: According to the graph above, Lab/Field natural are most common test locations. “field natural” is more common in 1990s and 2009 and lab is more common after 2000 (several exceptions).

11. Create a bar graph of Endpoint counts. What are the two most common end points, and how are they defined? Consult the ECOTOX_CodeAppendix for more information.

```
ggplot(Neonics, aes(x = Endpoint)) +
  geom_bar() +
  theme(axis.text.x = element_text(size = 5))
```



Answer: The two most common end points are NOEL (No-observable-effect-level) & LOEL (Lowest-observable-effect-level).

Explore your data (Litter)

- Determine the class of collectDate. Is it a date? If not, change to a date and confirm the new class of the variable. Using the `unique` function, determine which dates litter was sampled in August 2018.

```
library(lubridate)
```

```
##
## Attaching package: 'lubridate'
## The following objects are masked from 'package:base':
##
##   date, intersect, setdiff, union
```

```
class(Litter$collectDate)
```

```
## [1] "factor"
```

```
Litter$collectDate <- ymd(Litter$collectDate)
class(Litter$collectDate)
```

```
## [1] "Date"
```

```
unique(Litter$collectDate)
```

```
## [1] "2018-08-02" "2018-08-30"
```

13. Using the `unique` function, determine how many plots were sampled at Niwot Ridge. How is the information obtained from `unique` different from that obtained from `summary`?


```
unique(Litter$plotID)
```

```
## [1] NIWO_061 NIWO_064 NIWO_067 NIWO_040 NIWO_041 NIWO_063 NIWO_047 NIWO_051
## [9] NIWO_058 NIWO_046 NIWO_062 NIWO_057
## 12 Levels: NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 NIWO_057 ... NIWO_067
```

Answer: 12 plots were sampled at Niwot Ridge. “`unique()`” only counts levels while “`summary()`” also counts the frequency of these levels.


14. Create a bar graph of `functionalGroup` counts. This shows you what type of litter is collected at the Niwot Ridge sites. Notice that litter types are fairly equally distributed across the Niwot Ridge sites.

```
ggplot(Litter, aes(x=functionalGroup)) +
  geom_bar()
```



15. Using `geom_boxplot` and `geom_violin`, create a boxplot and a violin plot of `dryMass` by `functionalGroup`.

```
ggplot(data = Litter) +
  geom_boxplot(aes(x= functionalGroup, y = dryMass))
```




```
ggplot(data = Litter) +
  geom_violin(aes(x = functionalGroup, y = dryMass),
    draw_quantiles = c(0.25, 0.5, 0.75))
```

```
## Warning in regularize.values(x, y, ties, missing(ties), na.rm = na.rm):
## collapsing to unique 'x' values
```

```
## Warning in regularize.values(x, y, ties, missing(ties), na.rm = na.rm):
## collapsing to unique 'x' values
```

```
## Warning in regularize.values(x, y, ties, missing(ties), na.rm = na.rm):
## collapsing to unique 'x' values
```

Why is the boxplot a more effective visualization option than the violin plot in this case?

Answer: The boxplot could clearly show the outliers and the area of box itself (representing the distribution of dry mass in each functional group). On the contrary, the violin plot failed to show the distribution of groups like Mixed, Needles and Twigs/branches as these groups have relatively large ranges and shrink to single lines.

What type(s) of litter tend to have the highest biomass at these sites?

Answer: Needles are of the highest biomass.