
UAV Obstacle collision avoidance system

Subsystem integration for safer autonomous flights

By

ÁLVARO MELGOSA PASCUAL



Department of Bioengineering and Aerospace Engineering
UNIVERSIDAD CARLOS III DE MADRID

SEPTEMBER 2016

ABSTRACT

Here goes the abstract

DEDICATION AND ACKNOWLEDGEMENTS

Here goes the dedication.

TABLE OF CONTENTS

	Page
Abstract	ii
Dedication and Acknowledgements	iv
Table of Contents	v
List of Figures	vii
List of Tables	ix
1 Introduction	1
1.1 Section	1
1.1.1 Subsection	1
A Appendix A	3
Bibliography	5

LIST OF FIGURES

FIGURE	Page
1.1 Developmental zones of an Arabidopsis root.	1

LIST OF TABLES

TABLE	Page
-------	------

CHAPTER

1

INTRODUCTION

Begins a chapter.

APPENDIX



APPENDIX A

Begins an appendix

BIBLIOGRAPHY

- [GS02] C. Grierson and J. Schiefelbein.
The Arabidopsis Book.
American Society of Plant Biologist, 2002.
- [JS06] M. Jones and N. Smirnov.
Nuclear dynamics during the simultaneous and sustained tip growth of multiple root hairs arising from a single root epidermal cell.
J. of Exp. Bot., 57(15):4269–4275, 2006.
- [MS94] J. D. Masucci and J. W. Schiefelbein.
The *rh6* mutation of *arabidopsis thaliana* alters root-hair initiation through an auxin- and ethylene-associated process.
Plant. Physiol., 106:1335–1346, 1994.
- [PG09] R.J.H. Payne and C.S. Grierson.
A theoretical model for rop localisation by auxin in *arabidopsis* root hair cells.
PLoS ONE, 4(12):e8337. doi:10.1371/journal.pone.0008337, 2009.
- [RDH⁺01] S. Rigas, G. Debrosses, K. Haralampidis, F. Vicente-Angulo, K. A. Feldman, A. Grabov, L. Dolan, and P. Hatzopoulos.
Trh1 encodes a potassium transporter required for tip growth in *arabidopsis* root hairs.
The Plant Cell, 13:139–151, 2001.

