

Reperfusion therapies and in-hospital outcomes for ST-elevation myocardial infarction in Europe: the ACVC-EAPCI EORP STEMI Registry of the European Society of Cardiology

Uwe Zeymer^{1,*}, Peter Ludman², Nicolas Danchin³, Petr Kala⁴, Cécile Laroche⁵, Masoumeh Sadeghi⁶, Roberto Caporale⁷, Sameh Mohamed Shaheen⁸, Jacek Legutko⁹, Zaza Iakobsishvili¹⁰, Khalid F. Alhabib¹¹, Zuzana Motovska¹², Martin Studencan¹³, Jorge Mimoso¹⁴, David Becker¹⁵, Dimitrios Alexopoulos¹⁶, Zviad Kereseslidze¹⁷, Sinisa Stojkovic¹⁸, Parounak Zelveian¹⁹, Artan Goda²⁰, Erkin Mirrakhimov^{21,22}, Gani Bajraktari²³, Hasan Al-Farhan²⁴, Pranas Šerpytis²⁵, Bent Raungaard²⁶, Toomas Marandi²⁷, Alice May Moore²⁸, Martin Quinn²⁹, Pasi Paavo Karjalainen³⁰, Gabriel Tatu-Chitolu³¹, Chris P. Gale³², Aldo P. Maggioni³³, and Franz Weidinger³⁴; on behalf of the ACVC EAPCI EORP ACS STEMI investigators group of the ESC[†]

¹Klinikum der Stadt Ludwigshafen and Institut für Herzinfarktforschung, Bremsersstraße 79, 67063 Ludwigshafen am Rhein, Germany; ²Institute of Cardiovascular Sciences, University of Birmingham, Edgbaston 1, Birmingham, B15 2TT, United Kingdom; ³Hôpital Européen Georges Pompidou, Service de Cardiologie, 20 Rue Leblanc, 75015 Paris, France; ⁴University Hospital Brno, Medical Faculty of Masaryk University, Dept of Internal Medicine and Cardiology, Kamenice 5, 625 00 Bohunice, Brno, Czech Republic; ⁵EURObservational Research Programme, European Society of Cardiology, European Heart House, 2035 Route des Colles, Sophia Antipolis, France; ⁶Cardiac Rehabilitation Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Shahid Rahmani Alley, Moshtagh Sevom Street, Isfahan, Iran; ⁷Interventional Cardiology Unit, Annunziata Civil Hospital, Via Migliori 1, 87100 Cosenza, Italy; ⁸Ain Shams University, Ramsis Street, Abbassia Square, Cairo, Egypt; ⁹Jagiellonian University Medical College; Institute of Cardiology, Department of Interventional Cardiology, John Paul II Hospital, Prądnicza 80, 31-202 Kraków, Poland; ¹⁰Rabin Medical Center, Ze'ev Jabotinsky St 39, Petah Tikva, 49100 Israel; ¹¹Department of Cardiac Sciences, King Fahad Cardiac Center, College of Medicine, King Saud University, Riyadh, Saudi Arabia; ¹²Cardiocenter, Third Faculty of Medicine Charles Univ. and Univ. Hospital Kralovske Ruská 2411, 100 00 Praha 10-Vinohrady, Czech Republic; ¹³Teaching Hospital of J.A. Reiman, Cardiocentre Prešov, Jána Hollého 5898/14, 081 81 Prešov, Slovakia; ¹⁴Centro Hospitalar e Universitário do Algarve, R. Leao Penedo, 8000-386 Faro, Portugal; ¹⁵Deputy-Director of the Heart and Vascular Center of Semmelweis University, Gaál József út 9, 1122 Budapest, Hungary; ¹⁶National and Kapodistrian University of Athens, Attikon University Hospital, Rimini 1, Chaidari 124 62, Greece; ¹⁷Chapidze Emergency Cardiology Center, 0159, Didube-Chugureti District, Nino Javakhishvili str., Tbilisi, Georgia; ¹⁸Faculty of Medicine, University of Belgrade, Belgrade, Serbia. Department of Cardiology, Clinical Center of Serbia, Pasterova 2, Beograd 11000 Belgrade, Serbia; ¹⁹Scientific Research Institute of Cardiology named after Levon Hovhannissyan, 0014, Yerevan, Paruyr Sevaki St., 5 Building (Kanaker-Zeytun adm. district), Yerevan, Armenia; ²⁰Cardiology I & Cardiology II, University Hospital Center Mother Theresa, Rruga e Dibrës 372 Tirana AL, 1000, Albania; ²¹Kyrgyz State Medical Academy, 92 Akhunbayev St, Bishkek, Kyrgyzstan; ²²National Center of Cardiology and Internal Medicine, Togolok Moldo Str., 720040, Bishkek, Kyrgyzstan; ²³Medical Faculty, University of Prishtina "Hasan Prishtina", University Clinical Centre of Kosova, Pristina 10000, Kosovo; ²⁴Iraqi Scientific Council of Cardiology, Baghdad Heart Center, Medical City, Baghdad, Iraq; ²⁵Vilnius University Faculty of Medicine Hospital Santaros Klinikos, 08406, Santariškių g. 2, Vilnius 08410, Lithuania; ²⁶Aalborg University Hospital, Department of Cardiology, Hobrovej 18-22, 9100 Aalborg, Denmark; ²⁷North Estonia Medical Centre, J. Sütiste tee 19, 13419 Tallinn, Estonia; ²⁸Mater Dei Hospital, Triq Dun Karm, L-Imnsida, MSD2090 Msida, Malta; ²⁹St Vincent's University Hospital, Merrion Rd, Dublin 4, D04 N2E0, Ireland; ³⁰Satakunta Central Hospital, Sairaalan tie 3, 28500 Pori, Finland; ³¹Spitalul Clinic de Urgenta "Floreasca", Calea Floreasca 8, București 014461, Romania; ³²Leeds Institute for Cardiovascular and Metabolic Medicine, University of Leeds School of Medicine, Worsley Building, University of Leeds, LS2 9JT Leeds, UK; ³³Maria Cecilia Hospital, GVM Care & Research, Via Corriera, 1, 48033 Cotignola RA, Italy; and ³⁴Hospital Rudolfstiftung, Juchgasse 25, 1030 Wien, Austria

Received 22 December 2020; revised 18 April 2021; editorial decision 19 May 2021; accepted 5 June 2021; online publish-ahead-of-print 14 August 2021

See page 4550 for the editorial comment for this article 'ST-segment elevation myocardial infarction management: great strides but still room for improvement', by C.L. Grines and S. Mehta, <https://doi.org/10.1093/eurheartj/ehab596>.

* Corresponding author. Tel: +49 561 980 2941, Email: uwe.zeymer@t-online.de

† The members of the ACVC-EAPCI EORP ACS STEMI Investigators Group of the ESC are listed in the [Appendix A](#).

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2021. For permissions, please email: journals.permissions@oup.com.

Aims

The aim of this study was to determine the contemporary use of reperfusion therapy in the European Society of Cardiology (ESC) member and affiliated countries and adherence to ESC clinical practice guidelines in patients with ST-elevation myocardial infarction (STEMI).


Methods and results

Prospective cohort (EURObservational Research Programme STEMI Registry) of hospitalized STEMI patients with symptom onset <24 h in 196 centres across 29 countries. A total of 11 462 patients were enrolled, for whom primary percutaneous coronary intervention (PCI) (total cohort frequency: 72.2%, country frequency range 0–100%), fibrinolysis (18.8%; 0–100%), and no reperfusion therapy (9.0%; 0–75%) were performed. Corresponding in-hospital mortality rates from any cause were 3.1%, 4.4%, and 14.1% and overall mortality was 4.4% (country range 2.5–5.9%). Achievement of quality indicators for reperfusion was reported for 92.7% (region range 84.8–97.5%) for the performance of reperfusion therapy of all patients with STEMI <12 h and 54.4% (region range 37.1–70.1%) for timely reperfusion.

Conclusions

The use of reperfusion therapy for STEMI in the ESC member and affiliated countries was high. Primary PCI was the most frequently used treatment and associated total in-hospital mortality was below 5%. However, there was geographic variation in the use of primary PCI, which was associated with differences in in-hospital mortality.

Graphical Abstract



Current use of reperfusion therapies, achievement of quality indicators for reperfusion therapies and in-hospital mortality in Europe, North Africa and the Middle East. In Europe, about 85% of patients with ST-elevation myocardial infarction are treated with primary percutaneous coronary intervention, while total in-hospital mortality is around 4%. ESC, European Society of Cardiology.

Keywords

ST-elevation myocardial infarction • Primary percutaneous coronary intervention • Observational studies • Reperfusion therapy

Introduction

The incidence of ST-elevation myocardial infarction (STEMI) in European countries ranges between 40 and 140/100 000/year, which equates to over 500 000 patients with STEMI being admitted each year.^{1,2} Despite improvements in management, STEMI remains one of the leading causes of death in Europe and worldwide.^{3–5} STEMI is defined by chest pain or equivalent symptoms and ST-segment elevation or left bundle branch block on the diagnostic electrocardiogram (ECG) and subsequent confirmation of diagnosis by elevation of cardiac troponin.^{1,6} The European Society of Cardiology (ESC) has issued practice guidelines for patients with STEMI, the two latest versions published in 2012 and 2017.^{1,6} It has been shown that adherence to these guidelines improves outcomes.^{1,6} The cornerstone of treatment of STEMI is acute reperfusion therapy preferably with primary percutaneous coronary intervention (PCI). Previous surveys on acute coronary syndromes (ACS) within the EuroHeart Survey programme collecting data of ACS presentation, treatment and outcome in Europe in 2000, 2004, and 2008 and the Snapshot Registry in 2009 showed gaps between guideline recommendations and their implementation into clinical practice.^{7–10} In addition, wide variations in the treatment of STEMI between countries have been noticed, especially in the use of primary PCI. The 'Stent for life' initiative of the ESC has been created to increase the rate of patients treated with primary PCI within Europe, the Mediterranean basin and other regions worldwide.^{11–13} Therefore, this registry aimed to evaluate the evolution of treatment of STEMI in ESC member countries, adherence to guidelines and outcomes.

Methods

The design and methods of the registry have been published.^{14,15} This study describes the demographic, clinical, and biological characteristics and outcomes of patients with STEMI admitted to cardiology centres in ESC member and affiliated countries.

Study organization

This registry is a joint initiative of the Association for Acute CardioVascular Care (ACVC) and the European Association of Percutaneous Cardiovascular Interventions (EAPCI) under the umbrella of the ESC EURObservational Research Programme (EORP). In each country, centres with and without PCI facilities were invited to participate. Each centre was asked to enrol at least 30 and up to 60 consecutive patients presenting with STEMI in the community within 24 h after symptom onset.

The study complies with the Declaration of Helsinki, the locally appointed ethics committees approved the research protocol, and informed consent was obtained from all subjects (or their legally authorized representatives).

Patients

Patients aged >18 years with an initial diagnosis of STEMI according to the 2012 ESC STEMI guidelines admitted within 24 h after symptom onset were identified on admission to the hospital, in the emergency room, or directly in the catheterization laboratory and given a unique study number.

Data

Baseline data included demographics, patient history, and risk factors. Reperfusion therapies, time intervals, and the reasons why reperfusion was not utilized were also collected. Angiographic data and details of the revascularization procedures were collected. Medications given in the pre-hospital phase, during hospitalization, and at discharge were documented. Follow-up for clinical events was performed until hospital discharge.

Regions

To compare characteristics, treatments, and outcomes between different regions, the entire cohort was divided into seven regions according to the definition of the World Health Organization: Northern Europe (Denmark, Estonia, Finland, Ireland, Lithuania), Eastern Europe (Armenia, Czech Republic, Georgia, Hungary, Kyrgyzstan, Poland, Romania, Russia, Slovakia), Western Europe (Austria, Germany), Southern Europe (Albania, Greece, Italy, Kosovo, Malta, Montenegro, Portugal, Serbia, Spain), Africa (Egypt), Middle East (Israel), and non-ESC-Middle East (Iran, Iraq, Saudi Arabia).¹⁵

Definitions

Cardiogenic shock was defined according to the ESC STEMI guidelines.¹ Bleeding complications were classified according to the Bleeding Academic Research Consortium (BARC) definition.¹⁶

Quality indicators for reperfusion therapy

To assess the quality of care regarding reperfusion therapy, the following parameters were evaluated:^{1,17} proportion of patients with STEMI <12 h who received reperfusion therapy; proportion of patients with timely reperfusion (fibrinolysis within 30 min after first medical contact and for patients with primary PCI admitted to centres with catheterization laboratories <60 min from door to PCI, for transfer patients: qualifying ECG to PCI <120 min).

Statistics

Descriptive statistics are used to summarize frequency tabulations (*n*, %) and distributions (mean, standard deviation). All the results are summarized overall and by type of reperfusion therapy. For categorical data, frequency tabulations are presented (without missing values if applicable).

Results

From 1 January 2015 to 31 March 2018, 11 462 patients in 196 centres of 29 countries were enrolled in the registry. The numbers of patients per country ranged from 5 to 1356.¹⁵ The baseline demographics of the entire cohort and patients in the seven regions are given in Table 1. There were variations with respect to age (North Africa 55.4 years, Northern Europe 64.2 years), female sex (Non-ESC Middle East 17.9%, Eastern Europe 28.7%), diabetes (Northern Europe 15.9%, North Africa 40.8%), smoking (Northern Europe 36.8%, North Africa 59.1%), and hypertension (North Africa 37.2%, Eastern Europe 54.0%) between the seven regions. Anterior STEMI was present in 49.1% of the patients without significant differences among regions. The other relevant clinical findings on admission are summarized in Table 2. Less than 5% of patients presented with cardiogenic shock.

Table 1 Baseline characteristics of the total cohort and the seven World Health Organization regions

	Total	Northern Europe	Eastern Europe	Western Europe	Southern Europe	North Africa	Middle East	Non-ESC Middle East
N	11 462	242	3846	270	2613	1356	756	2379
Age (years), N = 11 334	61.0 ± 12.8	64.2 ± 12.3	62.8 ± 12.2	62.3 ± 11.9	63.3 ± 12.7	55.4 ± 11.3	60.4 ± 14.3	58.7 ± 13.0
Women, N = 11 462	2643	67	1102	59	595	250	144	426
	23.1%	27.7%	28.7%	21.9%	22.8%	18.4%	19.0%	17.9%
Prior myocardial infarction	1264/10 417	27/239	557/3765	29/252	252/2574	107/1346	NA	292/2241
	12.1%	11.3%	14.8%	11.5%	9.8%	7.9%		13.0%
Previous PCI	1152/11 374	30/239	360/3822	31/257	247/2603	96/1353	171/753	217/2347
	10.1%	12.6%	9.4%	12.1%	9.5%	7.1%	22.7%	9.2%
Previous CABG	193/11 405	11/242	55/3827	4/262	33/2609	11/1355	22/753	57/2357
	1.7%	4.5%	1.4%	1.5%	1.3%	0.8%	2.9%	2.4%
Previous stroke/TIA	599/11 375	10/240	260/3815	12/257	104/2606	54/1354	41/755	118/2348
	5.3%	4.2%	6.8%	4.7%	4.0%	4.0%	5.4%	5.0%
Peripheral artery disease	470/10 883	10/236	244/3508	14/254	127/2532	12/1342	28/755	35/2256
	4.3%	4.2%	7.0%	5.5%	5.0%	0.9%	3.7%	1.6%
History of atrial fibrillation	451/11 348	19/242	226/3811	17/259	113/2605	21/1355	35/754	20/2322
	4.0%	7.9%	5.9%	6.6%	4.3%	1.5%	4.6%	0.9%
Diabetes	2996/11 204	38/239	817/3802	49/263	579/2607	547/1341	235/751	731/2201
	26.7%	15.9%	21.5%	18.6%	22.2%	40.8%	31.3%	33.2%
Current smoker	5123/11 204	86/234	1582/3708	123/258	1090/2585	799/1353	350/756	1093/2310
	45.7%	36.8%	42.7%	47.7%	42.2%	59.1%	46.3%	47.3%
Hypercholesterolemia	3770/9792	113/218	1288/2931	100/253	1051/2486	141/1111	482/750	595/2043
	38.5%	51.8%	43.9%	39.5%	42.3%	12.7%	64.3%	29.1%
Treated hypertension	5348/11 167	114/239	2019/3737	127/260	1334/2599	503/1351	398/750	853/2231
	47.9%	47.7%	54.0%	48.8%	51.3%	37.2%	53.1%	38.2%
Cancer or other life limiting diseases	224/11 234	5/238	93/3719	5/253	59/2584	9/1348	41/745	12/2347
	2.0%	2.1%	2.5%	2.0%	2.3%	0.7%	5.5%	0.5%

CABG, coronary artery bypass graft; ESC, European Society of Cardiology; PCI, percutaneous coronary intervention; TIA, transient ischaemic attack.

Table 2 Clinical findings on admission in the total population and the seven World Health Organization regions

	Total	Northern Europe N = 242	Eastern Europe N = 3846	Western Europe N = 270	Southern Europe N = 2613	North Africa N = 1356	Middle East N = 756	Non-ESC Middle East N = 2379
Anterior STEMI	5260/10 706 49.1%	113/242 46.7%	1772/3846 46.1%	119/270 44.1%	1201/2613 46.0%	774/1356 57.1%	NA	1281/2379 53.8%
Other STEMI	5387/10 706 50.3%	128/242 52.9%	2048/3846 53.3%	151/270 55.9%	1399/2613 53.5%	577/1356 42.6%	NA	1084/2379 45.6%
LBBS	56/10 706 0.5%	1/242 0.4%	24/3846 0.6%	0/270 0.0%	12/2613 0.5%	5/1356 0.4%	NA	14/2379 0.6%
Atrial fibrillation on qualifying ECG Killip class	571/11 457 5.0%	21/242 8.7%	276/3846 7.2%	19/270 7.0%	132/2613 5.1%	47/1356 3.5%	22/756 2.9%	54/2374 2.3%
I	9158/11 419 80.2%	177/242 73.1%	2874/3846 74.7%	219/270 81.1%	2086/2602 80.2%	1137/1356 83.8%	652/724 90.1%	2013/2379 84.6%
II	1388/11 419 12.2%	40/242 16.5%	639/3846 16.6%	31/270 11.5%	309/2602 11.9%	134/1356 9.9%	40/724 5.5%	195/2379 8.2%
III	425/11 419 3.7%	12/242 5.0%	162/3846 4.2%	6/270 2.2%	94/2602 3.6%	46/1356 3.4%	15/724 2.1%	90/2379 3.8%
IV	448/11 419 3.9%	13/242 5.4%	171/3846 4.4%	14/270 5.2%	113/2602 4.3%	39/1356 2.9%	17/724 2.3%	81/2379 3.4%
Heart rate (b.p.m.), N = 11 429	79.8 ± 20.0	75.5 ± 19.1	79.0 ± 20.0	77.1 ± 20.8	77.8 ± 19.1	84.8 ± 19.4	80.4 ± 18.8	81.1 ± 21.2
Mean systolic blood pressure (mmHg), N = 11 431	133.2 ± 28.0	133.7 ± 29.3	135.4 ± 27.8	138.7 ± 29.5	133.7 ± 28.2	125.5 ± 25.8	138.1 ± 28.7	131.2 ± 28.0
Out-of-hospital cardiac arrest	482/10 963 4.4%	13/242 5.4%	255/3846 6.6%	22/269 8.2%	84/2613 3.2%	34/1356 2.5%	34/258 13.2%	40/2379 1.7%

ECG, electrocardiogram; LBBS, left bundle branch block; STEMI, ST-elevation myocardial infarction.




Figure 1 Rate of reperfusion therapies performed in the total cohort and the seven World Health Organization regions. ESC, European Society of Cardiology; PCI, percutaneous coronary intervention.

Reperfusion therapy

The intended treatment was primary PCI in the admission centre in 7338 (68.4%) cases of STEMI, transfer out for primary PCI at another hospital in 353 (3.1%), fibrinolysis in 1999 (17.4%), no acute reperfusion therapy in 630 (5.5%), and not determined in 643 (5.6%). Treatment actually received was primary PCI in 72.2% (country range 0–100%), fibrinolysis in 18.8% (country range 0–100%), and no acute reperfusion therapy in 9.0% (country range 0–75%). While primary PCI was performed in over 80% of patients in the ESC member countries in Europe and in 75% in the ESC Middle East, the rate was around 50% in North Africa and the non-ECS Middle East countries (Figure 1). In the European ESC members countries, the rates of primary PCI, fibrinolysis and no reperfusion therapy were 85.4% (5955/6971), 7.6% (530/6971), and 7.0% (486/6971). The reasons for not performing acute reperfusion therapy in the 1027 patients of the total cohort were as follows: clinically inappropriate (19.4%), contraindication to anticoagulation/antiplatelet therapy (5.2%), late presentation (38.9%), spontaneous reperfusion (15.6%), wrong diagnosis (3.7%), patient refusal (5.1%), and others (12.2%).

The mean time interval between symptom onset and first medical contact was 221.6 min and the mean time between first medical contact and primary PCI was 195.2 min (Table 3). Primary PCI was performed in 61.8% within 120 min after first medical contact. The majority of patients (55.8%) were admitted by an ambulance or emergency medical service, the remaining admissions were self-presenters, while 66.8% of patients presented directly to a PCI hospital.

Medical therapy

The acute antithrombotic therapies used according to reperfusion treatment are summarized in Table 4. The use of aspirin was over 97% and the most widely used P2Y₁₂ inhibitor was clopidogrel.

Intravenous antiplatelet agents were given in about 19% of patients, predominantly glycoprotein IIb/IIIa inhibitors. With respect to anticoagulation, unfractionated heparin was most frequently used followed by low molecular weight heparins, while bivalirudin and fondaparinux were administered only rarely. Beta-blockers were given in 86.4% of patients, angiotensin-converting enzyme inhibitors in 77.7% and angiotensin receptor blockers in 7.6%, sacubitril/valsartan in 0.1%, mineralocorticoid receptor antagonists in 16.7%, and ivabradine in 3.7%. With respect to low-density lipoprotein-lowering therapy a statin was given in 96.4% of patients, ezetimibe in 0.8% and fibrates in 0.3%. Proton pump inhibitors were prescribed to 70.3% of patients.

In-hospital procedures and outcomes

Median length of stay was 5 days and ranged between 3 days (North Africa) and 6 days (Eastern, Western, and Southern Europe). Emergency coronary artery bypass graft surgery was performed in only 88 (0.8%) patients. Additional revascularization procedures after day one were done in 11.3% of patients and are listed in Table 5. In addition, 10.0% of patients were scheduled for a staged PCI procedure after discharge.

In patients treated with primary PCI, radial access was used in 4815, femoral access in 3923 and both in 97 patients. Corresponding BARC 2–5 bleeding complications were 2.0%, 3.6%, and 9.2%, and in-hospital mortality rates 1.9%, 4.8%, and 7.2%, respectively.

A total of 212 (9.8%) of 2046 patients treated with fibrinolysis received PCI within 24 h after admission. In-hospital mortality was 1.0% vs. 4.8% compared to patients without early PCI.

Overall in-hospital mortality was 4.4%. It varied from 2.5% to 5.9% across the seven regions (Supplementary material online, Figure S1) and was 3.8% in the European and Middle East ESC member countries ($n = 7727$). In-hospital mortality in patients with the different reperfusion strategies is shown in Figure 2. Mortality in patients with

Table 3 Time intervals between symptom onset, first medical contact and percutaneous coronary intervention in the total population and the seven World Health Organization regions

	Total	Northern Europe	Eastern Europe	Western Europe	Southern Europe	North Africa	Middle East	Non-ESC Middle East
Symptom-onset to FMC (min), N = 11 214	221.6 ± 460.6	249.9 ± 812.5	257.3 ± 486.7	233.0 ± 408.7	227.6 ± 531.3	201.6 ± 255.6	251.1 ± 765.9	158.0 ± 218.0
FMC to PCI (min), N = 8165	195.2 ± 1105.6	227.5 ± 389.9	197.6 ± 565.2	193.9 ± 906.9	215.6 ± 1948.7	147.1 ± 241.6	131.9 ± 179.9	197.2 ± 237.4
FMC to PCI <30 min	291/8165 3.6%	2/214 0.9%	78/3197 2.4%	4/259 1.5%	100/2264 4.4%	38/663 5.7%	20/485 4.1%	49/1083 4.5%
FMC to PCI <60 min	1690/8165 20.7%	20/214 9.3%	598/3187 18.7%	60/259 23.2%	456/2264 20.1%	286/663 43.1%	106/485 21.9%	164/1083 15.1%
FMC to PCI <120 min	5044/8165 61.8%	119/214 55.6%	2028/3197 63.4%	174/259 67.2%	1404/2264 62.0%	466/663 70.3%	348/485 71.8%	505/1083 46.6%

FMC, first medical contact; PCI, percutaneous coronary intervention.

Table 4 Antithrombotic therapy according to reperfusion therapy during the first 24 h in the entire cohort and according to initial reperfusion therapy

	Total N = 11 462	Primary PCI N = 8275	Fibrinolysis N = 2160	No reperfusion therapy N = 1027
Aspirin	11 151/11 449 97.4%	8053/8263 97.5%	2141/2160 99.1%	957/1026 93.3%
Clopidogrel	7486/11 450 65.4%	4674/8263 56.6%	2054/2160 95.1%	758/1027 73.8%
Prasugrel	1246/11 450 10.9%	1171/8264 14.2%	25/2160 1.2%	50/1026 4.9%
Ticagrelor	2705/11 451 23.6%	2482/8264 30.0%	94/2160 4.4%	129/1027 12.6%
Dual antiplatelet therapy	10 995/11 448 96.0%	7967/8262 96.4%	2134/2160 98.8%	894/1026 87.1%
GP IIb/IIIa inhibitors	2078/10 696 19.4%	1864/7697 24.2%	172/2145 8.0%	42/854 4.9%
Cangrelor	8/10 696 0.1%	7/7697 0.1%	1/2145 0.05%	0/854 0%
Unfractionated heparin	7300/11 442 63.8%	5771/8255 69.9%	1010/2160 46.8%	519/1027 50.5%
Low molecular weight heparin	4253/11 443 37.2%	2634/8256 31.9%	1189/2160 55.0%	430/1027 41.9%
Bivalirudin	134/11 345 1.2%	128/8187 1.6%	4/2160 0.2%	2/998 0.2%
Fondaparinux	222/11 340 2.0%	163/8181 2.0%	33/2160 1.5%	26/999 2.6%

GP, glycoprotein; PCI, percutaneous coronary intervention.

cardiogenic shock was 10-fold higher than in patients without shock (35.5% vs. 3.1%) (Supplementary material online, Figure S1). Mechanical complications were reported in 0.7% of patients. Definite or probable stent thrombosis occurred in 1.2%, and a re-infarction in 1.0%. Cerebrovascular accidents were rare and mostly ischaemic. A BARC bleeding complication was reported in 5.8% of patients and a transfusion was given in 2.1% (Table 5). The in-hospital mortality of the 372 (3.2%) patients with BARC 2–5 bleeding complications was 15.3% vs. 4.0% in the 11 083 patients with no or BARC-1 bleeding complications. A total of 2584 (24.1%) patients experienced heart failure during the index hospitalization. The mean left ventricular ejection fraction was lower in the heart failure cohort compared to patients without heart failure ($39.9 \pm 11.6\%$ vs. $48.1 \pm 10.3\%$), and the corresponding in-hospital mortality was 13.5% vs. 1.4%. Patients with atrial fibrillation on the qualifying ECG ($n = 571$, 5%) received primary PCI in 73.7% vs. 72.1%, fibrinolysis in 13.7% vs. 19.1% and no early reperfusion therapy in 12.6% vs. 8.8%. During hospital stay, they had more often stroke (2.8% vs. 0.9%) and a higher in-hospital mortality (13.0% vs. 3.9%).

Achievement of quality indicators for reperfusion therapy

The proportion of patients reperfused among eligible patients with STEMI <12 h ranged from 84.8% to 97.5% in the seven regions (Figure

3), while timely reperfusion was achieved in 54.4% (region range 37.1–70.1%).

Discussion

The major finding of the current EORP STEMI registry is that primary PCI was the preferred reperfusion therapy in patients with STEMI in Europe and the participating Middle East countries. In the first EuroHeart Survey ACS registry in 2001, primary PCI was used in about 20% of patients, while in the current analysis the PCI rate in European ESC member countries has risen to about 80% (Figure 3). This increase may in part be due to the success of the 'Stent for Life' initiative of the ESC¹¹ and also the very strong guideline recommendation that primary PCI should be the preferred approach for the treatment of STEMI.^{1,6} In contrast, the use of fibrinolytic therapy decreased over time from 35% in 2001 and to about 20% in the whole cohort and less than 10% in the European countries now (Figure 4). About 10% of patients still did not receive early reperfusion therapy, with late presentation as the most often reported reason. The rate of patients without reperfusion therapy in the ESC member countries has declined the last 20 years, starting with over 40% in the first EuroHeart Survey on ACS and <10% in the current study. As in previous reports, patients receiving no reperfusion therapy had a high in-hospital mortality, which was nearly 15% in this registry (Graphical abstract). Public campaigns are therefore still warranted to

help minimize patient-related pre-hospital delays, to reduce the rate of patient presenting too late to benefit from early reperfusion therapy. Nevertheless, there will always remain some patients who are not treated with reperfusion therapy for appropriate clinical reasons such as frailty or severe concomitant diseases.

The increase in the use of primary PCI has been accompanied by a decrease in mortality, from 7.0% in the year 2000 to 4.0% in the years 2016–2018, supporting the widespread use of this therapy in STEMI, as also described by others.¹⁹

There remain variations in the use of primary PCI between regions ranging from 46% in the non-ESC Middle East countries to 75% and 98% in the European ESC member countries. These variations were associated with differences in total in-hospital mortality, from 5.9% in

the Middle East non-ESC countries to 2.5% in Northern Europe. Therefore, all efforts should be made to increase the use of primary PCI in all regions to improve outcome in STEMI. As recommended in the ESC guidelines,^{1,6} this can be achieved by creating STEMI networks with clearly defined patient pathways.

The ESC guidelines recommend a time of <120 min between first medical contact and primary PCI. This goal was achieved in only about 60% of cases, while only 20% were treated in less than 60 min. Accordingly, logistics of care should be continuously improved to further reduce total ischaemic time.

Quality indicators for reperfusion therapy were met for over 90% of patients with STEMI <12 h, while timely reperfusion was achieved in only about half of the patients underscoring the need for the improvement of logistics.

Aspirin is still the cornerstone of antiplatelet therapy in patients with STEMI regardless of the initial reperfusion therapy and was used in over 95% of the patients. Despite the recommendations in favour of the newer P2Y12 inhibitors prasugrel and ticagrelor in the ESC guidelines, clopidogrel was the most often used second antiplatelet agent. The reasons might include limitations in the availability of prasugrel and ticagrelor and cost. With respect to anticoagulation, unfractionated heparin and low molecular weight heparins were the preferred drugs, while bivalirudin and fondaparinux were used only rarely.

Current guidelines recommend dual antiplatelet therapy regardless of the initial reperfusion therapy and 96% of the patients actually received dual antiplatelet therapy at discharge. Statins were given in almost 96% of patients, which documents the widespread acceptance of the low-density lipoprotein lowering as very important therapy in patients with STEMI.

In patients without cardiogenic shock, mortality was as low as 3.9%, while the occurrence of cardiogenic shock was still associated

Table 5 In-hospital additional procedures and complications

PCI after fibrinolysis	892/2160 (41.3%)
PCI after initial no reperfusion	285/1027 (27.8%)
Staged PCI after primary PCI	863/7644 (11.3%)
Total CABG	327/11 458 (2.9%)
Stent thrombosis	
Definite	106/11 416 (0.9%)
Probable	34/11 416 (0.3%)
Reinfarction	120/11 460 (1.0%)
Stroke	115/11 460 (1.0%)
Heart failure	2584/10 702 (24.1%)
Total BARC bleeding complications	667/11 455 (5.8%)
BARC 2–5 bleeding complications	372/11 455 (3.2%)

BARC, Bleeding Academic Research Consortium; CABG, coronary artery bypass graft; PCI, percutaneous coronary intervention.




Figure 2 In-hospital mortality in the total cohort and subgroups of patients treated with percutaneous coronary intervention, fibrinolysis, and no initial reperfusion therapy.




Figure 3 Achievement of quality indicators for reperfusion therapies. Rate of patients with reperfusion therapy (percutaneous coronary intervention or fibrinolysis) among all patients with ST-elevation myocardial infarction <12 h and rate of patients with timely reperfusion (fibrinolysis within 30 min after first medical contact and for patients with primary percutaneous coronary intervention admitted to centres with catheterization laboratory <60 min from door to arterial access, for transfer patients: qualifying electrocardiogram to percutaneous coronary intervention <120 min).




Figure 4 Temporal trends in the use of reperfusion therapies in patients with ST-elevation myocardial infarction in Europe over the last 20 years in the EuroHeart Surveys I to III, the EuroHeart Surveys Snapshot Survey and the EORP ST-elevation myocardial infarction registry.

with a high mortality of 35%. This was somewhat lower than the mortality reported in randomized trials including patients with cardiogenic shock, which approximates 40%.¹⁸ Nevertheless, it suggests that major improvements in total mortality rates can only be achieved if mortality due to cardiogenic shock will be reduced.¹⁹

Limitations

Despite the large number of patients included, the representativeness of the patient population for Europe was somewhat limited. The national sites were selected by the National Societies of Cardiology with the aim to provide a representative sample within the given

country. However, since participation was voluntary, a selection bias with participation of ‘better’ centres cannot be excluded and therefore the reality might be less favourable. The larger European countries such as France, Germany and the United Kingdom did not participate or enrol a sufficient number of patients.

Actions to be taken

The findings of this registry imply further steps to be seriously taken into consideration to improve STEMI quality of care: continuous public campaigns should be performed to raise awareness and reduce the interval between symptom onset and first medical contact; improvement of logistics should be implemented to minimize delays so as to increase the number of patients treated with primary PCI within 120 min or preferably 60 min after first medical contact. A further increase in the use of primary PCI could be achieved by implementing patient pathways within STEMI networks and future research is needed to reduce mortality in patients with cardiogenic shock to substantially improve overall mortality.

Supplementary material

Supplementary material is available at *European Heart Journal* online.

Acknowledgements

This study was supported by the EORP Oversight Committee, The Registry Executive Committee of the EURObservational Research Programme (EORP). Data collection was conducted by the EORP department from the ESC by Marème Konte and Florian Larras as data managers, Elin Folkesson Lefrancq as project officer, and Souad Mekhaldi as clinical project manager. Statistical analyses were performed by Cécile Laroche. Overall activities are coordinated by Aldo P. Maggioni (Scientific Coordinator EORP). This study was also supported by the Saudi Heart Association, The Deanship of Scientific Research at King Saud University, Riyadh, Saudi Arabia (Research group number: RG-1436-013).

Funding

Since the start of EORP, the following companies have supported the programme: Abbott Vascular Int. (2011–2021), Amgen Cardiovascular (2009–2018), AstraZeneca (2014–2021), Bayer AG (2009–2018), Boehringer Ingelheim (2009–2019), Boston Scientific (2009–2012), The Bristol Myers Squibb and Pfizer Alliance (2011–2019), Daiichi Sankyo Europe GmbH (2011–2020), The Alliance Daiichi Sankyo Europe GmbH and Eli Lilly and Company (2014–2017), Edwards (2016–2019), Gedeon Richter Plc. (2014–2016), Menarini Int. Op. (2009–2012), MSD-Merck & Co. (2011–2014), Novartis Pharma AG (2014–2020), ResMed (2014–2016), Sanofi (2009–2011), SERVIER (2009–2021), and Vifor (2019–2022).

Conflict of interest: U.Z. reports personal fees from Amgen, grants and personal fees from AstraZeneca, Abiomed, Bayer, and BMS, personal fees from Novartis, Sanofi, and Boehringer Ingelheim, grants and personal fees from Tommsdorf, personal fees from Pfizer and The Medicines Company, outside the submitted work. P.L. has nothing to disclose. N.D. reports grants, personal fees and non-financial support from Amgen, AstraZeneca, Bayer, and BMS, personal fees from Boehringer Ingelheim, Intercept, MSD, Novartis, Novo-Nordisk, and Pfizer, grants, personal fees,

and non-financial support from Sanofi, personal fees from Servier, and personal fees from UCB, outside the submitted work. P.K. reports personal fees from Boston Scientific, grants from Novartis, personal fees from Edwards Lifesciences, Bayer, and Chiesi, and grants from Sanofi-Aventis, outside the submitted work. C.L., R.C., S.M.S., K.F.A., M.S., J.M., D.B., D.A., Z.K., S.S., P.Z., A.G., E.M., G.B., H.A.-F., B.R., A.M.M., M.Q., P.P.K., M.S., P.Š. have nothing to disclose. J.L. reports personal fees from AstraZeneca, Bayer AG, Balton, Philips Volcano, Abbott, Procardia, and Bracco, outside the submitted work. Z.I. reports personal fees from Bayer, grants and personal fees from Pfizer, personal fees from Boehringer Ingelheim, AstraZeneca, and Medison Israel, grants and personal fees from Novo-Nordisk, and personal fees from Sanofi, outside the submitted work. Z.M. reports personal fees from AstraZeneca, Bayer, Pfizer, Boehringer Ingelheim, and Chiesi, outside the submitted work. T.M. reports personal fees from AstraZeneca Eesti, Sanofi Aventis, Berlin and Chemie Menarini, outside the submitted work. G.T.-C. reports personal fees from AstraZeneca, Pfizer, and Berlin Chemie, outside the submitted work. C.P.G. reports personal fees from AstraZeneca, personal fees from Bayer, personal fees from Daiichi Sankyo, grants from BMS, and grants from Abbott outside the submitted work. A.P.M. reports personal fees from Bayer, personal fees from Fresenius, and personal fees from Novartis, outside the submitted work. F.W. reports personal fees from AstraZeneca, outside the submitted work.

Data availability

Data request can be send to EORP at the European Heart House, Sophia Antipolis France.

Appendix A—ACS STEMI

Executive Committee

Franz Weidinger AT (Co-Chair), Uwe Zeymer DE (Co-Chair), Nicolas Danchin FR (Co-Chair), Peter Ludman GB (Co-Chair), Peter Sinnaeve, BE, Petr Kala, CZ, Roberto Ferrari, IT, Aldo P. Maggioni, IT.

Steering Committee

Artan Goda, AL; Parounak Zelveian, AM; Franz Weidinger, AT; Kiril Karamfilov, BG; Zuzana Motovska, CZ; Uwe Zeymer, DE; Bent Raungaard, DK; Toomas Marandi, EE; Sameh Mohamed Shaheen, EG; Rosa-Maria Lidon, ES; Pasi Paavo Karjalainen, FI; Zviad Kereselidze, GE; Dimitrios Alexopoulos, GR; David Becker, HU; Martin Quinn, IE; Zaza Iakobishvili, IL; Hasan Al-Farhan, IQ; Masoumeh Sadeghi, IR; Roberto Caporale, IT; Erkin Mirrakhimov, KG; Pranas Serpytis, LT; Andrejs Erglis, LV; Sasko Kedev, MK; Alice May Moore, MT; Dariusz Dudek, PL; Jacek Legutko, PL; Jorge Mimoso, PT; Gabriel Tatu-Chitoiu, RO; Sinisa Stojkovic, RS; Evgeny Shlyakhto, RU; Khalid F AlHabib, SA; Matjaz Bunc, SI; Martin Studencan, SK; Mohamed Sami Murali, TN; Gani Bajraktari, XK.

EORP Team

Marème Konte, Florian Larras, Elin Folkesson Lefrancq, Souad Mekhaldi, Cécile Laroche, Aldo P. Maggioni.

Investigators

Albania: *Tirana:* A. Goda, N. Shuka, E. Pavli, E. Tafaj, T. Gishto, A. Dibra, A. Duka, A. Gjana, A. Kristo, G. Knuti, A. Demiraj, E. Dado,

- E. Hasimi, L. Simoni, M. Siqeca, **Armenia:** Yerevan: H. Sisakian, Yerevan: H. Hayrapetyan, S. Markosyan, L. Galustyan, N. Arustamyan, H. Kzhdryan, S. Pepoyan, **Austria:** Graz: A. Zirkik, D. Von Lewinski, S. Paetzold, I. Kienzl, K. Matyas, **Krems an der Donau:** T. Neunteufl, M. Nikfardjam, U. Neuhold, A. Mihalcz, F. Glaser, **Linz:** C. Steinwender, C. Reiter, M. Grund, D. Hrnčić, **Salzburg:** U. Hoppe, M. Hammerer, L. Hinterbuchner, **Vienna:** C. Hengstenberg, **Vienna:** G. Delle Karth, I. Lang, **Vienna:** F. Weidinger, W. Winkler, M. Hasun, J. Kastner, C. Havel, M. Derntl, G. Oberegger, J. Hajos, C. Adlbrecht, T. Publig, M.-C. Leitgeb, R. Wilfing, P. Jirak, C.-Y. Ho, L. Puskas, L. Schrutka, **Czech Republic:** Brno: J. Spinar, J. Parenica, Brno: O. Hlinomaz, V. Fendrychova, J. Semenka, J. Sikora, J. Sitar, L. Groch, M. Rezek, M. Novak, P. Kramarikova, **Hradec Kralove:** J. Stasek, J. Dusek, P. Zdrahal, **Liberec:** R. Polasek, J. Karasek, J. Seiner, N. Sukova, **Pardubice:** I. Varvarovsky, T. Lazarák, V. Novotny, J. Matejka, **Plzen:** R. Rokyta, S. Volovar, **Prague:** J. Belohlavek, **Prague:** Z. Motovska, M. Siranec, M. Kamenik, R. Kralik, **Denmark:** Aalborg: B. Raungaard, J. Ravkilde, S.E. Jensen, A. Villadsen, K. Villefrance, C. Schmidt Skov, **Aarhus:** M. Maeng, K. Moeller, **Egypt:** Assiut: H. Hasan-Ali, T.A. Ahmed, **Aswan:** M. Hassan, A. ElGuindy, **Badr city:** M. Farouk Ismail, A. Ibrahim Abd El-Aal, A. El-sayed Gaafar, H. Magdy Hassan, M. Ahmed Shafie, M. Nabil El-khouly, **Benha:** A. Bendary, M. Darwish, **Beni Suef:** Y. Ahmed, O. A. Amin, A. AbdElHakim, K. Abosaif, **Cairo:** H. Kandil, **Cairo:** M.A.G. Galal, **Cairo:** E.E. El Hefny, **Cairo:** M. El Sayed, K. Aly, M. Mokarrab, **Cairo:** M. Osman, M. Abdelhamid, S. Mantawy, M.R. Ali, S.D. Kaky, V.A. Khalil, M.E.A. Saraya, A. Talaat, **Cairo:** M. Nabil, W.M. Mounir, K. Mahmoud A. Aransa, G. Kazamel, S. Anwar, A. Al-Habbaa, M. Abd el Monem, A. Ismael, M. Amin Abu-Sheishaa, M.M. Abd Rabou, T. M. A. Hammouda, **Elkharga:** M. Moaaz, **Fayoum:** K. Elkhassab, T. Ragab, **Fayoum:** A. Rashwan, A. Rmdan, G. AbdelRazek, H. Ebeid, H. Soliman Ghareeb, **Giza:** N. Farag, **Giza:** M. Zaki, M. Seleem, A. Torki, M. Youssef, N.A. AlLah Nasser, A. Rafaat, H. Selim, M.M. Makram, M. Khayyal, K. Malasi, A. Madkour, M. Kolib, H. Alkady, H. Nagah, **Mansoura:** M. Yossef, A. Wafa, E. Mahfouz, G. Faheem, M. Magdy Moris, **Marsa Matruh:** A. Ragab, M. Ghazal, A. Mabrouk, M. Hassan, **Tanta:** M. El-Masry, M. Naseem, S. Samir, **Estonia:** Tallinn: T. Marandi, J. Reinmets, M. Allvee, A. Saar, T. Ainla, A. Vaide, M. Kisseljova, U. Pakosta, **Tartu:** J. Eha, K. Lotamois, **Finland:** **Kokkola:** J. Sia, J. Myllymaki, T. Pinola, **Por:** P.P. Karjalainen, T. Paana, J. Mikkelsen, M. Ampio, **Georgia:** Tbilisi: J. Tsvilasvili, **Tbilisi:** P. Zurab, **Tbilisi:** Z. Kereselidze, R. Agladze, A. Melia, D. Gogoberidze, N. Khubua, L. Totladze, I. Metreveli, A. Chikovani, **Germany:** Lübeck: I. Eitel, J. Pöss, M. Werner, A. Constantz, C. Ahrens, **Ludwigshafen am Rhein:** U. Zeymer, H. Tolsdorf, S. Klinger, **Munich:** S. Sack, T. Heer, **Greece:** Athens: J. Lekakis, Athens: I. Kanakakis, I. Xenogiannis, K. Ermidou, N. Makris, A. Ntalianis, F. Katsaros, **Kerkyra:** E. Revi, **Chania:** K. Kafkala, E. Mihelakis, G. Diakakis, K. Grammatikopoulos, D. Voutsinos, **Patras:** D. Alexopoulos, I. Xanthopoulou, V. Mplani, **Piraeus:** S. Foussas, N. Papakonstantinou, N. Patsourakos, A. Dimopoulos, A. Derventzis, K. Athanasiou, **Thessaloniki:** V.P. Vassilikos, C. Papadopoulos, S. Tzikas, **Verioia:** I. Vogiatzis, A. Datsios, I. Galitsianos, K. Koutsampasopoulos, S. Grigoriadis, **Volos:** A. Douras, N. Baka, S. Spathis, T. Kyrlidis, H. Hatzinikolaou, **Hungary:** Budapest: R.G. Kiss, **Budapest:** D. Becker, F. Nowotta, K. Tóth, S. Szabó, C. Lakatos, **Gyula:** Z. Jambrik, J. Ruzsa, **Kecskemet:** Z. Ruzsa, S. Róna, J. Toth, A. Vargane Kosik, K.S.B. Toth, **Miskolc:** G.G. Nagy, Z. Ondrejko, Z. Körömi, B. Botos, **Iran:** Isfahan: M. Pourmoghadas, **Isfahan:** A. Salehi, **Isfahan:** G. Massoumi, M. Sadeghi, A. Soleimani, N. Sarrafzadegan, H. Roohafza, M. Azarm, A. Mirmohammadsadeghi, D. Rajabi, **Kermanshab:** Y. Rahmani, S. Siabani, F. Najafi, B. Hamzeh, H. Karim, H. Siabani, N. Saleh, H. Charehjo, L. Zamzam, **Iraq:** Baghdad: G. Al-Temimi, H. Al-Farhan, A. Al-Yassin, A. Mohammad, A. Ridha, G. Al-Saedi, N. Atabi, O. Sabbar, S. Mahmood, Z. Dakhil, I.F. Yaseen, **Nasseria:** M. Almyahi, H. Alkenzawi, T. Alkinani, A. Alyacopy, **Ireland:** Cork: P. Kearney, K. Twomey, **Israel:** **Petah Tikva:** Z. Iakobishvili, **Ramat Gan:** N. Shlomo, R. Beigel, **Italy:** Bari: P. Caldarola, D. Rutigliano, L. Sublimi Saponetti, N. Locuratolo, V. Palumbo, **Benevento:** M. Scherillo, D. Formigli, **Bergamo:** P. Canova, G. Musumeci, F. Roncali, **Brescia:** M. Metra, C. Lombardi, E. Visco, L. Rossi, **Cagliari:** L. Meloni, R. Montisci, V. Pippia, M.F. Marchetti, M. Congia, C. Cacace, **Caltagirone:** G. Luca, G. Boscarelli, **Catanzaro:** C. Indolfi, G. Ambrosio, A. Mongiardo, C. Spaccarotella, S. De Rosa, G. Canino, C. Critelli, **Cosenza:** R. Caporale, D. Chiappetta, F. Battista, **Fermo:** D. Gabrielli, A. Marziali, **Genoa:** P. Bernabò, **Guastalla:** A. Navazio, E. Guerri, F. Manca, **Lugo:** M. Gobbi, **Messina:** G. Oretto, G. Andò, S. Carerj, F. Saporito, M. Cimmino, **Mestre-Venice:** F. Rigo, G. Zuin, **Naples:** B. Tuccillo, F. Scotto di Uccio, L. Irace, **Nuoro:** G. Lorenzoni, I. Meloni, P. Merella, **Partinico:** G.M. Polizzi, R. Pino, **Pisa:** M. Marzilli, D. Morrone, P. Caravelli, E. Orsini, S. Mosa, **Rimini:** G. Piovaccari, A. Santarelli, C. Cavazza, **Rome:** F. Romeo, **Rome:** F. Fedele, M. Mancone, M. Straito, N. Salvi, P. Scarparo, P. Severino, C. Razzini, G. Massaro, A. Cinque, **Rome:** C. Gaudio, F. Barilla, C. Torromeo, L. Porco, M. Mei, R. Iorio, **Saronno:** D. Nassiacos, B. Barco, **Trieste:** G. Sinagra, L. Falco, L. Priolo, A. Perkan, **Kosovo:** **Mitrovica:** M. Strana, G. Bajraktari, L. Percuku, G. Berisha, B. Mziu, **Kyrgyzstan:** **Bishkek:** M. Beishenkulov, T. Abdurashidova, A. Toktosunova, K. Kaliev, **Lithuania:** Vilnius: P. Serpytis, R. Serpytis, E. Butkute, M. Lizaitis, M. Broslavskytė, **Malta:** Msida: R.G. Xuereb, A.M. Moore, M. Mercieca Balbi, E. Paris, L. Buttigieg, **Poland:** **Bialystok:** W. Musial, S. Dobrzycki, A. Dubicki, E. Kazimierzczak, A. Tycinska, **Katowice:** W. Wojakowski, B. Kalanska-Lukasik, A. Ochala, W. Wanh, S. Dworowy, **Kielce:** J. Sielski, M. Janion, A. Janion-Sadowska, **Krakow:** D. Dudek, J. Wojtasik-Bakalarz, L. Bryniarski, **Lodz:** J.Z. Peruga, M. Jonczyk, L. Jankowski, **Nowy Targ:** A. Klecha, J. Legutko, J. Michalowska, M. Brzezinski, T. Kozmik, T. Kowalczyk, J. Adamczuk, **Ostrowiec Swietokrzyski:** M. Maliszewski, P. Kuziemka, P. Plaza, A. Jaros, A. Pawelec, J. Sledz, **Oswiecim:** S. Bartus, W. Zmuda, M. Bogusz, M. Wisnicki, G. Szastak, M. Adamczyk, M. Suska, P. Czunko, **Warsaw:** G. Opolski, J. Kochman, M. Tomaniak, S. Miernik, K. Paczwa, **Warsaw:** A. Witkowski, M.P. Opolski, A.D. Staruch, **Zabrze:** Z. Kalarus, G. Honisz, G. Mencil, M. Swierad, T. Podolecki, **Portugal:** Braga: J. Marques, P. Azevedo, M.A. Pereira, A. Gaspar, **Coimbra:** S. Monteiro, F. Goncalves, L. Leite, **Faro:** J. Mimoso, W. Manuel Lopes dos Santos, J. Amado, **Funchal:** D. Pereira, B. Silva, G. Caires, M. Neto, R. Rodrigues, A. Correia, D. Freitas, **Guimaraes:** A. Lourenco, F. Ferreira, F. Sousa, J. Portugues, L. Calvo, F. Almeida, **Santarem:** M. Alves, A. Silva, **Setubal:** R. Caria, F. Seixo, **Romania:** Craiova: C. Militaru, E. Ionica, G. Tatu-Chitoiu, O. Istratoaie, M. Florescu, **Russian Federation:** Astrakhan: E. Lipnitkaia, **Belgorod:** O. Osipova, **Belgorod:** S. Konstantinov, V. Bukator, **Cheboksary:** T. Vinokur, E. Egorova, E. Nefedova, **Chelyabinsk:** S. Levashov, A. Gorbunova, M. Redkina, N. Karaulovskaya, **Cherkessk:** F. Bijieva, **Ekaterinburg:** N. Babich, O. Smirnova, R. Filyanin, S. Eseva, A. Kutluev, A. Chlopenova, A. Shtanko, E. Kuppar, E. Shaekhmurzhina, M. Ibragimova, M. Mullahmetova, M. Chepisova, M. Kuzminykh, **Gelendzhik:** M. Betkaraeva, A. Namitokov, **Kazan:** N. Khasanov, **Kazan:** L. Baleeva, Z. Galeeva, F. Magamedkerimova, E. Ivantsov, **Kemerovo:** E. Tavlueva, A. Kochergina, D. Sedykh, **Krasnodar:** E. Kosmachova, **Krasnodar:** V. Skibitskiy, N. Porodenko, A. Namitokov, K. Litovka,

E. Ulbasheva, *Krasnoyarsk*: S. Niculina, *Krasnoyarsk*: M. Petrova, E. Harkov, N. Tsybulskaya, A. Lobanova, A. Chernova, A. Kuskaeva, A. Kuskaev, *Moscow*: M. Ruda, *Moscow*: D. Zateyshchikov, *Moscow*: M. Gilarov, E. Konstantinova, O. Koroleva, A. Averkova, N. Zhukova, D. Kalimullin, *Nizhny Novgorod*: N. Borovkova, A. Tokareva, M. Buyanova, *Rostov-on-Don*: L. Khaisheva, A. Pirozhenko, *Saint Petersburg*: T. Novikova, *Saint Petersburg*: A. Yakovlev, *Saint Petersburg*: T. Tyurina, K. Lapshin, N. Moroshkina, M. Kiseleva, S. Fedorova, L. Krylova, *Samara*: D. Duplyakov, *Seversk*: Y. Semenova, A. Rusina, *Tomsk*: V. Ryabov, A. Syrkina, S. Demianov, *Tyumen*: O. Reitblat, A. Artemchuk, *Ulyanovsk*: E. Efremova, E. Makeeva, M. Menzorov, A. Shutov, N. Klimova, *Voronezh*: I. Shevchenko, O. Elistratova, O. Kostyuckova, R. Islamov, V. Budyak, E. Ponomareva, *Saudi Arabia*: Al-Kharj: U. Ullah Jan, *Alkhobar*: A.M. Alshehri, E. Sedky, Z. Alsahati, *Jeddah*: L. Mimish, *Jeddah*: A. Selem, *Jeddah*: A. Malik, O. Majeed, I. Altnji, *Khamis Mushayt*: M. AlShehri, *Riyadh*: A. Aref, *Riyadh*: K. AlHabib, *Riyadh*: M. AlDosary, *Riyadh*: S. Tayel, M. Abd AlRahman, K.N. Asfina, *Tabuk*: G. Abdin Hussein, *Tabuk*: M. Butt, *Serbia*: *Belgrade*: N. Markovic Nikolic, *Belgrade*: S. Obradovic, N. Djenic, M. Brajovic, A. Davidovic, R. Romanovic, V. Novakovic, M. Dekleva, M. Spasic, B. Dzudovic, Z. Jovic, D. Cvijanovic, S. Veljkovic, *Sremska Kamenica*: I. Ivanov, M. Cankovic, M. Jarakovic, M. Kovacevic, M. Trajkovic, *Zajecar*: V. Mitov, A. Jovic, *Slovakia*: *Banska Bystrica*: M. Hudec, M. Gombasky, *Bratislava*: J. Sumbal, A. Bohm, E. Baranova, *Martin*: F. Kovar, M. Samos, J. Podoba, *Nitra*: P. Kurray, T. Obona, A. Remenarikova, B. Kollarik, D. Verebova, G. Kardosova, *Presov*: M. Studencan, D. Alusik, J. Macakova, M. Kozlej, *Spain*: *Barcelona*: A. Bayes-Genis, *Barcelona*: A. Sionis, C. Garcia Garcia, *Barcelona*: R-M. Lidon, A. Duran Cambra, C. Labata Salvador, F. Rueda Sobella, J. Sans Rosello, M. Vila Perales, T. Oliveras Vila, M. Ferrer Massot, J. Bañeras, *Galdakao*: I. Lekuona, G. Zugazabeitia, *Madrid*: A. Fernandez-Ortiz, A. Viana Tejedor, C. Ferrera, V. Alvarez, *Pontevedra*: O. Diaz-Castro, R.M. Agra-Bermejo, C. Gonzalez-Cambeiro, E. Gonzalez-Babarro, J. Domingo-Del Valle, *Santander*: N. Royuela, V. Burgos, A. Canteli, C. Castrillo, M. Cobo, M. Ruiz, *Santiago de Compostela*: E. Abu-Assi, J.M. Garcia Acuna.

References

1. Steg PG, James SK, Atar D, Badano LP, Blömostrom-Lundqvist C, Borger MA, Di Mario C, Dickstein K, Ducrocq G, Fernandez-Aviles F, Gershlick AH, Giannuzzi P, Halvorsen S, Huber K, Juni P, Kastrati A, Knuuti J, Lenzen MJ, Mahaffey KW, Valgimigli M, van 't Hof A, Widimsky P, Zahger D; Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology (ESC). ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2012;**33**:2569–2619.
2. Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, Bax JJ, Borger MA, Brotons C, Chew DP, Gencer B, Hasenfuss G, Kjeldsen K, Lancellotti P, Landmesser U, Mehilli J, Mukherjee D, Storey RF, Windecker S; ESC Scientific Document Group. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2016;**37**:267–315.
3. Anderson JL, Morrow DA. Acute myocardial infarction. *N Engl J Med* 2017;**376**:2053–2064.
4. Puymirat E, Simon T, Cayla G, Cottin Y, Elbaz M, Coste P, Lemesle G, Motreff P, Popovic B, Khalife K, Labèque JN, Perret T, Le Ray C, Orion L, Jouve B, Blanchard D, Peycher P, Silvain J, Steg PG, Goldstein P, Guéret P, Belle L, Aissaoui N, Ferrières J, Schiele F, Danchin N; USIK, USIC 2000, and FAST-MI investigators. Acute myocardial infarction: changes in patient characteristics, management, and 6-month outcomes over a period of 20 years in the FAST-MI Program (French Registry of Acute ST-Elevation or Non-ST-elevation Myocardial Infarction) 1995 to 2015. *Circulation* 2017;**136**:1908–1919.
5. Terkelsen CJ, Lassen JF, Nørgaard BL, Gerdes JC, Jensen T, Gøtzsche LB, Nielsen TT, Andersen HR. Mortality rates in patients with ST-elevation vs. non-ST-elevation acute myocardial infarction: observations from an unselected cohort. *Eur Heart J* 2005;**26**:18–26.
6. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, Caforio ALP, Crea F, Goudevenos JA, Halvorsen S, Hindricks G, Kastrati A, Lenzen MJ, Prescott E, Roffi M, Valgimigli M, Varenhorst C, Vranckx P, Widimský P; ESC Scientific Document Group. 2017 ESC guidelines on the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2018;**39**:119–177.
7. Hasdai D, Behar S, Wallentin L, Danchin N, Gitt AK, Boersma E, Fioretti PM, Simoons ML, Battler A. A prospective survey of the characteristics, treatments and outcomes of patients with acute coronary syndromes in Europe and the Mediterranean basin: the Euro Heart Survey on Acute Coronary Syndromes (Euro Heart Survey ACS). *Eur Heart J* 2002;**23**:1190–1201.
8. Mandelzweig L, Battler A, Boyko V, Bueno H, Danchin N, Filippatos G, Gitt A, Hasdai D, Hasin Y, Marrugat J, Van de Werf F, Wallentin L, Behar S; Euro Heart Survey Investigators. The second Euro Heart Survey on Acute Coronary Syndromes: characteristics, treatment, and outcome of patients with ACS in Europe and the Mediterranean Basin in 2004. *Eur Heart J* 2006;**27**:2285–2293.
9. Schiele F, Hochadel M, Tubaro M, Meneveau N, Wojakowski W, Gierlotka M, Polonski L, Bassand JP, Fox KA, Gitt AK. Reperfusion strategy in Europe: temporal trends in performance measures for reperfusion therapy in ST-elevation myocardial infarction. *Eur Heart J* 2010;**31**:2614–2624.
10. Puymirat E, Battler A, Birkhead J, Bueno H, Clemmensen P, Cottin Y, Fox KA, Gorenk B, Hamm C, Huber K, Lettino M, Lindahl B, Müller C, Parkhomenko A, Price S, Quinn T, Schiele F, Simoons M, Tatu-Chitoiu G, Tubaro M, Vrints C, Zahger D, Zeymer U, Danchin N; EHS 2009 Snapshot Participants. Euro Heart Survey 2009 Snapshot: regional variations in presentation and management of patients with AMI in 47 countries. *Eur Heart J Acute Cardiovasc Care* 2013;**2**:359–370.
11. Kaifoszova Z, Kala P, Alexander T, Zhang Y, Huo Y, Snyders A, Delport R, Alcocer-Gamba MA, Gavidia LM. Stent for Life Initiative: leading example in building STEMI systems of care in emerging countries. *EuroIntervention* 2014;**10** Suppl T:T87–95.
12. Widimsky P, Wijns W, Fajadet J, de Belder M, Knot J, Aaberge L, Andrikopoulos G, Baz JA, Betriu A, Claeys M, Danchin N, Djambazov S, Erne P, Hartikainen J, Huber K, Kala P, Klineva M, Kristensen SD, Ludman P, Ferre JM, Merkely B, Milicic D, Morais J, Noc M, Opolski G, Ostojic M, Radovanovic D, De Servi S, Stenestrand U, Studencan M, Tubaro M, Vasiljevic Z, Weidinger F, Witkowski A, Zeymer U; European Association for Percutaneous Cardiovascular Interventions. Reperfusion therapy for ST elevation acute myocardial infarction in Europe: description of the current situation in 30 countries. *Eur Heart J* 2010;**31**:943–957.
13. Kristensen SD, Laut KG, Fajadet J, Kaifoszova Z, Kala P, Di Mario C, Wijns W, Clemmensen P, Agladze V, Antoniadis L, Alhabib KF, De Boer MJ, Claeys MJ, Deleanu D, Dudek D, Erglis A, Gilard M, Goktekin O, Guagliumi G, Gudnason T, Hansen KW, Huber K, James S, Janota T, Jennings S, Kajander O, Kanakakis J, Karamfiloff KK, Kedev S, Kornowski R, Ludman PF, Merkely B, Milicic D, Najafov R, Nicolini FA, Noč M, Ostojic M, Pereira H, Radovanovic D, Sabaté M, Sobhy M, Sokolov M, Studencan M, Terzic I, Wahler S, Widimsky P; European Association for Percutaneous Cardiovascular Interventions. Reperfusion therapy for ST elevation acute myocardial infarction 2010/2011. Current status in 37 ESC countries. *Eur Heart J* 2014;**35**:1957–1970.
14. Zeymer U, Ludman P, Danchin N, Kala P, Maggioni AP, Weidinger F. Background and design of the ACCA-EAPCI registry on ST-segment elevation myocardial infarction of the European Society of Cardiology. *Eur Heart J Acute Cardiovasc Care* 2019;**8**:63–67.
15. Zeymer U, Ludman P, Danchin N, Kala P, Maggioni AP, Weidinger F; ACS STEMI Investigators. The ESC ACCA EAPCI EORP acute coronary syndrome ST-elevation myocardial infarction registry. *Eur Heart J Qual Care Clin Outcomes* 2020;**6**:100–104.
16. Mehran R, Rao SV, Bhatt DL, Gibson CM, Caixeta A, Eikelboom J, Kaul S, Wiwiot SD, Menon V, Nikolsky E, Serebruany V, Valgimigli M, Vranckx P, Taggart D, Sabik JF, Cutlip DE, Krucoff MV, Ohman EM, Steg PG, White H. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation* 2011;**123**:2736–2747.

17. Schiele F, Gale CP, Bonnefoy E, Capuano F, Claeys MJ, Danchin N, Fox KA, Huber K, Iakobishvili Z, Lettino M, Quinn T, Rubini Gimenez M, Bøtker HE, Swahn E, Timmis A, Tubaro M, Vrints C, Walker D, Zahger D, Zeymer U, Bueno H. Acute Cardiovascular Care Association. Quality indicators for acute myocardial infarction: a position paper of the Acute Cardiovascular Care Association. *Eur Heart J Acute Cardiovasc Care* 2017;**6**:34–59.
18. Zeymer U, Bueno H, Granger CB, Hochman J, Huber K, Lettino M, Price S, Schiele F, Tubaro M, Vranckx P, Zahger D, Thiele H. Acute Cardiovascular Care Association position statement for the diagnosis and treatment of patients with acute myocardial infarction complicated by cardiogenic shock: a document of the Acute Cardiovascular Care Association of the European Society of Cardiology. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:183–197.
19. Dondo TB, Hall M, Munyombwe T, Wilkinson C, Yadegarfar ME, Timmis A, Batin PD, Jernberg T, Fox KA, Gale CP. A nationwide causal mediation analysis of survival following ST-elevation myocardial infarction. *Heart* 2020;**106**:765–771.