













# Ticagrelor monotherapy in patients with chronic kidney disease undergoing percutaneous coronary intervention: TWILIGHT-CKD

Giulio G. Stefanini <sup>1,2†</sup>, Carlo Briguori <sup>3†</sup>, Davide Cao <sup>4</sup>, Usman Baber <sup>5</sup>, Samantha Sartori <sup>3</sup>, Zhongjie Zhang <sup>3</sup>, George Dangas <sup>4</sup>, Dominick J. Angiolillo <sup>6</sup>, Shamir Mehta <sup>7</sup>, David J. Cohen <sup>8,9</sup>, Timothy Collier <sup>10</sup>, Dariusz Dudek <sup>11</sup>, Javier Escaned <sup>12</sup>, C. Michael Gibson <sup>13</sup>, Robert Gil <sup>14</sup>, Kurt Huber <sup>15</sup>, Upendra Kaul <sup>16</sup>, Ran Kornowski <sup>17</sup>, Mitchell W. Krucoff <sup>18</sup>, Vijay Kunadian <sup>19</sup>, David J. Moliterno <sup>20</sup>, E. Magnus Ohman <sup>18</sup>, Keith G. Oldroyd <sup>21</sup>, Gennaro Sardella <sup>22</sup>, Samin K. Sharma <sup>4</sup>, Richard Shlofmitz <sup>9</sup>, Giora Weisz <sup>23</sup>, Bernhard Witzenbichler <sup>24</sup>, Stuart Pocock <sup>10</sup> and Roxana Mehran <sup>4\*</sup>

<sup>1</sup>Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan 20090, Italy; <sup>2</sup>IRCCS Humanitas Research Hospital, Rozzano, Milan 20089, Italy; <sup>3</sup>Mediterranea Cardiocentro, Napoli 80122, Italy; <sup>4</sup>The Zena and Michael A. Wiener Cardiovascular Institute, Mount Sinai Hospital, One Gustave L. Levy Place, Box 1030, New York, NY 10029-6574, USA; <sup>5</sup>The University of Oklahoma Health Sciences Center, Oklahoma City, OK 73104, USA; <sup>6</sup>University of Florida College of Medicine, Jacksonville, FL 32209, USA; <sup>7</sup>Hamilton Health Sciences, Hamilton, ON L8N 3Z5, Canada; <sup>8</sup>Cardiovascular Research Foundation, New York, NY 10019, USA; <sup>9</sup>St. Francis Hospital, Roslyn, NY 11576, USA; <sup>10</sup>London School of Hygiene and Tropical Medicine, London WC1E 7HT, UK; <sup>11</sup>Jagiellonian University Medical College, Krakow 31-008, Poland; <sup>12</sup>Instituto de Investigacion Sanitaria del Hospital Clinico San Carlos and Complutense University, Madrid 28040, Spain; <sup>13</sup>Beth Israel Deaconess Medical Center, Boston, MA 02215, USA; <sup>14</sup>Center of Postgraduate Medical Education, Central Clinical Hospital of the Ministry of Interior and Administration, Warsaw 02-507, Poland; <sup>15</sup>Wilhelminen hospital, Wien 1160, Austria; <sup>16</sup>Batra Hospital and Medical Research Centre, New Delhi 110062, India; <sup>17</sup>Rabin Medical Center, Petah Tikva 49100, Israel; <sup>18</sup>Duke University Medical Center-Duke Clinical Research Institute, Durham, NC 27710, USA; <sup>19</sup>Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University and Freeman Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne NE7 7DN, UK; <sup>20</sup>University of Kentucky, Lexington, KY 40506, USA; <sup>21</sup>The West of Scotland Heart and Lung Centre, Golden Jubilee National Hospital, Clydebank G81 4DY, UK; <sup>22</sup>Policlinico Umberto I University, Roma 00161, Italy; <sup>23</sup>New York Presbyterian Hospital, Columbia University Medical Center, New York, NY 10032, USA; and <sup>24</sup>Helios Amper-Klinikum, Dachau 85221, Germany

Received 13 April 2021; revised 7 June 2021; editorial decision 5 July 2021; accepted 26 July 2021; online publish-ahead-of-print 23 August 2021

See page 4694 for the editorial comment for this article ‘Ticagrelor monotherapy in CKD: better safety at what price?’, by W.A.E. Parker and R.F. Storey, <https://doi.org/10.1093/eurheartj/ehab607>.

## Aims

The aim of this study was to assess the impact of chronic kidney disease (CKD) on the safety and efficacy of ticagrelor monotherapy among patients undergoing percutaneous coronary intervention (PCI).

## Methods and results

In this prespecified subanalysis of the TWILIGHT trial, we evaluated the treatment effects of ticagrelor with or without aspirin according to renal function. The trial enrolled patients undergoing drug-eluting stent implantation who fulfilled at least one clinical and one angiographic high-risk criterion. Chronic kidney disease, defined as an estimated glomerular filtration rate (eGFR) <60 mL/min/1.73 m<sup>2</sup>, was a clinical study entry criterion. Following a 3-month period of ticagrelor plus aspirin, event-free patients were randomly assigned to aspirin or placebo on top of ticagrelor for an additional 12 months. Of the 6835 patients randomized and with available eGFR at baseline, 1111 (16.3%) had CKD. Ticagrelor plus placebo reduced the primary endpoint of Bleeding Academic Research Consortium (BARC) type 2, 3, or 5 bleeding as compared with ticagrelor plus aspirin in both patients with [4.6% vs. 9.0%; hazard ratio (HR) 0.50, 95% confidence interval (CI) 0.31–0.80] and without (4.0% vs. 6.7%; HR 0.59, 95% CI 0.47–0.75;  $P_{\text{interaction}} = 0.508$ ) CKD, but the absolute risk reduction was greater in the former group. Rates of

\* Corresponding author. Tel: +1 212 659 9649, Fax: +1 646 537 8547, Email: [roxana.mehran@mountsinai.org](mailto:roxana.mehran@mountsinai.org)

† The first two authors contributed equally to the study.


Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2021. For permissions, please email: [journals.permissions@oup.com](mailto:journals.permissions@oup.com).

death, myocardial infarction, or stroke were not significantly different between the two randomized groups irrespective of the presence (7.9% vs. 5.7%; HR 1.40, 95% CI 0.88–2.22) or absence of (3.2% vs. 3.6%; HR 0.90, 95% CI 0.68–1.20;  $P_{\text{interaction}} = 0.111$ ) CKD.

Conclusion

Among CKD patients undergoing PCI, ticagrelor monotherapy reduced the risk of bleeding without a significant increase in ischaemic events as compared with ticagrelor plus aspirin.

Graphical Abstract



Keywords

Chronic kidney disease • Ticagrelor monotherapy • Aspirin • Bleeding • Thrombosis • PCI

Introduction

Impaired renal function is an established risk factor for incident and recurrent coronary events with cardiovascular disease being the leading cause of death in patients with chronic kidney disease (CKD).<sup>1–3</sup> The degree of CKD severity is associated with a progressive increase in the risk of both thrombotic and bleeding complications.<sup>4–6</sup> The pathophysiology behind these observations is multifactorial and relates to abnormalities of both platelet function and the coagulation cascade. As a result, the clinical implications of antithrombotic therapies may be different in patients with CKD as compared to those with normal renal function.<sup>7,8</sup>

A combination of aspirin and P2Y<sub>12</sub> inhibitor, commonly referred as dual antiplatelet therapy (DAPT), is the standard of care after percutaneous coronary intervention (PCI). Dual antiplatelet therapy effectively prevents ischaemic events, including stent thrombosis, but at the cost of an increase in bleeding.<sup>9</sup> Bleeding complications have been shown to negatively correlate with

patient survival after PCI, thereby putting the ischaemic benefits of DAPT in jeopardy.<sup>10–12</sup> This risk–benefit trade-off is further enhanced by more potent P2Y<sub>12</sub> inhibitors (such as prasugrel and ticagrelor), reflecting the incremental extent of platelet inhibition.<sup>13</sup>

Recently, a strategy of ticagrelor monotherapy after a short course of DAPT has emerged as an alternative treatment for high-risk patients undergoing PCI. In the TWILIGHT (Ticagrelor With Aspirin or Alone in High-Risk Patients after Coronary Intervention) trial, this approach was shown to significantly reduce clinically relevant bleeding without compromising antithrombotic efficacy.<sup>14</sup> It is unknown whether the clinical benefits of ticagrelor monotherapy are preserved in patients with CKD undergoing PCI, who—being intrinsically characterized by high thrombotic and bleeding risks—might particularly benefit from an optimization of antithrombotic strategies. Therefore, we conducted a prespecified analysis of the TWILIGHT trial to examine the safety and efficacy of ticagrelor monotherapy according to renal function.

## Methods

### Trial design and oversight

TWILIGHT was a randomized, placebo-controlled trial conducted at 187 sites in 11 countries from July 2015 through July 2019. The trial rationale, design, and principal results have been reported previously.<sup>15</sup> TWILIGHT was designed, coordinated, and sponsored by The Icahn School of Medicine at Mount Sinai. AstraZeneca provided an investigator-initiated grant and supplied ticagrelor for the trial but had no role in the design, collection, analysis, or interpretation of the data. The executive and steering committees were responsible for trial conduct, integrity of data analysis, and reporting of results. National regulatory agencies and institutional review boards or ethics committees of participating centres approved the trial protocol. An independent data safety monitoring board provided external oversight to ensure the safety of trial participants.

### Trial population

Patients undergoing successful PCI with at least one drug-eluting stent were eligible if they had at least one clinical and one angiographic feature associated with a high risk of ischaemic or bleeding events. Chronic kidney disease, defined as an estimated glomerular filtration rate  $<60$  mL/min/1.73 m<sup>2</sup>, was a clinical study entry criterion; other clinical criteria included age  $\geq 65$  years, female sex, troponin positive acute coronary syndrome (ACS), atherosclerotic vascular disease [prior myocardial infarction (MI), coronary revascularization or peripheral arterial disease], and diabetes mellitus requiring medication. Angiographic criteria included multivessel coronary artery disease, total stent length  $>30$  mm, thrombotic target lesion, bifurcation lesion requiring two stents, obstructive left main or proximal left anterior descending lesion, and calcified target lesion requiring debulking devices. Key exclusion criteria included dialysis-dependent renal failure, presentation with ST-elevation MI, cardiogenic shock, prior stroke, or need for oral anticoagulation.

All enrolled patients received open-label ticagrelor (90 mg twice daily) and enteric-coated aspirin (81–100 mg daily) after the index PCI. At 3 months, patients without major bleeding or ischaemic events were randomized 1:1 in a double-blind fashion to aspirin or matching placebo for 12 months in addition to open-label ticagrelor. Patients who experienced Bleeding Academic Research Consortium (BARC) type 3b or higher bleeds or ischaemic events (stroke, MI, or coronary revascularization) between the index PCI and 3 months were not eligible for randomization. Moreover, patients were ineligible for randomization if they were non-adherent to ticagrelor or aspirin. Randomization was performed using a secure web-based system; an independent statistician not involved with the trial generated the allocation sequence, which was stratified by site with randomly varying block sizes of 4, 6, or 8. Follow-up occurred 1 month after randomization via telephone and in-person at 6 and 12 months after randomization. After 12 months of protocol-mandated therapy, patients were switched to a standard-of-care antiplatelet regimen at the discretion of their treating physician followed by final telephone follow-up 3 months later.

### Endpoints

The primary endpoint was the composite of BARC type 2, 3, or 5 bleeding up to 1 year after randomization. The key secondary endpoint was the composite of all-cause death, MI, or stroke. Secondary bleeding endpoints included BARC types 3 or 5 bleeding; Thrombolysis in Myocardial Infarction (TIMI) major or minor bleeding; Global Use of Strategies to Open Occluded Arteries (GUSTO) moderate, severe, or life-threatening bleeding; or major bleeding as defined by the International Society on Thrombosis and Hemostasis (ISTH).<sup>16–19</sup> Other secondary endpoints

included cardiovascular death, non-fatal MI, ischaemic stroke, and definite or probable stent thrombosis. MI was defined according to the third universal definition, and revascularization and stent thrombosis were classified according to the Academic Research Consortium.<sup>20,21</sup> All clinical events were adjudicated by an independent committee, blinded to treatment assignment.

### Renal function assessment

Laboratory tests were performed locally at each site and collected during the enrolment procedure. Renal function was assessed using the most recent value of serum creatinine preceding index PCI, up to 4 weeks before. Glomerular filtration rate (eGFR) was estimated according to the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation.<sup>22</sup> The prespecified eGFR cut-point to define CKD was  $<60$  mL/min/1.73 m<sup>2</sup>. The Kidney Disease Outcomes Quality Initiative of the National Kidney Foundation classification was used to further stratify patients into mild to moderate (stage 3a: eGFR 45–59 mL/min/1.73 m<sup>2</sup>) or moderate to severe CKD (stage  $\geq 3b$ : eGFR  $<45$  mL/min/1.73 m<sup>2</sup>).<sup>23</sup>

### Statistical analyses

Baseline characteristics and clinical outcomes were evaluated in relation to renal function. In the primary prespecified analysis, the treatment effects of ticagrelor monotherapy vs. ticagrelor plus aspirin were evaluated in patients with CKD (eGFR  $<60$  mL/min/1.73 m<sup>2</sup>) and without CKD (eGFR  $\geq 60$  mL/min/1.73 m<sup>2</sup>), with formal interaction testing to assess for effect modification. Patients with unknown eGFR were excluded from the analysis. Clinical and procedural features are summarized by CKD status and randomized group using means (standard deviation) and frequencies for continuous and categorical variables, respectively. The cumulative incidence of primary and secondary endpoints was estimated using the Kaplan–Meier method. Patients without a primary endpoint between randomization and 1 year after randomization were censored at the time of death, last known contact, or 365 days, whichever came first. Hazard ratios (HRs) and 95% confidence intervals (CIs) were generated using Cox proportional hazards models.

Exploratory analyses were performed to examine the effects of ticagrelor monotherapy in the following clinically relevant subgroups within the CKD cohort: age ( $\geq 65$  vs.  $<65$  years), sex (male vs. female), body mass index (above vs. below median), non-ST-segment elevation ACS indication for PCI, prior MI, diabetes mellitus, anaemia, and multivessel disease. Clinical outcomes were also evaluated according to the degree of CKD severity using eGFR as a three-level categorical variable ( $<45$ , 45–59, and  $\geq 60$  mL/min/1.73 m<sup>2</sup>). In addition, cubic splines fitted with four equally spaced knots were used to plot the 1-year rate of ischaemic and bleeding events according to eGFR as a continuous variable in the overall population and in the two treatment arms, separately. For all analyses, bleeding outcomes were assessed in the intention-to-treat cohort, while ischaemic outcomes were analysed using the per-protocol cohort. A two-sided *P*-value of  $<0.05$  was considered statistically significant. All analyses were performed using Stata version 16.0 (StataCorp., College Station, TX, USA).

## Results

### Patient characteristics

Baseline serum creatinine levels were not available in 284 (4.0%) of the 7119 randomized patients. Therefore, the final cohort for the present analysis comprised 6835 patients, 1111 (16.3%) of whom had CKD (Supplementary material online, Figure S1). Of these patients, 554 patients (49.9%) were randomly assigned to ticagrelor plus

**Table 1** Baseline clinical characteristics

|                               | CKD (eGFR < 60) (n = 1111)          |                                     |         | No CKD (eGFR ≥ 60) (n = 5724)        |                                      |         |
|-------------------------------|-------------------------------------|-------------------------------------|---------|--------------------------------------|--------------------------------------|---------|
|                               | Tica + Placebo<br>(n = 554) (49.9%) | Tica + Aspirin<br>(n = 557) (50.1%) | P-value | Tica + Placebo<br>(n = 2856) (49.9%) | Tica + Aspirin<br>(n = 2868) (50.1%) | P-value |
| Age (years)                   | 70.1 ± 9.1                          | 70.6 ± 8.7                          | 0.370   | 62.1 ± 9.7                           | 61.9 ± 9.9                           | 0.626   |
| Female sex                    | 154 (27.8%)                         | 193 (34.6%)                         | 0.014   | 662 (23.2%)                          | 627 (21.9%)                          | 0.233   |
| Non-white race                | 113 (20.4%)                         | 113 (20.3%)                         | 0.964   | 954 (33.4%)                          | 920 (32.1%)                          | 0.285   |
| BMI (kg/m <sup>2</sup> )      | 28.9 ± 5.8                          | 29.3 ± 6.0                          | 0.228   | 28.5 ± 5.5                           | 28.4 ± 5.5                           | 0.534   |
| Enrolling region              |                                     |                                     | 0.163   |                                      |                                      | 0.574   |
| North America                 | 271 (48.9%)                         | 300 (53.9%)                         |         | 1174 (41.1%)                         | 1156 (40.3%)                         |         |
| Europe                        | 215 (38.8%)                         | 186 (33.4%)                         |         | 966 (33.8%)                          | 1008 (35.1%)                         |         |
| Asia                          | 68 (12.3%)                          | 71 (12.7%)                          |         | 716 (25.1%)                          | 704 (24.5%)                          |         |
| Diabetes                      | 251 (45.3%)                         | 270 (48.5%)                         | 0.290   | 1013 (35.5%)                         | 992 (34.6%)                          | 0.485   |
| Diabetes treated with insulin | 82 (32.7%)                          | 115 (42.6%)                         | 0.020   | 240 (23.7%)                          | 248 (25.0%)                          | 0.495   |
| Anaemia                       | 193 (35.2%)                         | 205 (37.1%)                         | 0.508   | 473 (16.7%)                          | 444 (15.6%)                          | 0.260   |
| Current smoker                | 71 (12.8%)                          | 77 (13.8%)                          | 0.621   | 629 (22.0%)                          | 720 (25.1%)                          | 0.006   |
| Hypercholesterolaemia         | 382 (69.0%)                         | 412 (74.0%)                         | 0.064   | 1697 (59.4%)                         | 1667 (58.1%)                         | 0.320   |
| Hypertension                  | 468 (84.5%)                         | 473 (84.9%)                         | 0.838   | 2010 (70.4%)                         | 2006 (70.0%)                         | 0.735   |
| Peripheral arterial disease   | 66 (11.9%)                          | 69 (12.4%)                          | 0.809   | 172 (6.0%)                           | 166 (5.8%)                           | 0.707   |
| Previous MI                   | 156 (28.2%)                         | 162 (29.1%)                         | 0.733   | 820 (28.7%)                          | 818 (28.5%)                          | 0.874   |
| Previous PCI                  | 265 (47.8%)                         | 264 (47.4%)                         | 0.884   | 1187 (41.6%)                         | 1182 (41.2%)                         | 0.789   |
| Previous CABG                 | 99 (17.9%)                          | 86 (15.4%)                          | 0.277   | 255 (8.9%)                           | 253 (8.8%)                           | 0.883   |
| Multivessel CAD               | 391 (70.6%)                         | 381 (68.4%)                         | 0.431   | 1779 (62.3%)                         | 1734 (60.5%)                         | 0.155   |
| Previous major bleed          | 6 (1.1%)                            | 9 (1.6%)                            | 0.442   | 23 (0.8%)                            | 22 (0.8%)                            | 0.870   |
| Indication for PCI            |                                     |                                     | 0.294   |                                      |                                      | 0.250   |
| Stable CAD                    | 244 (44.0%)                         | 228 (40.9%)                         |         | 974 (34.1%)                          | 937 (32.7%)                          |         |
| NSTEMI-ACS                    | 310 (56.0%)                         | 329 (59.1%)                         |         | 1881 (65.9%)                         | 1930 (67.3%)                         |         |

BMI, body mass index; CABG, coronary artery bypass graft; CAD, coronary artery disease; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; MI, myocardial infarction; NSTEMI-ACS, non-ST-elevation acute coronary syndrome; PCI, percutaneous coronary intervention; Tica, ticagrelor.

placebo and 557 (50.1%) to ticagrelor plus aspirin. Baseline clinical and procedural characteristics of patients with and without CKD are summarized in [Supplementary material online, Tables S1 and S2](#). Baseline characteristics were balanced across treatment groups, except for a lower prevalence of female sex (27.8% vs. 34.6%;  $P=0.014$ ) and insulin-dependent diabetes mellitus (32.7% vs. 42.6%;  $P=0.020$ ) in CKD patients receiving ticagrelor plus placebo and a lower prevalence of smokers (22.0% vs. 25.1%;  $P=0.006$ ) in patients without CKD receiving ticagrelor plus placebo, respectively, compared with those on ticagrelor plus aspirin ([Table 1](#)). There were no significant differences in procedural characteristics between randomized treatment groups ([Table 2](#)).

In the cohort of CKD patients, adherence to ticagrelor at 1 year was similar among those randomized to ticagrelor plus placebo compared with ticagrelor plus aspirin (80.5% vs. 83.0%;  $P=0.288$ ). Corresponding rates of adherence to blinded study drug were 74.7% and 78.7%, respectively ( $P=0.110$ ). Among patients without CKD, adherence rates to ticagrelor were higher in the placebo group (88.6% vs. 86.6%;  $P=0.025$ ), while there were no significant differences with respect to study drug (84.7% vs. 83.0%;  $P=0.087$ ).

### Bleeding events

During the trial, a total of 74 (6.8%) primary endpoint events were reported in patients with CKD as compared with 302 (5.4%) in those

without CKD ( $P=0.052$ ) ([Supplementary material online, Figure S2A](#)). As shown in [Figure 1](#), in the CKD cohort, the primary endpoint of BARC 2, 3, or 5 bleeding occurred in 25 patients (4.6%) randomized to ticagrelor plus placebo vs. 50 patients (9.0%) randomized to ticagrelor plus aspirin (HR 0.50, 95% CI 0.31–0.80;  $P=0.004$ ). Treatment effects on BARC 2, 3, or 5 bleeding were consistent among patients without CKD (4.0% vs. 6.7%; HR 0.59, 95% CI 0.47–0.75;  $P<0.001$ ) with no evidence of heterogeneity ( $P_{\text{interaction}}=0.508$ ). Ticagrelor plus placebo resulted in lower bleeding rates also with respect to more severe BARC 3 or 5 bleeding events and across different bleeding scales, including TIMI, GUSTO, and ISTH ([Figure 2](#)). There was no significant interaction between CKD and treatment arm for any of the bleeding endpoints (all  $P_{\text{interaction}}>0.1$ ), but the absolute risk difference in BARC 2, 3, or 5 bleeding associated with ticagrelor monotherapy was greater in the CKD group (−4.4%, 95% CI −7.3% to −1.4%) compared with the no CKD group (−2.7%, 95% CI −3.8% to −1.5%).

### Ischaemic events

A total of 74 (6.8%) key secondary endpoint events were reported in patients with CKD as compared with 190 (3.4%) in those without CKD ( $P<0.001$ ) ([Supplementary material online, Figure S2B](#)). In the CKD cohort, the key secondary endpoint of all-cause death, MI, or stroke occurred in 43 patients (7.9%) randomized to ticagrelor plus



**Table 2** Baseline procedural characteristics

|                                      | CKD (eGFR < 60) (n = 1111)          |                                     |         | No CKD (eGFR ≥ 60) (n = 5724)        |                                      |         |
|--------------------------------------|-------------------------------------|-------------------------------------|---------|--------------------------------------|--------------------------------------|---------|
|                                      | Tica + Placebo<br>(n = 554) (49.9%) | Tica + Aspirin<br>(n = 557) (50.1%) | P-value | Tica + Placebo<br>(n = 2856) (49.9%) | Tica + Aspirin<br>(n = 2868) (50.1%) | P-value |
| Radial artery access                 | 375 (67.7%)                         | 347 (62.3%)                         | 0.060   | 2124 (74.4%)                         | 2141 (74.7%)                         | 0.807   |
| Multivessel CAD                      | 391 (70.6%)                         | 381 (68.4%)                         | 0.431   | 1779 (62.3%)                         | 1734 (60.5%)                         | 0.155   |
| Target vessel                        |                                     |                                     |         |                                      |                                      |         |
| Left main                            | 34 (6.1%)                           | 33 (5.9%)                           | 0.882   | 117 (4.1%)                           | 148 (5.2%)                           | 0.055   |
| LAD                                  | 290 (52.3%)                         | 283 (50.8%)                         | 0.608   | 1611 (56.4%)                         | 1652 (57.6%)                         | 0.362   |
| LCX                                  | 199 (35.9%)                         | 183 (32.9%)                         | 0.282   | 906 (31.7%)                          | 924 (32.2%)                          | 0.688   |
| RCA                                  | 193 (34.8%)                         | 204 (36.6%)                         | 0.534   | 998 (34.9%)                          | 991 (34.6%)                          | 0.757   |
| Number of vessels treated            | 1.3 ± 0.5                           | 1.3 ± 0.5                           | 0.327   | 1.3 ± 0.5                            | 1.3 ± 0.5                            | 0.080   |
| Number of lesions treated            | 1.5 ± 0.8                           | 1.5 ± 0.7                           | 0.365   | 1.5 ± 0.7                            | 1.5 ± 0.8                            | 0.821   |
| Lesion morphology <sup>a</sup>       |                                     |                                     |         |                                      |                                      |         |
| Moderate/severe calcification        | 97 (17.5%)                          | 111 (19.9%)                         | 0.301   | 374 (13.1%)                          | 360 (12.6%)                          | 0.539   |
| Bifurcation                          | 66 (11.9%)                          | 49 (8.8%)                           | 0.088   | 341 (11.9%)                          | 365 (12.7%)                          | 0.365   |
| Total occlusion                      | 25 (4.5%)                           | 28 (5.0%)                           | 0.688   | 189 (6.6%)                           | 184 (6.4%)                           | 0.757   |
| Thrombotic                           | 48 (8.7%)                           | 50 (9.0%)                           | 0.854   | 312 (10.9%)                          | 317 (11.1%)                          | 0.876   |
| Total stent length <sup>b</sup> (mm) | 39.6 ± 25.7                         | 38.1 ± 22.6                         | 0.310   | 40.0 ± 23.9                          | 39.9 ± 24.6                          | 0.787   |
| Minimum stent diameter (mm)          | 2.8 ± 0.5                           | 2.8 ± 0.5                           | 0.764   | 2.8 ± 0.5                            | 2.9 ± 0.5                            | 0.320   |

CAD, coronary artery disease; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; LAD, left anterior descending coronary artery; LCX, left circumflex artery; RCA, right coronary artery; Tica, ticagrelor.

<sup>a</sup>Lesion morphology assessed by operators.

<sup>b</sup>Stent length calculated by operators.

placebo vs. 31 patients (5.7%) randomized to ticagrelor plus aspirin (HR 1.40, 95% CI 0.88–2.22;  $P = 0.153$ ) for an absolute risk difference of 2.2% (95% CI -0.8 to 5.2%) (Figure 3). Individual rates of MI (5.0% vs. 3.9%), ischaemic stroke (0.9% vs 0.0%), and definite/probable stent thrombosis (0.9% vs. 0.6%) were numerically higher with ticagrelor plus placebo but not significantly different between treatment groups, with the exception of ischaemic stroke ( $P = 0.025$ ) (Figure 4). Similar treatment effects on ischaemic events were observed in patients without CKD for the composite of all-cause death, MI, or stroke (3.2% vs. 3.6%; absolute risk difference -0.4%, 95% CI -1.3% to 0.6%; HR 0.90, 95% CI 0.68–1.20;  $P = 0.477$ ) and other secondary end-points, with no significant interaction between CKD status and treatment arm (all  $P_{\text{interaction}} > 0.1$ ).

## Exploratory analyses

Ticagrelor monotherapy reduced the risk of BARC 2, 3, or 5 bleeding without any increase in death, MI, or stroke across different subgroups of CKD patients, with no evidence of effect modification driven by the presence of additional risk features (Supplementary material online, Table S3 and S4).



Of the 1111 patients with an eGFR <60 mL/min/1.73 m<sup>2</sup>, 796 (71.6%) had mild-to-moderate CKD (eGFR 45–59 mL/min/1.73 m<sup>2</sup>) while 315 (28.4%) had moderate-to-severe CKD (eGFR < 45 mL/min/1.73 m<sup>2</sup>). There was a graded relationship between the severity of CKD and the risk of bleeding and ischaemic events (Supplementary material online, Figure S3), but the treatment effects of ticagrelor monotherapy on the primary and key secondary endpoints were preserved across all CKD categories (Supplementary material online, Tables S5 and S6).


When the estimated event rates were plotted against eGFR used as a continuous variable, the absolute risk reduction in BARC 2, 3, or 5 bleeding with ticagrelor monotherapy progressively increased with worsening renal function (Supplementary material online, Figure S4A). Meanwhile, the rates of death, MI, or stroke were numerically lower with ticagrelor monotherapy compared with ticagrelor plus aspirin for eGFR values >80 mL/min/1.73 m<sup>2</sup> and numerically higher below the same eGFR cut-off (Supplementary material online, Figure S4B).

## Discussion

The principal findings from this prespecified subgroup analysis of the TWILIGHT trial suggest that the treatment effects of ticagrelor monotherapy on bleeding and ischaemic outcomes observed in the overall trial are preserved irrespective of CKD. Of note, withdrawing aspirin after 3 months of DAPT with ticagrelor reduced clinically relevant BARC 2, 3, or 5 bleeding and major BARC 3 or 5 bleeding by 50% and 60%, respectively, among patients with CKD. This translated into an absolute reduction in bleeding risk more pronounced in patients with CKD than in those without CKD. Furthermore, ticagrelor monotherapy, as compared with ticagrelor plus aspirin, was not associated with significant differences in the composite outcome of all-cause death, MI, or stroke, despite a numerical increase in the rates of ischaemic events with worsening renal function (Graphical abstract).

Chronic kidney disease is a prevalent comorbid condition in patients undergoing PCI, an epidemiology that reflects the






**Figure 3** Rates of death, myocardial infarction, or stroke at 1 year after randomization. Kaplan–Meier curves for all-cause death, myocardial infarction, or stroke with ticagrelor plus placebo vs. ticagrelor plus aspirin in patients with and without CKD (estimated glomerular filtration rate < 60 mL/min/1.73 m<sup>2</sup>) in the per-protocol cohort. CI, confidence interval; CKD, chronic kidney disease; HR, hazard ratio; MI, myocardial infarction.

progressive ageing of the general population as well as the broadening of the indications to PCI to higher risk cohorts. The increased haemorrhagic and thrombotic risk profile of patients with CKD has been extensively characterized, with clinical studies demonstrating a gradient in the rates of adverse events that parallels the degree of renal dysfunction.<sup>4–6</sup> Several homeostatic modifications, including impaired platelet-vessel interaction, platelet aggregation and secretion abnormalities, and a pro-coagulant state with higher levels of fibrinogen and tissue factor and reduced anti-thrombin activity, have been implicated in these clinical manifestations.<sup>24</sup> Accelerated atherosclerosis, systemic inflammation, and oxidative stress are other key contributors to the enhanced cardiovascular risk associated with CKD.<sup>25,26</sup> Additional pharmacological issues relating to altered pharmacodynamic response, drug accumulation, and modified drug interactions further compound the management of these patients.<sup>7,8</sup>

In the early era of DAPT, subgroup analyses from randomized trials suggested that the benefits of adjunctive therapy with clopidogrel over aspirin alone were attenuated in patients with renal dysfunction.<sup>27,28</sup> Similar findings were reported in those with diabetic nephropathy randomized to monotherapy with clopidogrel instead of aspirin for secondary cardiovascular prevention.<sup>29</sup> This apparent lack of benefit was partly attributed to the higher levels of platelet reactivity observed in patients with CKD during treatment with clopidogrel, thus providing a rationale for the use of alternative antithrombotic regimens in this cohort.<sup>30,31</sup>

Compared with clopidogrel, potent P2Y<sub>12</sub> inhibitors (prasugrel and ticagrelor) have demonstrated a significant benefit in terms of ischaemic protection among ACS patients, including those with eGFR < 60 mL/min/1.73 m<sup>2</sup>. In the PLATO trial, the absolute and relative risk reduction for the primary endpoint of death, MI, or stroke associated with ticagrelor was enhanced in patients with CKD, although significant interaction with renal function was only achieved in a sensitivity analysis with the MDRD (Modification of Diet in Renal Disease) equation.<sup>32</sup> Yet, the incremental antithrombotic efficacy of ticagrelor over clopidogrel was evaluated on a background aspirin therapy, and this treatment combination also generated a remarkable increase in severe bleeding, which is known to negatively affect survival to a similar extent as thrombotic events.<sup>11,12</sup>

The idea that withdrawing aspirin after a short course of DAPT could reduce bleeding without compromising antithrombotic efficacy upon continuation of ticagrelor alone recently started to emerge.<sup>33</sup> This approach was also supported by pharmacodynamic data suggesting a marginal antiplatelet effect of aspirin when added to potent P2Y<sub>12</sub> inhibitors.<sup>34,35</sup> Hence, a number of PCI trials have investigated the safety and efficacy of P2Y<sub>12</sub> inhibitor monotherapy after a DAPT course as short as 1–3 months.<sup>36–39</sup> In GLOBAL LEADERS, which randomized nearly 16 000 all-comer patients to ticagrelor monotherapy after 1-month DAPT vs. standard of care, the prevalence of CKD was 13.7%.<sup>40</sup> The trial results showed no significant differences in the primary endpoint of all-cause mortality or new Q-wave MI and BARC 3 or 5 bleeding at 2 years between the experimental and the



**Figure 4** Risk of ischaemic events at 1 year after randomization. Forest plots showing the effect of ticagrelor plus placebo vs. ticagrelor plus aspirin on the ischaemic endpoints in relation to CKD (estimated glomerular filtration rate < 60 mL/min/1.73 m<sup>2</sup>). Event rates at 1 year were estimated using the Kaplan–Meier method. Hazard ratios and 95% confidence intervals (CI) with interaction *P*-values generated using Cox regression. In the CKD cohort, the stroke rate was 0.9% (five events) in patients randomized to ticagrelor plus placebo vs. 0% (no events) in those randomized to ticagrelor plus aspirin (log-rank *P* = 0.025, hazard ratio not applicable); Corresponding stroke rates in patients without CKD were 0.4% (11 events) vs. 0.3% (8 events) (log-rank *P* = 0.481; hazard ratio 1.38; 95% CI 0.55–3.43). ASA, aspirin; CI, confidence interval; CKD, chronic kidney disease; CV, cardiovascular; HR, hazard ratio; MI, myocardial infarction, ST, stent thrombosis, Tica, ticagrelor.

control strategy, irrespective of renal function.<sup>40</sup> While these findings must be interpreted in the context of an overall negative trial, exploratory analyses using eGFR as a continuous variable suggested a lower bleeding risk with ticagrelor monotherapy with decreasing eGFR, consistent with what reported in the present analysis. Compared to GLOBAL LEADERS, however, the TWILIGHT trial enrolled patients enriched with clinical and angiographic features of high risk for bleeding or ischaemia. Chronic kidney disease, an established risk factor for both these types of events, represented a clinical study entry criterion. Most of the available risk scores developed in PCI cohorts identify CKD as a qualifying condition to define patients as high bleeding risk.<sup>41–43</sup> Building on this prior evidence, our results support the relevance of CKD when evaluating the bleeding-related benefits of an aspirin withdrawal strategy. The absolute and relative bleeding risk reductions that we observed were larger than in other trials evaluating a similar treatment regimen and underscore the importance of implementing bleeding-avoidance strategies in such vulnerable cohorts.

It is also noteworthy that, in TWILIGHT, the reduction in bleeding risk was not counterbalanced by a trade-off in antithrombotic efficacy with ticagrelor monotherapy. Nonetheless, there was a numerical increase in ischaemic events among CKD patients on ticagrelor monotherapy, mainly driven by an excess in stroke rates. Such trend, although not statistically significant, was also suggested by visual assessment of the relationship between ischaemic event rates and eGFR used as a continuous variable. Whether aspirin serves an important platelet inhibitory role in the prothrombotic milieu of CKD, however, remains unproven. Hence, while these data seem to reassure on both the safety and efficacy of ticagrelor monotherapy after PCI among CKD patients, the limited sample size of our subgroup

analysis warrants prospective confirmation from adequately powered studies.

**Study limitations**

Randomization was not stratified by CKD status, and residual confounders between treatment groups may exist. Furthermore, type II error cannot be excluded in the context of an underpowered subgroup analysis. Hence, our findings must be considered hypothesis generating and dedicated prospective research is needed to assess the optimal treatment combination in high-risk CKD patients undergoing PCI. Moreover, the TWILIGHT trial excluded subjects with dialysis-dependent renal failure as well as those undergoing primary PCI for ST-segment elevation ACS. Therefore, the safety and efficacy of ticagrelor monotherapy observed in our study cannot be generalized to patient cohorts that would not otherwise be eligible for enrolment in the TWILIGHT trial.

**Conclusions**

Among high-risk patients undergoing PCI, a strategy of withdrawing aspirin and continuing ticagrelor alone after 3 months of DAPT significantly reduced clinically relevant as well as major bleeding without increasing ischaemic events as compared with ticagrelor plus aspirin, irrespective of renal function. However, owing to their worse risk profile, patients with CKD experienced a greater absolute risk reduction in bleeding events, but also a marginal increase in the rates of ischaemic events with ticagrelor monotherapy. Future research should explore the safety and efficacy of this treatment strategy across all kidney function categories.



## Supplementary material

Supplementary material is available at *European Heart Journal* online.

## Funding

This work was supported by the investigator-initiated grant from AstraZeneca.

**Conflict of interest:** G.G.S. reports consulting fees or honoraria from Abbott Vascular and Boston Scientific and that his institution has received a research grant from Boston Scientific. C.B. reports speaker honoraria from AstraZeneca and Boston Scientific. G.D. reports receiving consulting fees and advisory board fees from AstraZeneca, consulting fees from Biosensors, and previously holding stock in Medtronic. D.J.A. has received payment as an individual for: (i) consulting fee or honorarium from Abbott, Amgen, Aralez, AstraZeneca, Bayer, Biosensors, Boehringer Ingelheim, Bristol Myers Squibb, Chiesi, Daiichi Sankyo, Eli Lilly, Haemonetics, Janssen, Merck, PhaseBio, PLx Pharma, Pfizer, Sanofi, and The Medicines Company and (ii) participation in review activities from CeloNova and St. Jude Medical. Institutional payments for grants from Amgen, AstraZeneca, Bayer, Biosensors, CeloNova, CSL Behring, Daiichi Sankyo, Eisai, Eli Lilly, Gilead, Idorsia, Janssen, Matsutani Chemical Industry Co., Merck, Novartis, Osprey Medical, Renal Guard Solutions, and the Scott R. MacKenzie Foundation. S.M. has received research grants to the institution from AstraZeneca, Abbott, Boston Scientific, and Sanofi and has received honoraria for consultancy from AstraZeneca, Bayer, Biosensors, and Sanofi. D.J.C. reports receiving grant support, paid to his institution, and consulting fees from AstraZeneca, Medtronic, and Abbott Vascular, and grant support, paid to his institution, from Boston Scientific. J.E. reports receiving consulting fees and lecture fees from Abbott, Philips, Boston Scientific, and Medtronic, and lecture fees from Abiomed, Terumo, and Biosensors. C.M.G. reports receiving grant support and consulting fees from Angel Medical, Bayer, CSL Behring, Janssen Pharmaceuticals, Johnson & Johnson, and Portola Pharmaceuticals, consulting fees from the Medicines Company, Eli Lilly, Gilead Sciences, Novo Nordisk, WebMD, UpToDate Cardiovascular Medicine, Amarin Pharma, Amgen, Boehringer Ingelheim, Chiesi, Merck, PharmaMar, Sanofi, Somahlution, Verreson Corporation, Boston Scientific, Impact Bio, MedImmune, Medtelligence, MicroPort, PERT Consortium, and GE Healthcare, holding equity in nference, serving as chief executive officer of Baim Institute, and receiving grant support, paid to Baim Institute, from Bristol Myers Squibb. K.H. reports receiving lecture fees from AstraZeneca and Bayer. M.W.K. reports grants and/or personal fees from Abbott Vascular, Biosensors, Boston Scientific, CeloNova, Medtronic, OrbusNeich, and Terumo. V.K. has received personal fees/honoraria from Bayer, AstraZeneca, Abbott, Amgen, and Daiichi Sankyo. D.J.M. reports grants from AstraZeneca, during the conduct of the study. E.M.O. reports research grants from Abiomed and Chiesi, and consulting fees from AstraZeneca, Cara Therapeutics, Faculty Connection, Imbria, Impulse Medical, Janssen Pharmaceuticals, Milestone Pharmaceuticals, Xylocor, and Zoll Medical. K.G.O. reports receiving grant support and lecture fees from AstraZeneca and is employed by Biosensors. S.K.S. has received consulting fees or honoraria from Abbott, Boston Scientific, Abiomed, and Cardiovascular Systems, Inc. G.W. reports receiving grant support and advisory board fees from and holding equity in Corindus, advisory board fees from and holding equity in Filterlex, serving on an advisory board for and holding options in Trisol, and receiving grant support from Abbott, CSI, and RenalGuard. R.M. reports institutional research grants from Abbott, Abiomed, Applied Therapeutics, Arena, AstraZeneca, Bayer, Biosensors, Boston Scientific, Bristol Myers Squibb, CardiaWave, CellAegis, CERC, Chiesi, Concept Medical, CSL Behring, DSI, Insel Gruppe AG, Medtronic, Novartis Pharmaceuticals,

OrbusNeich, Philips, Transverse Medical, and Zoll; personal fees from ACC, Boston Scientific, California Institute for Regenerative Medicine (CIRM), Cine-Med Research, Janssen, WebMD, and SCAI; consulting fees paid to the institution from Abbott, Abiomed, AM-Pharma, Alleviant Medical, Bayer, Beth Israel Deaconess, CardiaWave, CeloNova, Chiesi, Concept Medical, DSI, Duke University, Idorsia Pharmaceuticals, Medtronic, Novartis, and Philips; Equity <1% in Applied Therapeutics, Elixir Medical, STEL, CONTROLRAD (spouse); and Scientific Advisory Board for AMA, Biosensors (spouse). All other authors declared no conflict of interest.

## Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

## References

- Matsushita K, Coresh J, Sang Y, Chalmers J, Fox C, Guallar E, Jafar T, Jassal SK, Landman GW, Muntner P, Roderick P, Sainruchi T, Schöttker B, Shankar A, Shlipak M, Tonelli M, Townsend J, van Zuilen A, Yamagishi K, Yamashita K, Gansevoort R, Sarnak M, Warnock DG, Woodward M, Ärnlöv J; CKD Prognosis Consortium. Estimated glomerular filtration rate and albuminuria for prediction of cardiovascular outcomes: a collaborative meta-analysis of individual participant data. *Lancet Diabetes Endocrinol* 2015;3:514–525.
- Bansal N, Katz R, Robinson-Cohen C, Odden MC, Dalrymple L, Shlipak MG, Sarnak MJ, Siscovick DS, Zelnick L, Psaty BM, Kestenbaum B, Correa A, Afkarian M, Young B, de Boer IH. Absolute rates of heart failure, coronary heart disease, and stroke in chronic kidney disease: an analysis of 3 community-based cohort studies. *JAMA Cardiol* 2017;2:314–318.
- Stefanini GG, Taniwaki M, Kalesan B, Räber L, Stortecky S, Pilgrim T, Onuma Y, Silber S, Serruys PW, Meier B, Jüni P, Windecker S. The impact of renal impairment on long-term safety and effectiveness of drug-eluting stents. *PLoS One* 2014;9:e106450.
- Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004;351:1296–1305.
- Molnar AO, Bota SE, Garg AX, Harel Z, Lam N, McArthur E, Nesrallah G, Perl J, Sood MM. The risk of major hemorrhage with CKD. *J Am Soc Nephrol* 2016;27:2825–2832.
- Ocak G, Rookmaaker MB, Algra A, de Borst GJ, Doevendans PA, Kappelle LJ, Verhaar MC, Visseren FL; the SMART Study Group. Chronic kidney disease and bleeding risk in patients at high cardiovascular risk: a cohort study. *J Thromb Haemost* 2018;16:65–73.
- Capodanno D, Angiolillo DJ. Antithrombotic therapy in patients with chronic kidney disease. *Circulation* 2012;125:2649–2661.
- Bonello L, Angiolillo DJ, Aradi D, Sibbing D. P2Y<sub>12</sub>-ADP receptor blockade in chronic kidney disease patients with acute coronary syndromes. *Circulation* 2018;138:1582–1596.
- Cao D, Chandiramani R, Chiarito M, Claessen BE, Mehran R. Evolution of antithrombotic therapy in patients undergoing percutaneous coronary intervention: a 40-year journey. *Eur Heart J* 2021;42:339–351.
- Genereux P, Giustino G, Witzensbichler B, Weisz G, Stuckey TD, Rinaldi MJ, Neumann FJ, Metzger DC, Henry TD, Cox DA, Duffy PL, Mazzaferri E, Yadav M, Francese DP, Palmerini T, Kirtane AJ, Litherland C, Mehran R, Stone GW. Incidence, predictors, and impact of post-discharge bleeding after percutaneous coronary intervention. *J Am Coll Cardiol* 2015;66:1036–1045.
- Mehran R, Pocock SJ, Stone GW, Clayton TC, Dangas GD, Feit F, Manoukian SV, Nikolsky E, Lansky AJ, Kirtane A, White HD, Colombo A, Ware JH, Moses JW, Ohman EM. Associations of major bleeding and myocardial infarction with the incidence and timing of mortality in patients presenting with non-ST-elevation acute coronary syndromes: a risk model from the ACUITY trial. *Eur Heart J* 2009;30:1457–1466.
- Valgimigli M, Costa F, Lokhnygina Y, Clare RM, Wallentin L, Moliterno DJ, Armstrong PW, White HD, Held C, Aylward PE, Van de Werf F, Harrington RA, Mahaffey KW, Tricoci P. Trade-off of myocardial infarction vs. bleeding types on mortality after acute coronary syndrome: lessons from the Thrombin Receptor Antagonist for Clinical Event Reduction in Acute Coronary Syndrome (TRACER) randomized trial. *Eur Heart J* 2017;38:804–810.
- Bhatt DL. Intensifying platelet inhibition—navigating between Scylla and Charybdis. *N Engl J Med* 2007;357:2078–2081.
- Mehran R, Baber U, Sharma SK, Cohen DJ, Angiolillo DJ, Briguori C, Cha JY, Collier T, Dangas G, Dudek D, Dzavik V, Escaned J, Gil R, Gurbel P, Hamm CW, Henry T, Huber K, Kastrati A, Kaul U, Kornowski R, Krucoff M, Kunadian V,

- Marx SO, Mehta SR, Moliterno D, Ohman EM, Oldroyd K, Sardella G, Sartori S, Shlofmitz R, Steg PG, Weisz G, Witzencbichler B, Han YL, Pocock S, Gibson CM. Ticagrelor with or without aspirin in high-risk patients after PCI. *N Engl J Med* 2019;**381**:2032–2042.
15. Baber U, Dangas G, Cohen DJ, Gibson CM, Mehta SR, Angiolillo DJ, Pocock SJ, Krucoff MW, Kastrati A, Ohman EM, Steg PG, Badimon J, Zafar MU, Chandrasekhar J, Sartori S, Aquino M, Mehran R. Ticagrelor with aspirin or alone in high-risk patients after coronary intervention: rationale and design of the TWILIGHT study. *Am Heart J* 2016;**182**:125–134.
  16. Mehran R, Rao SV, Bhatt DL, Gibson CM, Caixeta A, Eikelboom J, Kaul S, Wiviott SD, Menon V, Nikolsky E, Serebruany V, Valgimigli M, Vranckx P, Taggart D, Sabik JF, Cutlip DE, Krucoff MW, Ohman EM, Steg PG, White H. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation* 2011;**123**:2736–2747.
  17. Bovill EG, Terrin ML, Stump DC, Berke AD, Frederick M, Collen D, Feit F, Gore JM, Hillis LD, Lambrew CT, Leiboff R, Mann KG, Markis JE, Pratt CM, Sharkey SW, Sopko G, Tracy RP, Chesebro JH. Hemorrhagic events during therapy with recombinant tissue-type plasminogen activator, heparin, and aspirin for acute myocardial infarction. Results of the Thrombolysis in Myocardial Infarction (TIMI), phase II trial. *Ann Intern Med* 1991;**115**:256–265.
  18. GUSTO Investigators. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. *N Engl J Med* 1993;**329**:673–682.
  19. Kaatz S, Ahmad D, Spyropoulos AC, Schulman S; Subcommittee on Control of Anticoagulation. Definition of clinically relevant non-major bleeding in studies of anticoagulants in atrial fibrillation and venous thromboembolic disease in non-surgical patients: communication from the SSC of the ISTH. *J Thromb Haemost* 2015;**13**:2119–2126.
  20. Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD; Writing Group on the Joint ESC/ACCF/AHA/WHF Task Force for the Universal Definition of Myocardial Infarction, Thygesen K, Alpert JS, White HD, Jaffe AS, Katus HA, Apple FS, Lindahl B, Morrow DA, Chaitman BR, Clemmensen PM, Johanson P, Hod H, Underwood R, Bax JJ, Bonow JJ, Pinto F, Gibbons RJ, Fox KA, Atar D, Newby LK, Galvani M, Hamm CW, Uretsky BF, Steg PG, Wijns W, Bassand JP, Menasche P, Ravkilde J, Ohman EM, Antman EM, Wallentin LC, Armstrong PW, Simoons ML, Januzzi JL, Nieminen MS, Gheorghiade M, Filippatos G, Luepker RV, Fortmann SP, Rosamond WD, Levy D, Wood D, Smith SC, Hu D, Lopez-Sendon JL, Robertson RM, Weaver D, Tendera M, Bove AA, Parkhomenko AN, Vasilieva EJ, Mendis S; ESC Committee for Practice Guidelines (CPG). Third universal definition of myocardial infarction. *J Am Coll Cardiol* 2012;**60**:1581–1598.
  21. Cutlip DE, Windecker S, Mehran R, Boam A, Cohen DJ, van Es GA, Steg PG, Morel MA, Mauri L, Vranckx P, McFadden E, Lansky A, Hamon M, Krucoff MW, Serruys PW; Academic Research Consortium. Clinical end points in coronary stent trials: a case for standardized definitions. *Circulation* 2007;**115**:2344–2351.
  22. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J; for the CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;**150**:604–612.
  23. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002;**39**:S1–S266.
  24. Lutz J, Menke J, Sollinger D, Schinzel H, Thümel K. Haemostasis in chronic kidney disease. *Nephrol Dial Transplant* 2014;**29**:29–40.
  25. Baber U, Stone GW, Weisz G, Moreno P, Dangas G, Maehara A, Mintz GS, Cristea E, Fahy M, Xu K, Lansky AJ, Wennerblom B, Mathey DG, Templin B, Zhang Z, Serruys PW, Mehran R. Coronary plaque composition, morphology, and outcomes in patients with and without chronic kidney disease presenting with acute coronary syndromes. *JACC Cardiovasc Imaging* 2012;**5**:553–561.
  26. Daenen K, Andries A, Mekahli D, Van Schepdael A, Joret F, Bammens B. Oxidative stress in chronic kidney disease. *Pediatr Nephrol* 2019;**34**:975–991.
  27. Keltai M, Tonelli M, Mann JF, Sitkei E, Lewis BS, Hawken S, Mehta SR, Yusuf S; CURE Trial Investigators. Renal function and outcomes in acute coronary syndrome: impact of clopidogrel. *Eur J Cardiovasc Prev Rehabil* 2007;**14**:312–318.
  28. Best PJ, Steinhilber SR, Berger PB, Dasgupta A, Brennan DM, Szczec LA, Califf RM, Topol EJ. The efficacy and safety of short- and long-term dual antiplatelet therapy in patients with mild or moderate chronic kidney disease: results from the Clopidogrel for the Reduction of Events During Observation (CREDO) trial. *Am Heart J* 2008;**155**:687–693.
  29. Dasgupta A, Steinhilber SR, Bhatt DL, Berger PB, Shao M, Mak KH, Fox KA, Montalescot G, Weber MA, Haffner SM, Dimas AP, Steg PG, Topol EJ; CHARISMA Investigators. Clinical outcomes of patients with diabetic nephropathy randomized to clopidogrel plus aspirin versus aspirin alone (a post hoc analysis of the clopidogrel for high atherothrombotic risk and ischemic stabilization, management, and avoidance [CHARISMA] trial). *Am J Cardiol* 2009;**103**:1359–1363.
  30. Morel O, El Ghannudi S, Jesel L, Radulescu B, Meyer N, Wiesel ML, Caillard S, Campia U, Moulin B, Gachet C, Ohlmann P. Cardiovascular mortality in chronic kidney disease patients undergoing percutaneous coronary intervention is mainly related to impaired P2Y12 inhibition by clopidogrel. *J Am Coll Cardiol* 2011;**57**:399–408.
  31. Baber U, Mehran R, Kirtane AJ, Gurbel PA, Christodoulidis G, Maehara A, Witzencbichler B, Weisz G, Rinaldi MJ, Metzger DC, Henry TD, Cox DA, Duffy PL, Mazzaferri EL Jr, Xu K, Parise H, Brodie BR, Stuckey TD, Stone GW. Prevalence and impact of high platelet reactivity in chronic kidney disease: results from the Assessment of Dual Antiplatelet Therapy with Drug-Eluting Stents registry. *Circ Cardiovasc Interv* 2015;**8**:e001683.
  32. James S, Budaj A, Aylward P, Buck KK, Cannon CP, Cornel JH, Harrington RA, Horrow J, Katus H, Keltai M, Lewis BS, Parikh K, Storey RF, Szummer K, Wojdyla D, Wallentin L. Ticagrelor versus clopidogrel in acute coronary syndromes in relation to renal function: results from the Platelet Inhibition and Patient Outcomes (PLATO) trial. *Circulation* 2010;**122**:1056–1067.
  33. Chiarito M, Sanz-Sánchez J, Cannata F, Cao D, Sturla M, Panico C, Godino C, Regazzoli D, Reimers B, De Caterina R, Condorelli G, Ferrante G, Stefanini GG. Monotherapy with a P2Y12 inhibitor or aspirin for secondary prevention in patients with established atherosclerosis: a systematic review and meta-analysis. *Lancet* 2020;**395**:1487–1495.
  34. Armstrong PC, Leadbeater PD, Chan MV, Kirkby NS, Jakubowski JA, Mitchell JA, Warner TD. In the presence of strong P2Y12 receptor blockade, aspirin provides little additional inhibition of platelet aggregation. *J Thromb Haemost* 2011;**9**:552–561.
  35. Baber U, Zafar MU, Dangas G, Escolar G, Angiolillo DJ, Sharma SK, Kini AS, Sartori S, Joyce L, Vogel B, Farhan S, Gurbel P, Gibson CM, Fuster V, Mehran R, Badimon JJ. Ticagrelor with or without aspirin after PCI: the TWILIGHT platelet substudy. *J Am Coll Cardiol* 2020;**75**:578–586.
  36. Vranckx P, Valgimigli M, Juni P, Hamm C, Steg PG, Heg D, van Es GA, McFadden EP, Onuma Y, van Meijeren C, Chichareon P, Benit E, Mollmann H, Janssens L, Ferrario M, Moschovitis A, Zurakowski A, Dominici M, Van Geuns RJ, Huber K, Slagboom T, Serruys PW, Windecker S; GLOBAL LEADERS Investigators. Ticagrelor plus aspirin for 1 month, followed by ticagrelor monotherapy for 23 months vs aspirin plus clopidogrel or ticagrelor for 12 months, followed by aspirin monotherapy for 12 months after implantation of a drug-eluting stent: a multicentre, open-label, randomised superiority trial. *Lancet* 2018;**392**:940–949.
  37. Hahn JY, Song YB, Oh JH, Chun WJ, Park YH, Jang WJ, Im ES, Jeong JO, Cho BR, Oh SK, Yun KH, Cho DK, Lee JY, Koh YY, Bae JW, Choi JW, Lee WS, Yoon HJ, Lee SU, Cho JH, Choi WG, Rha SW, Lee JM, Park TK, Yang JH, Choi JH, Choi SH, Lee SH, Gwon HC; SMART-CHOICE Investigators. Effect of P2Y12 inhibitor monotherapy vs dual antiplatelet therapy on cardiovascular events in patients undergoing percutaneous coronary intervention: the SMART-CHOICE randomized clinical trial. *JAMA* 2019;**321**:2428–2437.
  38. Watanabe H, Domei T, Morimoto T, Natsuaki M, Shiomi H, Toyota T, Ohya M, Suwa S, Takagi K, Nanasato M, Hata Y, Yagi M, Suematsu N, Yokomatsu T, Takamisawa I, Doi M, Noda T, Okayama H, Seino Y, Tada T, Sakamoto H, Hibi K, Abe M, Kawai K, Nakao K, Ando K, Tanabe K, Ikari Y, Hanaoka KI, Morino Y, Kozuma K, Kadota K, Furukawa Y, Nakagawa Y, Kimura T; STOPDAPT-2 Investigators. Effect of 1-month dual antiplatelet therapy followed by clopidogrel vs 12-month dual antiplatelet therapy on cardiovascular and bleeding events in patients receiving PCI: the STOPDAPT-2 randomized clinical trial. *JAMA* 2019;**321**:2414–2427.
  39. Kim BK, Hong SJ, Cho YH, Yun KH, Kim YH, Suh Y, Cho JY, Her AY, Cho S, Jeon DW, Yoo SY, Cho DK, Hong BK, Kwon H, Ahn CM, Shin DH, Nam CM, Kim JS, Ko YG, Choi D, Hong MK, Jang Y; TICO Investigators. Effect of ticagrelor monotherapy vs ticagrelor with aspirin on major bleeding and cardiovascular events in patients with acute coronary syndrome: the TICO randomized clinical trial. *JAMA* 2020;**323**:2407–2416.
  40. Tomaniak M, Chichareon P, Klimczak-Tomaniak D, Takahashi K, Kogame N, Modolo R, Wang R, Ono M, Hara H, Gao C, Kawashima H, Rademaker-Havinga T, Garg S, Curzen N, Haude M, Kochman J, Gori T, Montalescot G, Angiolillo DJ, Capodanno D, Storey RF, Hamm C, Vranckx P, Valgimigli M, Windecker S, Onuma Y, Serruys PW, Anderson R. Impact of renal function on clinical outcomes after PCI in ACS and stable CAD patients treated with ticagrelor: a pre-specified analysis of the GLOBAL LEADERS randomized clinical trial. *Clin Res Cardiol* 2020;**109**:930–943.
  41. Baber U, Mehran R, Giustino G, Cohen DJ, Henry TD, Sartori S, Ariti C, Litherland C, Dangas G, Gibson CM, Krucoff MW, Moliterno DJ, Kirtane AJ, Stone GW, Colombo A, Chieffo A, Kini AS, Witzencbichler B, Weisz G, Steg PG, Pocock S. Coronary thrombosis and major bleeding after pci with drug-eluting stents: risk scores from PARIS. *J Am Coll Cardiol* 2016;**67**:2224–2234.

42. Costa F, van Klaveren D, James S, Heg D, Raber L, Feres F, Pilgrim T, Hong MK, Kim HS, Colombo A, Steg PG, Zanchin T, Palmerini T, Wallentin L, Bhatt DL, Stone GW, Windecker S, Steyerberg EW, Valgimigli M; PRECISE-DAPT Study Investigators. Derivation and validation of the predicting bleeding complications in patients undergoing stent implantation and subsequent dual antiplatelet therapy (PRECISE-DAPT) score: a pooled analysis of individual-patient datasets from clinical trials. *Lancet* 2017;**389**: 1025–1034.
43. Urban P, Mehran R, Collieran R, Angiolillo DJ, Byrne RA, Capodanno D, Cuisset T, Cutlip D, Eerdmans P, Eikelboom J, Farb A, Gibson CM, Gregson J, Haude M, James SK, Kim HS, Kimura T, Konishi A, Laschinger J, Leon MB, Magee PFA, Mitsutake Y, Mylotte D, Pocock S, Price MJ, Rao SV, Spitzer E, Stockbridge N, Valgimigli M, Varenne O, Windhoevel U, Yeh RW, Krucoff MW, Morice MC. Defining high bleeding risk in patients undergoing percutaneous coronary intervention: a consensus document from the academic research consortium for high bleeding risk. *Circulation* 2019;**40**:2632–2653.