

Ideal Alternatives


- Function (Model): $f(x)$
 $x \rightarrow \begin{cases} g(x) > 0 & \text{Output = class 1} \\ \text{else} & \text{Output = class 2} \end{cases}$
- Loss function:

$$L(f) = \sum_n \delta(f(x^n) \neq \hat{y}^n)$$

The number of times f get incorrect results on training data.
- Find the best function:
 - Example: Perceptron, SVM Not Today

Two Classes

Estimating the Probabilities
From training data




Given an x , which class does it belong to

$$P(C_1|x) = \frac{P(x|C_1)P(C_1)}{P(x|C_1)P(C_1) + P(x|C_2)P(C_2)}$$

Generative Model $P(x) = P(x|C_1)P(C_1) + P(x|C_2)P(C_2)$

Maximum Likelihood $f_{\mu,\Sigma}(x) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\Sigma|^{1/2}} \exp\left\{-\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu)\right\}$



The Gaussian with any mean μ and covariance matrix Σ can generate these points. → Different Likelihood

Likelihood of a Gaussian with mean μ and covariance matrix Σ
 = the probability of the Gaussian samples $x^1, x^2, x^3, \dots, x^{79}$

$$L(\mu, \Sigma) = f_{\mu,\Sigma}(x^1) f_{\mu,\Sigma}(x^2) f_{\mu,\Sigma}(x^3) \dots f_{\mu,\Sigma}(x^{79})$$

Maximum Likelihood

We have the “Water” type Pokémons: $x^1, x^2, x^3, \dots, x^{79}$

We assume $x^1, x^2, x^3, \dots, x^{79}$ generate from the Gaussian (μ^*, Σ^*) with the **maximum likelihood**

$$L(\mu, \Sigma) = f_{\mu,\Sigma}(x^1) f_{\mu,\Sigma}(x^2) f_{\mu,\Sigma}(x^3) \dots f_{\mu,\Sigma}(x^{79})$$

$$f_{\mu,\Sigma}(x) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\Sigma|^{1/2}} \exp\left\{-\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu)\right\}$$

$$\mu^*, \Sigma^* = \arg \max_{\mu, \Sigma} L(\mu, \Sigma)$$

$$\mu^* = \frac{1}{79} \sum_{n=1}^{79} x^n \quad \Sigma^* = \frac{1}{79} \sum_{n=1}^{79} (x^n - \mu^*) (x^n - \mu^*)^T$$

average

Now we can do classification 😊

$$f_{\mu^1, \Sigma^1}(x) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\Sigma^1|^{1/2}} \exp\left\{-\frac{1}{2}(x - \mu^1)^T (\Sigma^1)^{-1} (x - \mu^1)\right\}$$

$$\mu^1 = \begin{bmatrix} 75.0 \\ 71.3 \end{bmatrix} \quad \Sigma^1 = \begin{bmatrix} 874 & 327 \\ 327 & 929 \end{bmatrix}$$

$$P(C_1) = 79 / (79 + 61) = 0.56$$

$$P(C_1|x) = \frac{P(x|C_1)P(C_1)}{P(x|C_1)P(C_1) + P(x|C_2)P(C_2)}$$

$$f_{\mu^2, \Sigma^2}(x) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\Sigma^2|^{1/2}} \exp\left\{-\frac{1}{2}(x - \mu^2)^T (\Sigma^2)^{-1} (x - \mu^2)\right\}$$

$$\mu^2 = \begin{bmatrix} 55.6 \\ 59.8 \end{bmatrix} \quad \Sigma^2 = \begin{bmatrix} 847 & 422 \\ 422 & 685 \end{bmatrix}$$

$$P(C_2) = 61 / (79 + 61) = 0.44$$

If $P(C_1|x) > 0.5$ ➡ x belongs to class 1 (Water)