Sensitivity Analysis: Flood Factor Groupings

Alvin Sheng

Contents

Helper Functions	. 2
CAR model results, Coronary Heart Disease Stratified on RPL_THEMES Credible Interval plots for the coefficients, in ggplot	. 4
CAR model results, High Blood Pressure Stratified on RPL_THEMES Credible Interval plots for the coefficients, in ggplot	. 7
CAR model results, Asthma Stratified on RPL_THEMES Credible Interval plots for the coefficients, in ggplot	. 10
CAR model results, Poor Mental Health Stratified on RPL_THEMES Credible Interval plots for the coefficients, in ggplot	. 13

```
library(here)
## Warning in readLines(f, n): line 1 appears to contain an embedded nul
## Warning in readLines(f, n): incomplete final line found on '/Volumes/
## ALVINDRIVE2/flood-risk-health-effects/._flood-risk-health-effects.Rproj'
## here() starts at /Volumes/ALVINDRIVE2/flood-risk-health-effects
library(coda)
library(CARBayes)
## Loading required package: MASS
## Loading required package: Rcpp
## Registered S3 method overwritten by 'GGally':
    method from
##
    +.gg
           ggplot2
library(ggplot2)
library(tidyverse)
## -- Attaching packages -----
                                                   ----- tidyverse 1.3.1 --
## v tibble 3.1.8
                       v dplyr 1.0.10
## v tidyr 1.2.1
                       v stringr 1.4.0
           2.1.1
                       v forcats 0.5.1
## v readr
## v purrr
           0.3.4
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()
                  masks stats::lag()
## x dplyr::select() masks MASS::select()
fhs_model_df <- readRDS(here("intermediary_data/sensitivity_analysis/fhs_model_df_ff_grouped.rds"))</pre>
var_names <- c("Intercept", "flood_ff_group1", "flood_ff_group2", "flood_ff_group3",</pre>
              "EP_UNINSUR", "pollute_conc_pc1", "pollute_conc_pc2",
              "pollute_conc_pc3", "tmmx", "rmax", "Data_Value_CSMOKING")
names_ff_grouped_strat <- c(paste("strat0", var_names, sep = ":"),</pre>
                         paste("strat1", var_names, sep = ":"))
```

Helper Functions

Function for post-processing the inference

```
# extract the flood risk PC coefficients
# pc_idx is the vector of indices of the flood risk PC coefficients, after splitting data frame by stra
beta_data_frames_extract <- function(beta_inference_df, pc_idx) {

beta_inference_df_strat0 <- beta_inference_df[1:(nrow(beta_inference_df)/2),]

beta_inference_df_strat1 <- beta_inference_df[(nrow(beta_inference_df)/2 + 1):nrow(beta_inference_df)

beta_pcs_strat0 <- beta_inference_df_strat0[pc_idx, ]</pre>
```

```
beta_pcs_strat1 <- beta_inference_df_strat1[pc_idx, ]

beta_pcs_strat0 <- mutate(beta_pcs_strat0, var_idx = factor(1:nrow(beta_pcs_strat0)))
beta_pcs_strat1 <- mutate(beta_pcs_strat1, var_idx = factor(1:nrow(beta_pcs_strat1)))

return(list(beta_pcs_strat0 = beta_pcs_strat0, beta_pcs_strat1 = beta_pcs_strat1))
}</pre>
```

CAR model results, Coronary Heart Disease Stratified on RPL_THEMES

Inference is based on 3 markov chains, each of which has been run for 110000 samples, the first 10000 of which has been removed for burn-in. The remaining 100000 samples are thinned by 2, resulting in 150000 samples for inference across the 3 Markov chains.

```
load(here("modeling_files/sensitivity_analysis/ff_grouped/all_census_tract_CHD.RData"))
beta_samples_matrix <- rbind(chain1$samples$beta, chain2$samples$beta, chain3$samples$beta)
colnames(beta_samples_matrix) <- names_ff_grouped_strat
beta_inference <- round(t(apply(beta_samples_matrix, 2, quantile, c(0.5, 0.025, 0.975))),5)
beta_inference</pre>
```

```
##
                                 50%
                                               97.5%
                                        2.5%
## strat0:Intercept
                             6.23243 6.21298 6.25191
## strat0:flood_ff_group1
                             0.01407 -0.00782 0.03585
## strat0:flood_ff_group2
                             0.01823 -0.00077 0.03706
## strat0:flood_ff_group3
                             0.08804 0.07019 0.10586
## strat0:EP UNINSUR
                            -0.01044 -0.03871 0.01806
## strat0:pollute conc pc1
                            -0.44536 -0.50037 -0.38884
## strat0:pollute_conc_pc2
                            -0.50906 -0.58969 -0.42809
## strat0:pollute_conc_pc3
                            -0.20659 -0.28198 -0.12932
## strat0:tmmx
                            -0.00061 -0.12178 0.12021
## strat0:rmax
                             0.14099 0.04885 0.23051
## strat1:Intercept
                             6.86954 6.85271 6.88619
## strat1:flood_ff_group1
                             0.00488 -0.01267 0.02231
## strat1:flood_ff_group2
                             0.01922 0.00378 0.03465
## strat1:flood_ff_group3
                             0.06069 0.04108 0.08031
## strat1:EP_UNINSUR
                            -0.13900 -0.15783 -0.12044
## strat1:pollute_conc_pc1
                            -0.28413 -0.34123 -0.22545
## strat1:pollute conc pc2
                            -0.40869 -0.48847 -0.32956
## strat1:pollute_conc_pc3
                            -0.27477 -0.35068 -0.19704
## strat1:tmmx
                             0.08883 -0.03322 0.21065
## strat1:rmax
                             0.13936 0.04691 0.22889
## strat1:Data Value CSMOKING 0.85763 0.83422 0.88101
```


```
saveRDS(beta_inference, file = here("modeling_files/sensitivity_analysis/ff_grouped/beta_inference_file
List of significant beta coefficients:
row.names(beta_inference)[sign(beta_inference[, 2]) == sign(beta_inference[, 3])]
  [1] "strat0:Intercept"
                                      "strat0:flood_ff_group3"
## [3] "strat0:pollute_conc_pc1"
                                      "strat0:pollute_conc_pc2"
## [5] "strat0:pollute_conc_pc3"
                                      "strat0:rmax"
## [7] "strat0:Data_Value_CSMOKING" "strat1:Intercept"
                                      "strat1:flood_ff_group3"
## [9] "strat1:flood_ff_group2"
## [11] "strat1:EP_UNINSUR"
                                      "strat1:pollute_conc_pc1"
## [13] "strat1:pollute_conc_pc2"
                                      "strat1:pollute_conc_pc3"
## [15] "strat1:rmax"
                                      "strat1:Data_Value_CSMOKING"
Credible Interval plots for the coefficients, in ggplot
# first, process the beta_inference matrix in a form ggplot can understand
beta_inference_df <- as.data.frame(beta_inference)</pre>
beta_inference_df <- mutate(beta_inference_df, var_name = row.names(beta_inference_df))
beta_inference_df <- rename(beta_inference_df,</pre>
                            post_median = `50%`,
                            post_2.5 = 2.5\%,
                            post_97.5 = `97.5\%`)
beta_inference_df$var_name <- substring(beta_inference_df$var_name, first = 8)</pre>
beta_inference_df$var_name <- factor(beta_inference_df$var_name,</pre>
                                      levels = unique(beta_inference_df$var_name))
beta_inference_df$strat <- factor(c(rep("Low SV", (nrow(beta_inference_df)/2)),
                                     rep("High SV", (nrow(beta_inference_df)/2))), levels = c("Low SV",
Splitting up the beta coefficients for each strata
beta_inference_df_strat0 <- beta_inference_df[1:(nrow(beta_inference_df)/2),]
beta_inference_df_strat1 <- beta_inference_df[(nrow(beta_inference_df)/2 + 1):nrow(beta_inference_df),]
Note: The intercept for both strata is not included.
p <- ggplot(beta_inference_df_strat0[-1, ], aes(x = var_name, y = post_median, color = strat)) +
  geom_point() +
  ylim(c(-1, 2)) +
  theme(axis.text.x = element_text(angle = 45, vjust = 1, hjust=1), axis.title.x = element_blank(), axi
        axis.text=element_text(size=12),
        plot.margin = margin(5.5, 5.5, 5.5, 25)) +
  geom_errorbar(aes(ymin = post_2.5, ymax = post_97.5, width = 0.4), col = "#F8766D") +
  geom_vline(xintercept = c(3.5, 4.5, 7.5, 9.5), col = "blue") +
  geom_hline(yintercept = 0, col = "red") +
  annotate(geom = "text", x = 2, y = 1.45, label = "Flood\nRisk",
           col = "blue", size = 4.5) +
  scale_x_discrete(labels = c("FF Group 1", "FF Group 2", "FF Group 3", "Uninsured",
                               "Pollution PC 1", "Pollution PC 2", "Pollution PC 3",
                              "Maximum Temperature", "Maximum Humidity",
```

geom_errorbar(data = beta_inference_df_strat1[-1,], aes(ymin = post_2.5, ymax = post_97.5, width = 0


geom_point(data = beta_inference_df_strat1[-1,], col = "#00BFC4") + # strat 1

"Smoking")) + ggtitle("95% Credible Intervals, CHD, Stratified on All RPL


95% Credible Intervals, CHD, Stratified on All RPL Themes, Flood Factor (



```
ggsave(here("figures/final_figures/sensitivity_analysis/ff_grouped/CHD_CI_rpls.pdf"),
       plot = p, device = "pdf",
       width = 8, height = 6, units = "in")
pc_extract_idx <- 2:4</pre>
beta_CHD_pcs <- beta_data_frames_extract(beta_inference_df, pc_extract_idx)</pre>
p <- ggplot(beta_CHD_pcs$beta_pcs_strat0, aes(x = var_idx, y = post_median, color = strat)) +
  geom_point() +
  ylim(c(-0.04, 0.13)) +
  theme(axis.text.x = element_text(angle = 45, vjust = 1, hjust=1), axis.title.x = element_blank(), axi
        axis.text=element_text(size=12),
        plot.margin = margin(5.5, 5.5, 5.5, 25)) +
  geom_errorbar(aes(ymin = post_2.5, ymax = post_97.5, width = 0.4), col = "#F8766D") +
  geom_hline(yintercept = 0, col = "red") +
  scale_x_discrete(labels = c("FF Group 1", "FF Group 2", "FF Group 3"), 6) + ggtitle("95% Credible Int
  geom_point(data = beta_CHD_pcs$beta_pcs_strat1, col = "#00BFC4") + # strat 1
  geom_errorbar(data = beta_CHD_pcs$beta_pcs_strat1, aes(ymin = post_2.5, ymax = post_97.5, width = 0.4
  scale_color_manual(name = "Strata",
                     values = c("#F8766D", "#00BFC4"),
```



95% Credible Intervals for Flood Risk Groupings, Coronary Heart Diseas



```
ggsave(here("figures/final_figures/sensitivity_analysis/ff_grouped/CHD_cred_intervals_fr_only.pdf"),
    plot = p, device = "pdf",
    width = 8, height = 6, units = "in")
```

CAR model results, High Blood Pressure Stratified on RPL_THEMES

```
load(here("modeling_files/sensitivity_analysis/ff_grouped/all_census_tract_BPHIGH.RData"))
beta_samples_matrix <- rbind(chain1$samples$beta, chain2$samples$beta, chain3$samples$beta)
colnames(beta_samples_matrix) <- names_ff_grouped_strat</pre>
beta_inference <- round(t(apply(beta_samples_matrix, 2, quantile, c(0.5, 0.025, 0.975))),5)
beta_inference
##
                                   50%
                                           2.5%
                                                   97.5%
## strat0:Intercept
                              31.05190 30.99444 31.10923
## strat0:flood_ff_group1
                              -0.01803 -0.08435 0.04823
## strat0:flood_ff_group2
                               0.01329 -0.04422 0.07048
## strat0:flood_ff_group3
                               0.20867 0.15438 0.26284
## strat0:EP_UNINSUR
                              -0.09963 -0.18484 -0.01406
```

```
## strat0:pollute_conc_pc1
                            -1.66675 -1.83825 -1.49119
## strat0:pollute_conc_pc2
                            -2.39616 -2.64341 -2.14154
## strat0:pollute_conc_pc3
                             0.37656 0.14324 0.62196
## strat0:tmmx
                             -0.03531 -0.40350 0.34643
                             0.19324 -0.10026 0.47757
## strat0:rmax
## strat0:Data_Value_CSMOKING 0.73863 0.64527 0.83213
## strat1:Intercept
                    32.84111 32.79177 32.88981
## strat1:EP_UNINSUR
                             -0.47514 -0.53158 -0.41884
## strat1:pollute_conc_pc1
                             -0.74064 -0.91870 -0.55859
## strat1:pollute_conc_pc2
                             -2.22077 -2.46508 -1.97155
## strat1:pollute_conc_pc3
                             -0.17637 -0.41242 0.07120
## strat1:tmmx
                              0.49442 0.12109 0.88122
## strat1:rmax
                              0.28215 -0.01326 0.56640
## strat1:Data_Value_CSMOKING 2.12807 2.05691 2.19935
saveRDS(beta_inference, file = here("modeling_files/sensitivity_analysis/ff_grouped/beta_inference_file
List of significant beta coefficients:
row.names(beta_inference)[sign(beta_inference[, 2]) == sign(beta_inference[, 3])]
## [1] "strat0:Intercept"
                                    "strat0:flood_ff_group3"
## [3] "strat0:EP_UNINSUR"
                                   "strat0:pollute_conc_pc1"
## [5] "strat0:pollute_conc_pc2"
                                   "strat0:pollute_conc_pc3"
## [7] "strat0:Data_Value_CSMOKING"
                                   "strat1:Intercept"
## [9] "strat1:EP_UNINSUR"
                                    "strat1:pollute_conc_pc1"
                                   "strat1:tmmx"
## [11] "strat1:pollute_conc_pc2"
## [13] "strat1:Data_Value_CSMOKING"
Credible Interval plots for the coefficients, in ggplot
# first, process the beta_inference matrix in a form ggplot can understand
beta_inference_df <- as.data.frame(beta_inference)</pre>
beta_inference_df <- mutate(beta_inference_df, var_name = row.names(beta_inference_df))
beta_inference_df <- rename(beta_inference_df,</pre>
```


```
post_median = `50%`,
                              post_2.5 = 2.5\%,
                              post_97.5 = `97.5\%`)
beta_inference_df$var_name <- substring(beta_inference_df$var_name, first = 8)</pre>
beta_inference_df$var_name <- factor(beta_inference_df$var_name,</pre>
                                        levels = unique(beta_inference_df$var_name))
beta_inference_df$strat <- factor(c(rep("Low SV", (nrow(beta_inference_df)/2)),</pre>
                                      rep("High SV", (nrow(beta_inference_df)/2))), levels = c("Low SV",
Splitting up the beta coefficients for each strata
```

```
beta_inference_df_strat0 <- beta_inference_df[1:(nrow(beta_inference_df)/2),]
beta_inference_df_strat1 <- beta_inference_df[(nrow(beta_inference_df)/2 + 1):nrow(beta_inference_df),]
```

Note: The intercept for both strata is not included.

```
p <- ggplot(beta_inference_df_strat0[-1, ], aes(x = var_name, y = post_median, color = strat)) +
  geom_point() +
  ylim(c(-3, 5)) +
  theme(axis.text.x = element_text(angle = 45, vjust = 1, hjust=1), axis.title.x = element_blank(), axi
        axis.text=element_text(size=12),
        plot.margin = margin(5.5, 5.5, 5.5, 25)) +
  geom_errorbar(aes(ymin = post_2.5, ymax = post_97.5, width = 0.4), col = "#F8766D") +
  geom_vline(xintercept = c(3.5, 4.5, 7.5, 9.5), col = "blue") +
  geom_hline(yintercept = 0, col = "red") +
  annotate(geom = "text", x = 2, y = 3.95, label = "Flood\nRisk",
           col = "blue", size = 4.5) +
  scale_x_discrete(labels = c("FF Group 1", "FF Group 2", "FF Group 3", "Uninsured",
                              "Pollution PC 1", "Pollution PC 2", "Pollution PC 3",
                              "Maximum Temperature", "Maximum Humidity",
                              "Smoking")) + ggtitle("95% Credible Intervals, BPHIGH, Stratified on All
  geom_point(data = beta_inference_df_strat1[-1, ], col = "#00BFC4") + # strat 1
  geom_errorbar(data = beta_inference_df_strat1[-1, ], aes(ymin = post_2.5, ymax = post_97.5, width = 0
  scale_color_manual(name = "Strata",
                     values = c("#F8766D", "#00BFC4"),
                     drop = FALSE)
р
```


95% Credible Intervals, BPHIGH, Stratified on All RPL Themes, Flood Fac



```
ggsave(here("figures/final_figures/sensitivity_analysis/ff_grouped/BPHIGH_CI_rpls.pdf"),
    plot = p, device = "pdf",
    width = 8, height = 6, units = "in")
```

```
pc_extract_idx <- 2:4</pre>
beta BPHIGH pcs <- beta data frames extract(beta inference df, pc extract idx)
p <- ggplot(beta_BPHIGH_pcs$beta_pcs_strat0, aes(x = var_idx, y = post_median, color = strat)) +
  geom_point() +
  ylim(c(-0.23, 0.28)) +
  theme(axis.text.x = element_text(angle = 45, vjust = 1, hjust=1), axis.title.x = element_blank(), axi
        axis.text=element_text(size=12),
        plot.margin = margin(5.5, 5.5, 5.5, 25)) +
  geom_errorbar(aes(ymin = post_2.5, ymax = post_97.5, width = 0.4), col = "#F8766D") +
  geom_hline(yintercept = 0, col = "red") +
  scale_x_discrete(labels = c("FF Group 1", "FF Group 2", "FF Group 3"), 6) + ggtitle("95% Credible Int
  geom_point(data = beta_BPHIGH_pcs$beta_pcs_strat1, col = "#00BFC4") + # strat 1
  geom_errorbar(data = beta_BPHIGH_pcs$beta_pcs_strat1, aes(ymin = post_2.5, ymax = post_97.5, width = 0
  scale_color_manual(name = "Strata",
                     values = c("#F8766D", "#00BFC4"),
                     drop = FALSE)
```

95% Credible Intervals for Flood Risk Groupings, High Blood Pressure



ggsave(here("figures/final_figures/sensitivity_analysis/ff_grouped/BPHIGH_cred_intervals_fr_only.pdf"),
 plot = p, device = "pdf",
 width = 8, height = 6, units = "in")

CAR model results, Asthma Stratified on RPL THEMES


```
load(here("modeling_files/sensitivity_analysis/ff_grouped/all_census_tract_CASTHMA.RData"))
beta_samples_matrix <- rbind(chain1$samples$beta, chain2$samples$beta, chain3$samples$beta)
colnames(beta_samples_matrix) <- names_ff_grouped_strat</pre>
beta_inference <- round(t(apply(beta_samples_matrix, 2, quantile, c(0.5, 0.025, 0.975))),5)
beta_inference
                                          2.5%
##
                                  50%
                                                  97.5%
## strat0:Intercept
                              9.76095 9.75423 9.76770
## strat0:flood_ff_group1
                             -0.02519 -0.03306 -0.01736
## strat0:flood_ff_group2
                              0.00145 -0.00539 0.00825
## strat0:flood_ff_group3
                             -0.00239 -0.00883 0.00405
## strat0:EP_UNINSUR
                             -0.02874 -0.03881 -0.01864
## strat0:pollute_conc_pc1
                              0.12192 0.10144 0.14282
## strat0:pollute_conc_pc2
                             -0.17891 -0.20813 -0.14853
## strat0:pollute_conc_pc3
                             -0.04918 -0.07685 -0.01945
## strat0:tmmx
                             -0.00275 -0.04600 0.04327
## strat0:rmax
                             -0.08532 -0.12032 -0.05156
## strat0:Data_Value_CSMOKING 1.12509 1.11403 1.13615
## strat1:Intercept 9.94379 9.93804 9.94947
## strat1:flood_ff_group1
## strat1:flood_ff_group2
## strat1:flood_ff_group3
## strat1:flood_ff_group1
                             -0.00499 -0.01134 0.00132
                              0.00200 -0.00354 0.00756
                             -0.02300 -0.03007 -0.01599
## strat1:EP_UNINSUR
                             -0.08558 -0.09225 -0.07890
## strat1:pollute_conc_pc2
                             -0.16418 -0.19330 -0.13449
## strat1:pollute_conc_pc3
                             -0.10285 -0.13069 -0.07291
## strat1:tmmx
                              0.04393 0.00036 0.09082
## strat1:rmax
                             -0.12043 -0.15573 -0.08667
## strat1:Data_Value_CSMOKING 1.31047 1.30207 1.31894
saveRDS(beta_inference, file = here("modeling_files/sensitivity_analysis/ff_grouped/beta_inference_file
List of significant beta coefficients:
row.names(beta_inference)[sign(beta_inference[, 2]) == sign(beta_inference[, 3])]
  [1] "strat0:Intercept"
                                    "strat0:flood_ff_group1"
##
   [3] "strat0:EP_UNINSUR"
                                    "strat0:pollute_conc_pc1"
## [5] "strat0:pollute_conc_pc2"
                                    "strat0:pollute_conc_pc3"
## [7] "strat0:rmax"
                                    "strat0:Data_Value_CSMOKING"
   [9] "strat1:Intercept"
                                    "strat1:flood_ff_group3"
## [11] "strat1:EP_UNINSUR"
                                    "strat1:pollute_conc_pc1"
                                    "strat1:pollute_conc_pc3"
## [13] "strat1:pollute_conc_pc2"
## [15] "strat1:tmmx"
                                    "strat1:rmax"
```

[17] "strat1:Data_Value_CSMOKING"


Credible Interval plots for the coefficients, in ggplot


```
# first, process the beta_inference matrix in a form applot can understand
beta_inference_df <- as.data.frame(beta_inference)</pre>
beta_inference_df <- mutate(beta_inference_df, var_name = row.names(beta_inference_df))</pre>
beta_inference_df <- rename(beta_inference_df,</pre>
                             post_median = `50%`,
                             post_2.5 = 2.5\%,
                             post_97.5 = `97.5\%`)
beta_inference_df$var_name <- substring(beta_inference_df$var_name, first = 8)</pre>
beta_inference_df$var_name <- factor(beta_inference_df$var_name,</pre>
                                      levels = unique(beta_inference_df$var_name))
beta_inference_df$strat <- factor(c(rep("Low SV", (nrow(beta_inference_df)/2)),</pre>
                                     rep("High SV", (nrow(beta_inference_df)/2))), levels = c("Low SV",
Splitting up the beta coefficients for each strata
beta inference df strat0 <- beta inference df[1:(nrow(beta inference df)/2),]
beta_inference_df_strat1 <- beta_inference_df[(nrow(beta_inference_df)/2 + 1):nrow(beta_inference_df),]
Note: The intercept for both strata is not included.
p <- ggplot(beta_inference_df_strat0[-1, ], aes(x = var_name, y = post_median, color = strat)) +
  geom_point() +
  ylim(c(-1, 1.5)) +
  theme(axis.text.x = element_text(angle = 45, vjust = 1, hjust=1), axis.title.x = element_blank(), axi
        axis.text=element text(size=12),
        plot.margin = margin(5.5, 5.5, 5.5, 25)) +
  geom_errorbar(aes(ymin = post_2.5, ymax = post_97.5, width = 0.4), col = "#F8766D") +
  geom_vline(xintercept = c(3.5, 4.5, 7.5, 9.5), col = "blue") +
  geom_hline(yintercept = 0, col = "red") +
  annotate(geom = "text", x = 2, y = 0.95, label = "Flood\nRisk",
           col = "blue", size = 4.5) +
  scale_x_discrete(labels = c("FF Group 1", "FF Group 2", "FF Group 3", "Uninsured",
                               "Pollution PC 1", "Pollution PC 2", "Pollution PC 3",
                               "Maximum Temperature", "Maximum Humidity",
                               "Smoking")) + ggtitle("95% Credible Intervals, CASTHMA, Stratified on All
  geom_point(data = beta_inference_df_strat1[-1, ], col = "#00BFC4") + # strat 1
  geom_errorbar(data = beta_inference_df_strat1[-1, ], aes(ymin = post_2.5, ymax = post_97.5, width = 0
  scale_color_manual(name = "Strata",
                     values = c("#F8766D", "#00BFC4"),
                     drop = FALSE)
```

95% Credible Intervals, CASTHMA, Stratified on All RPL Themes, Flood



```
ggsave(here("figures/final_figures/sensitivity_analysis/ff_grouped/CASTHMA_CI_rpls.pdf"),
      plot = p, device = "pdf",
       width = 8, height = 6, units = "in")
pc_extract_idx <- 2:4</pre>
beta CASTHMA pcs <- beta data frames extract(beta inference df, pc extract idx)
p <- ggplot(beta_CASTHMA_pcs$beta_pcs_strat0, aes(x = var_idx, y = post_median, color = strat)) +
  geom_point() +
  ylim(c(-0.035, 0.0285)) +
  theme(axis.text.x = element_text(angle = 45, vjust = 1, hjust=1), axis.title.x = element_blank(), axi
        axis.text=element_text(size=12),
        plot.margin = margin(5.5, 5.5, 5.5, 25)) +
  geom_errorbar(aes(ymin = post_2.5, ymax = post_97.5, width = 0.4), col = "#F8766D") +
  geom_hline(yintercept = 0, col = "red") +
  scale_x_discrete(labels = c("FF Group 1", "FF Group 2", "FF Group 3"), 6) + ggtitle("95% Credible Int
  geom_point(data = beta_CASTHMA_pcs$beta_pcs_strat1, col = "#00BFC4") + # strat 1
  geom_errorbar(data = beta_CASTHMA_pcs$beta_pcs_strat1, aes(ymin = post_2.5, ymax = post_97.5, width =
  scale_color_manual(name = "Strata",
                     values = c("#F8766D", "#00BFC4"),
                     drop = FALSE)
```





```
ggsave(here("figures/final_figures/sensitivity_analysis/ff_grouped/CASTHMA_cred_intervals_fr_only.pdf")
    plot = p, device = "pdf",
    width = 8, height = 6, units = "in")
```

CAR model results, Poor Mental Health Stratified on RPL_THEMES


```
## strat0:rmax
                            -0.09012 -0.15538 -0.02700
## strat0:Data_Value_CSMOKING  3.32658  3.30587  3.34740
## strat1:Intercept 14.34677 14.33603 14.35735
## strat1:EP_UNINSUR
                            0.04707 0.03457 0.05962
## strat1:pollute_conc_pc1
                           0.68425 0.64453 0.72515
## strat1:pollute_conc_pc2
                            0.38458 0.32977 0.44015
## strat1:pollute_conc_pc3
                            -0.48958 -0.54141 -0.43260
## strat1:tmmx
                             0.07988 -0.00084 0.16941
## strat1:rmax
                            -0.09826 -0.16434 -0.03545
## strat1:Data_Value_CSMOKING 3.28702 3.27128 3.30301
saveRDS(beta_inference, file = here("modeling_files/sensitivity_analysis/ff_grouped/beta_inference_file
List of significant beta coefficients:
row.names(beta_inference)[sign(beta_inference[, 2]) == sign(beta_inference[, 3])]
## [1] "strat0:Intercept"
                                   "strat0:flood_ff_group1"
## [3] "strat0:pollute_conc_pc1"
                                   "strat0:pollute_conc_pc2"
## [5] "strat0:pollute_conc_pc3"
                                   "strat0:rmax"
## [7] "strat0:Data_Value_CSMOKING" "strat1:Intercept"
## [9] "strat1:EP_UNINSUR"
                                   "strat1:pollute_conc_pc1"
## [11] "strat1:pollute_conc_pc2"
                                   "strat1:pollute_conc_pc3"
## [13] "strat1:rmax"
                                   "strat1:Data_Value_CSMOKING"
```

Credible Interval plots for the coefficients, in ggplot

```
# first, process the beta_inference matrix in a form ggplot can understand
beta_inference_df <- as.data.frame(beta_inference)</pre>
beta_inference_df <- mutate(beta_inference_df, var_name = row.names(beta_inference_df))
beta_inference_df <- rename(beta_inference_df,</pre>
                             post_median = `50%`,
                             post_2.5 = 2.5\%,
                             post_97.5 = `97.5\%`)
beta_inference_df$var_name <- substring(beta_inference_df$var_name, first = 8)</pre>
beta_inference_df$var_name <- factor(beta_inference_df$var_name,</pre>
                                      levels = unique(beta_inference_df$var_name))
beta_inference_df$strat <- factor(c(rep("Low SV", (nrow(beta_inference_df)/2)),
                                     rep("High SV", (nrow(beta_inference_df)/2))), levels = c("Low SV",
Splitting up the beta coefficients for each strata
beta_inference_df_strat0 <- beta_inference_df[1:(nrow(beta_inference_df)/2),]
beta_inference_df_strat1 <- beta_inference_df[(nrow(beta_inference_df)/2 + 1):nrow(beta_inference_df),]
Note: The intercept for both strata is not included.
p <- ggplot(beta_inference_df_strat0[-1, ], aes(x = var_name, y = post_median, color = strat)) +
  geom_point() +
  ylim(c(-1.5, 4)) +
  theme(axis.text.x = element_text(angle = 45, vjust = 1, hjust=1), axis.title.x = element_blank(), axi
    axis.text=element_text(size=12),
```

```
plot.margin = margin(5.5, 5.5, 5.5, 25)) +
  geom_errorbar(aes(ymin = post_2.5, ymax = post_97.5, width = 0.4), col = "#F8766D") +
  geom_vline(xintercept = c(3.5, 4.5, 7.5, 9.5), col = "blue") +
  geom_hline(yintercept = 0, col = "red") +
  annotate(geom = "text", x = 2, y = 3.75, label = "Flood\nRisk",
           col = "blue", size = 4.5) +
  scale_x_discrete(labels = c("FF Group 1", "FF Group 2", "FF Group 3", "Uninsured",
                              "Pollution PC 1", "Pollution PC 2", "Pollution PC 3",
                              "Maximum Temperature", "Maximum Humidity",
                              "Smoking")) + ggtitle("95% Credible Intervals, CASTHMA, Stratified on All
  geom_point(data = beta_inference_df_strat1[-1, ], col = "#00BFC4") + # strat 1
  geom_errorbar(data = beta_inference_df_strat1[-1, ], aes(ymin = post_2.5, ymax = post_97.5, width = 0
  scale_color_manual(name = "Strata",
                     values = c("#F8766D", "#00BFC4"),
                     drop = FALSE)
p
```

95% Credible Intervals, CASTHMA, Stratified on All RPL Themes, Flood Fa



95% Credible Intervals for Flood Risk Groupings, Poor Mental Health

