

Homework 1

1 Exercise 1

2 Exercise 2

2.1

In this exercise we will prove the following equality:

$$\int_0^T t dW(t) = TW(T) - \int_0^T W(t) dt \quad (1)$$

Look at the left hand side, by taking it's forward Euler we obtain

$$I_1 = \int_0^T t dW(t) = \sum_{n=0}^{N-1} t_n (W(t_{n+1}) - W(t_n)). \quad (2)$$

Applying the hint, i.e. using the Abel's summation by parts we get

$$\begin{aligned} \sum_{n=0}^{N-1} t_n (W(t_{n+1}) - W(t_n)) &= t_N W(t_N) - t_0 W(t_0) - \sum_{k=1}^{N-1} W(t_k) (t_k - t_{k-1}) \\ &= TW(T) - \sum_{k=1}^{N-1} W(t_k) (t_k - t_{k-1}). \end{aligned}$$

What is left to prove is the convergence in L_2 of the right hand side, i.e. the following equation.

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\left(\int_0^T W(t) dt - \sum_{k=1}^{N-1} W(t_k) (t_k - t_{k-1}) \right)^2 \right] = 0. \quad (3)$$

Let's fix $t_n - t_{n-1} = \Delta t$ for every $n \geq 0$. Using the linearity of the integral, we can rewrite it as follows

$$\begin{aligned} \mathbb{E} \left[\left(\sum_{n=0}^{N-1} \int_{t_{n-1}}^{t_n} W(t) dt - \sum_{k=1}^{N-1} W(t_k) (t_k - t_{k-1}) \right)^2 \right] &= \mathbb{E} \left[\left(\sum_{n=0}^{N-1} \int_{t_{n-1}}^{t_n} W(t) - W(t_n) dt \right)^2 \right] \\ &= \sum_{n=0}^{N-1} \mathbb{E} \left[\int_{t_{n-1}}^{t_n} W(t) - W(t_n) dt \right]^2 \\ &= \sum_{n=0}^{N-1} \mathbb{E} [e_n^2]. \end{aligned}$$

Where e_n is n-th error term. Moreover, in the last equation we dropped the cross terms since $\mathbb{E}[e_i e_j] = 0$ by the properties of the Brownian motion.

Next, we compute $\mathbb{E}[e_n^2]$ for every $n \geq 0$.

$$\begin{aligned}\mathbb{E}[e_n^2] &= \mathbb{E}\left[\left(\int_{t_n}^{t_{n+1}} W(t) - W(t_n) dt\right)^2\right] \\ &= \mathbb{E}\left[\int_{t_n}^{t_{n+1}} \int_{t_n}^{t_{n+1}} (W(t) - W(t_n)) (W(s) - W(t_n)) ds dt\right] \\ &= \mathbb{E}\left[\int_{t_n}^{t_{n+1}} \int_{t_n}^{t_{n+1}} (W(t) - W(t_n)) (W(s) - W(t_n)) dt ds\right].\end{aligned}$$

Where we used twice Fubini's Theorem. We also use Fubini for the following equality.

$$\begin{aligned}\mathbb{E}\left[\int_{t_n}^{t_{n+1}} \int_{t_n}^{t_{n+1}} (W(t) - W(t_n)) (W(s) - W(t_n)) dt ds\right] &= \int_{t_n}^{t_{n+1}} \int_{t_n}^{t_{n+1}} \mathbb{E}[(W(t) - W(t_n)) (W(s) - W(t_n))] dt ds \\ &= \int_{t_n}^{t_{n+1}} \int_{t_n}^{t_{n+1}} (\min(t, s) - t_n) dt ds \\ &= \frac{1}{3}(\Delta t)^3.\end{aligned}$$

Finally, by summing all the values, we get

$$\sum_{n=0}^N \mathbb{E}[e_n] = \sum_{n=0}^N \frac{1}{3}(\Delta t)^3 = \frac{T^3}{n^2}. \quad (4)$$

It follows that by letting n go to infinity, the Forward Euler converges in L_2 .

2.2

As in the hint, we now prove the following

$$\sum_{n=0}^{N-1} W(t_n)(W(t_{n+1}) - W(t_n)) = \sum_{n=0}^{N-1} \frac{W(t_{n+1})^2 - W(t_n)^2}{2} - \frac{(W(t_{n+1}) - W(t_n))^2}{2} \quad (5)$$

We start by looking at the right hand side.

$$\begin{aligned}
\sum_{n=0}^{N-1} \frac{W(t_{n+1})^2 - W(t_n)^2}{2} - \frac{(W(t_{n+1}) - W(t_n))^2}{2} &= \\
&= \sum_{n=0}^{N-1} \frac{W(t_{n+1})^2 - W(t_n)^2}{2} - \frac{W(t_{n+1})^2 - 2W(t_{n+1})W(t_n) + W(t_n)^2}{2} \\
&= \sum_{n=0}^{N-1} \frac{2W(t_n)W(t_{n+1}) - 2W(t_n)^2}{2} = \sum_{n=0}^{N-1} W(t_n)(W(t_{n+1}) - W(t_n))
\end{aligned}$$

Note that the

$$\sum_{n=0}^{N-1} \frac{W(t_{n+1})^2 - W(t_n)^2}{2} = \frac{W(T)}{2} \tag{6}$$

as it is a telescopic sum. To finalize the proof, we need to show that the second part converges to $\frac{T}{2}$ or in other words that

$$\mathbb{E} \left[\left(\sum_{n=0}^{N-1} [W(t_{n+1}) - W(t_n)]^2 - T \right)^2 \right] \rightarrow 0 \tag{7}$$