

 Un sitio web oficial del gobierno de los Estados Unidos. [Así es como lo sabes](#)

Institutos Nacionales de Salud / Biblioteca Nacional de Medicina



[Inicio](#) → [Medicamentos, hierbas y suplementos](#) → Repaglinida

URL de esta página: <https://medlineplus.gov/druginfo/meds/a600010.html>

Repaglinida

pronunciado como (re pag' lin ide)

¿Por qué se prescribe este medicamento?

La repaglinida se utiliza para tratar la diabetes tipo 2 (una afección en la que el cuerpo no utiliza la insulina de forma normal y, por lo tanto, no puede controlar la cantidad de azúcar en la sangre). La repaglinida ayuda al cuerpo a regular la cantidad de glucosa (azúcar) en la sangre. Disminuye la cantidad de glucosa al estimular al páncreas para que libere insulina.

Con el tiempo, las personas que tienen diabetes y niveles altos de azúcar en sangre pueden desarrollar complicaciones graves o potencialmente mortales, como enfermedades cardíacas, accidentes cerebrovasculares, problemas renales, daño a los nervios y problemas oculares. Tomar medicamentos, hacer cambios en el estilo de vida (por ejemplo, dieta, ejercicio, dejar de fumar) y controlar regularmente el nivel de azúcar en sangre puede ayudar a controlar la diabetes y mejorar la salud. Esta terapia también puede reducir las probabilidades de sufrir un ataque cardíaco, un accidente cerebrovascular u otras complicaciones relacionadas con la diabetes, como insuficiencia renal, daño a los nervios (piernas o pies entumecidos o fríos; disminución de la capacidad sexual en hombres y mujeres), problemas oculares, incluidos cambios o pérdida de la visión, o enfermedad de las encías. Su médico y otros proveedores de atención médica hablarán con usted sobre la mejor manera de controlar la diabetes.

Este medicamento a veces se prescribe para otros usos; consulte a su médico o farmacéutico para obtener más información.

¿Cómo se debe utilizar este medicamento?

La repaglinida se presenta en forma de comprimidos para tomar por vía oral. Los comprimidos se toman antes de las comidas, en cualquier momento desde 30 minutos antes de una comida hasta justo antes de la comida. Si se saltea una comida, debe omitir la dosis de repaglinida. Si agrega una comida extra, debe tomar una dosis adicional de repaglinida. Su médico puede aumentar gradualmente su dosis, dependiendo de su respuesta a la repaglinida. Siga atentamente las instrucciones de la etiqueta de su receta y pida a su médico o farmacéutico que le explique cualquier parte que no comprenda. Tome repaglinida exactamente como se le indique. No tome más ni menos del medicamento ni lo tome con más frecuencia que lo indicado en la etiqueta del envase o que lo prescriba su médico.

Continúe tomando repaglinida incluso si se siente bien. No deje de tomar repaglinida sin consultar con su médico.

¿Qué precauciones especiales debo seguir?

Antes de tomar repaglinida,

- tell your doctor and pharmacist if you are allergic to repaglinide or any other drugs.
- tell your doctor if you are taking gemfibrozil (Lopid). Your doctor will probably tell you not to take repaglinide if you are taking this medication.
- tell your doctor and pharmacist what prescription and nonprescription medications, vitamins, nutritional supplements, and herbal products you are taking or plan to take. Be sure to mention any of the following: acetophenazine (Tindal), aspirin, blood pressure medicines, carbamazepine (Tegretol), chloramphenicol (Chloromycetin), chlorpromazine (Thorazine), corticosteroids, diuretics ('water pills'), drugs for arthritis, erythromycin, troglitazone (Rezulin), estrogens, fluphenazine (Prolixin), isoniazid (Rifamate), ketoconazole (Nizoral), mesoridazine (Serentil), oral contraceptives, perphenazine (Trilafon), phenelzine (Nardil), phenobarbital (Luminal), phenytoin (Dilantin), probenecid (Benemid), prochlorperazine (Compazine), promazine (Sparine), promethazine (Phenergan), rifampin (Rifadin, Rimactane), thioridazine (Mellaril), tranlycypromine (Parnate), trifluoperazine (Stelazine), triflupromazine (Vesprin), trimeprazine (Temaril), vitamins, or warfarin (Coumadin). Your doctor may need to change the doses of your medications or monitor you carefully for side effects.
- tell your doctor if you have or have ever had liver or kidney disease or if you have been told you have type I diabetes mellitus.
- tell your doctor if you are pregnant, plan to become pregnant, or are breast-feeding. If you become pregnant while taking repaglinide, call your doctor.
- if you are having surgery, including dental surgery, tell the doctor or dentist that you are taking repaglinide.

What special dietary instructions should I follow?

Be sure to follow all exercise and dietary recommendations made by your doctor or dietitian. It is important to eat a healthful diet.

Alcohol may cause a decrease in blood sugar. Ask your doctor about the use of alcoholic beverages while you are taking repaglinide.

What should I do if I forget a dose?

If you have just begun to eat a meal, take the missed dose as soon as you remember it. However, if you have finished eating, skip the missed dose and continue your regular dosing schedule. Do not take a double dose to make up for a missed one.

What side effects can this medication cause?

This medication may cause changes in your blood sugar. You should know the symptoms of low and high blood sugar and what to do if you have these symptoms.

You may experience hypoglycemia (low blood sugar) while you are taking this medication. Your doctor will tell you what you should do if you develop hypoglycemia. He or she may tell you to check your blood sugar, eat or drink a food

or beverage that contains sugar, such as hard candy or fruit juice, or get medical care. Follow these directions carefully if you have any of the following symptoms of hypoglycemia:

- shakiness
- dizziness or lightheadedness
- sweating
- nervousness or irritability
- sudden changes in behavior or mood
- headache
- numbness or tingling around the mouth
- weakness
- pale skin
- hunger
- clumsy or jerky movements

If hypoglycemia is not treated, severe symptoms may develop. Be sure that your family, friends, and other people who spend time with you know that if you have any of the following symptoms, they should get medical treatment for you immediately.

- confusion
- seizures
- loss of consciousness

Call your doctor immediately if you have any of the following symptoms of hyperglycemia (high blood sugar):

- extreme thirst
- frequent urination
- extreme hunger
- weakness
- blurred vision

If high blood sugar is not treated, a serious, life-threatening condition called diabetic ketoacidosis could develop. Call your doctor immediately if you have any of these symptoms:

- dry mouth
- upset stomach and vomiting
- shortness of breath
- breath that smells fruity
- decreased consciousness

Repaglinide may cause side effects. Tell your doctor if any of these symptoms are severe or do not go away:

- headache
- nasal congestion
- joint aches
- back pain
- constipation
- diarrhea

What should I know about storage and disposal of this medication?

Keep this medication in the container it came in, tightly closed and out of reach of children. Store it at room temperature and away from excess heat and moisture (not in the bathroom).

Unneeded medications should be disposed of in special ways to ensure that pets, children, and other people cannot consume them. However, you should not flush this medication down the toilet. Instead, the best way to dispose of your medication is through a medicine take-back program. Talk to your pharmacist or contact your local garbage/recycling department to learn about take-back programs in your community. See the FDA's Safe Disposal of Medicines website (<https://goo.gl/c4Rm4p> [<https://goo.gl/c4Rm4p>]) for more information if you do not have access to a take-back program.

It is important to keep all medication out of sight and reach of children as many containers (such as weekly pill minders and those for eye drops, creams, patches, and inhalers) are not child-resistant and young children can open them easily. To protect young children from poisoning, always lock safety caps and immediately place the medication in a safe location – one that is up and away and out of their sight and reach. <https://www.upandaway.org> [<https://www.upandaway.org>]

In case of emergency/overdose

In case of overdose, call the poison control helpline at 1-800-222-1222. Information is also available online at <https://www.poisonhelp.org/help> [<https://www.poisonhelp.org/help>] . If the victim has collapsed, had a seizure, has trouble breathing, or can't be awakened, immediately call emergency services at 911.

What other information should I know?

No falte a ninguna cita con su médico ni con el laboratorio. Debe controlar periódicamente su nivel de azúcar en sangre y de hemoglobina glucosilada (HbA1c) para determinar su respuesta a la repaglinida. Su médico también le indicará cómo controlar su respuesta a este medicamento midiendo sus niveles de azúcar en sangre o en orina en casa. Siga estas instrucciones cuidadosamente.

Usted siempre debe usar una pulsera de identificación diabética para asegurarse de recibir el tratamiento adecuado en caso de emergencia.

No deje que nadie más tome su medicamento. Pregúntele a su farmacéutico cualquier duda que tenga sobre cómo renovar su receta.

Es importante que usted mantenga una lista escrita de todos los medicamentos que está tomando, ya sean recetados o no, incluyendo vitaminas, minerales y suplementos dietéticos. Debe tener la lista cada vez que visita su médico o cuando es admitido en un hospital. También es una información importante para llevar consigo en casos de emergencia.

Nombres de marca

- Prandin®

Nombres comerciales de productos combinados

- Prandimet ® (contiene metformina, repaglinida)

Última revisión: 15/07/2016

[Aprenda a citar esta página](#)



[Descargo de responsabilidad de la Sociedad Estadounidense de Farmacéuticos de Sistemas de Salud, Inc.](#)

Información sobre medicamentos para pacientes de AHFS® . © Copyright, 2024. The American Society of Health-System Pharmacists® · 4500 East-West Highway, Suite 900, Bethesda, Maryland. Todos los derechos reservados.

La duplicación para uso comercial debe contar con la autorización de ASHP.

Biblioteca Nacional de Medicina 8600 Rockville Pike, Bethesda, Maryland 20894

Departamento de Salud y Servicios Humanos de los Estados Unidos Institutos Nacionales de Salud