

# Time complexity analysis of Largest sub-array sum

Dr. Anand M

Id: 102763

Assistant Professor

Department of Computer Science and Engineering

SRM Institute of Science and Technology

# Max Subarray: Algorithm Efficiency

We finally have:

$$T(n) = \begin{cases} \Theta(1) & \text{if } n \leq 1, \\ 2T(n/2) + \Theta(n) & \text{if } n > 1. \end{cases}$$

The recurrence has the same form as that for MERGESORT, and thus we should expect it to have the same solution  $T(n) = \Theta(n \lg n)$ .

This algorithm is clearly substantially faster than any of the brute-force methods. It required some cleverness, and the programming is a little more complicated – but the payoff is large.

# Time complexity analysis

$$T(n) = \begin{cases} O(1) & \text{if } n = 1 \\ 2T(n/2) + O(n) & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &\leq 2T\left(\frac{n}{2}\right) + cn \\ &\leq 2\left[2T\left(\frac{n}{4}\right) + c\frac{n}{2}\right] + cn = 4T\left(\frac{n}{4}\right) + 2cn && \text{1st expansion} \\ &\leq 4\left[2T\left(\frac{n}{8}\right) + c\frac{n}{4}\right] + 2cn = 8T\left(\frac{n}{8}\right) + 3cn && \text{2nd expansion} \\ &\vdots \\ &\leq 2^k T\left(\frac{n}{2^k}\right) + kcn && \text{kth expansion} \end{aligned}$$

The expansion stops when  $2^k = n$

$$\begin{aligned} T(n) &\leq nT(1) + cn \log_2 n \\ &= O(n) + O(n \log n) \\ &= O(n \log n) \end{aligned}$$

# Time complexity analysis

- Theorem

$$T(n) = \begin{cases} O(1) & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + T(\lfloor n/2 \rfloor) + O(n) & \text{if } n \geq 2 \end{cases} \Rightarrow T(n) = O(n \log n)$$

- Proof

- There exists positive constant  $a, b$  s.t.  $T(n) \leq \begin{cases} a & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + T(\lfloor n/2 \rfloor) + b \cdot n & \text{if } n \geq 2 \end{cases}$
- Use induction to prove  $T(n) \leq 2b \cdot n \log_2 n + a \cdot n$

- $n = 1$ , trivial

- $n > 1, \lceil \frac{n}{2} \rceil \leq \frac{n}{\sqrt{2}}$

$$T(n) \leq T(\lceil n/2 \rceil) + T(\lfloor n/2 \rfloor) + b \cdot n$$

Inductive hypothesis  $\leq 2b \cdot (\lceil n/2 \rceil \log_2 \lceil n/2 \rceil) + a \cdot \lceil n/2 \rceil + 2b \cdot (\lfloor n/2 \rfloor \log_2 \lfloor n/2 \rfloor) + a \cdot \lfloor n/2 \rfloor + b \cdot n$

$$\leq 2b \cdot (\lceil n/2 \rceil \log_2 \frac{n}{\sqrt{2}}) + a \cdot \lceil n/2 \rceil + 2b \cdot (\lfloor n/2 \rfloor \log_2 \frac{n}{\sqrt{2}}) + a \cdot \lfloor n/2 \rfloor + b \cdot n$$

$$= 2b \cdot n(\log n - \log_2 \sqrt{2}) + a \cdot n + b \cdot n = 2b \cdot n \log_2 n + a \cdot n$$