

Postal address:
Postnet Suite 490
Private Bag x1
Die Wilgers
Republic of South Africa



Physical address:
Tshwane University of Technology campus
175 Nelson Mandela drive
Pretoria
Republic of South Africa

A38/01/06

ABSTRACT FOR A38/01/06

DETERMINATION OF THE ACUTE DERMAL TOXICITY OF F10 DISINFECTANT INSECTICIDE SPRAY (BATCH: Bn 282)

Experimental Starting Date: 28 Aug 2006
Experimental Completion Date: 02 Nov 2006

The test system was in accordance with the OPPTS 870.1200: Acute Dermal Toxicity Standards.

Fifteen female Sprague Dawley rats, 50 days of age were used. Only young healthy animals with no skin lesions were admitted to the study. No animals were withdrawn from the study.

A total of three dosage groups were used in this study. The dosage groups were chosen on the basis of previous literature, which indicates that the dermal toxicity of F10 superconcentrate is very low.

The first group was treated with a concentration of 300mg/kg F10 Disinfectant Insecticide Spray, the second group with 2000 mg/kg and the third with 5000 mg/kg.

The animals were observed 2 hours after exposure and once every day thereafter for a period of 14 days for signs of toxicity.

No toxicological symptoms were recorded in any of the experimental groups during the study period. Furthermore no pathological lesions were recorded in any of the animals at the post mortems. There were also no significant weight loss recorded

This study indicated that the dermal toxicity of the F10 Disinfectant Insecticide Spray lies above 5000 mg/kg.

According to EPA Toxicity Categories this falls within category III.

Dr D J Goosen (BVSc Hons)

Acute Toxicity Categories for Pesticide Products

Hazard Indicators	I	II	III	IV
Dermal LD50.....	Up to and including 200 mg/kg	200 thru 2000 mg/kg	2000 thru 20,000 mg/kg	>20,000 mg/kg