

Postal address:
Postnet Suite 490
Private Bag x1
Die Wilgers
Republic of South Africa



Physical address:
Tshwane University of Technology campus
175 Nelson Mandela drive
Pretoria
Republic of South Africa

A37/01/06

ABSTRACT FOR A37/01/06

DETERMINATION OF THE ACUTE ORAL TOXICITY OF F10 DISINFECTANT INSECTICIDE SPRAY (BATCH: Bn/282)

Experimental Starting Date: 28 Jun 2006
Experimental Completion Date: 16 Nov 2006

The test system was in accordance with the OPPTS 870.1100: Acute Oral Toxicity Standards.

Eleven female Sprague Dawley rats, 50 days of age were used. Only young healthy animals with no skin lesions were admitted to the study. No animals were withdrawn from the study.

A total of three dosage groups were used in this study. The dosage groups were chosen on the basis of previous literature, which indicates that the dermal toxicity of F10 superconcentrate is very low.

Sighting Study:

This procedure is necessary to identify the correct dosage to be used in the main study and to eliminate unnecessary stress on the animals.

Three female Sprague Dawley rats were used during the sighting study. The dosage groups used in this study were 300, 2000 and 5000 mg/kg. The observation period after dosing was 14 days.

It was concluded that the dosage level to be used in the main study was 2000 mg/kg

Main Study:

Four female Sprague Dawley rats were dosed at a level of 2000 mg/kg. None of the rats in this study presented with overt toxicological symptoms.

This study indicated that the oral toxicity of the F10 Disinfectant Insecticide Spray between 2000 and 5000 mg/kg.

According to EPA Toxicity Categories this falls within category III.


Study Director: Mrs. A. Kleynhans (MSc)


Dr D J Goosen (BVSc Hons)

Acute Toxicity Categories for Pesticide Products

Hazard Indicators	I	II	III	IV
Oral LD50.....	Up to and including 50 mg/kg	50 thru 500 mg/kg	500 thru 5,000 mg/kg	>5,000 mg/kg