

Fact Sheet: Emergency Use Authorization of F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) for New World Screwworm (NWS)

F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) is intended for use in cattle, horses, minor species¹ of hoof stock, raptors and other wild birds, pet birds, and captive and captured² wild, exotic, and zoo mammals and contains benzalkonium chloride 0.405 g/100 mL, polyhexanide 0.03 g/100 mL, and cypermethrin 0.25 g/100 g.

Original EUA Authorized Date: 03/10/2026

Emergency Use Authorization for F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) for NWS

The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) for the emergency use of the indexed product F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) for the prevention and treatment of infestations caused by New World screwworm (*Cochliomyia hominivorax*) larvae (myiasis) in cattle, horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals. F10 Antiseptic Wound Spray with Insecticide is not approved or indexed for this use.

F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) is indexed for use as a topical antiseptic for surface wounds, to repel flies and to treat infestations due to fly strike in raptors, pet birds, captive small mammals, captive reptiles and captive exotic/zoo mammals (Minor Species Index File (MIF) 900-010).³

Limitations of Authorized Use

It is a violation of Federal law to use this drug product other than as directed in this fact sheet.

Cattle, goats, and sheep must not be slaughtered for human consumption within 30 days of treatment.

For non-domesticated minor species hoof stock,⁴ use only when there is a reasonable certainty that the treated animal will not be slaughtered or harvested for human consumption within 30 days of treatment.

Milk taken from cows, goats, or sheep during treatment and for 10 days after treatment must not be used for human consumption.

¹ Minor species are all animals, other than humans, that are not one of the major species. They include animals such as zoo animals, parrots, ferrets, and guinea pigs. Some animals of agricultural importance are also minor species. These include sheep, goats, and game birds, among others. The term 'major species' means cattle, horses, swine, chickens, turkeys, dogs, and cats.

² Definitions for "captive wildlife" and "captured wildlife" can be found in the last section of the USDA APHIS New World Screwworm Response Playbook, accessible at <https://www.aphis.usda.gov/animal-emergencies/nws>.

³ See FDA's webpage "The Index of Legally Marketed Unapproved New Animal Drugs for Minor Species" at <https://www.fda.gov/animal-veterinary/minor-use/minor-species/index-legally-marketed-unapproved-new-animal-drugs-minor-species>

⁴ For example, deer, elk, antelope, and nilgai.

A withdrawal period has not been established for this product in pre-ruminating calves. Treated calves and calves born to treated cows must not be processed for veal.

Do not use in horses intended for human consumption.

Do not use in domestic dogs and cats.

Do not use in food-producing species that have not been assigned a withdrawal period, as listed in the Withdrawal Periods and Residue Warnings section.

F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) is authorized for this use only for the duration of the declaration that circumstances exist justifying the authorization of the emergency use of F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) under Section 564(b)(1) of the Federal Food, Drug, and Cosmetic Act (FD&C Act), 21 U.S.C. § 360bbb-3(b)(1), unless the declaration is terminated or the authorization is revoked sooner.

Product Description

F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) is an amber colored topical spray with a slight alcohol odor.

Directions

Use disposable gloves during application and wash hands thoroughly after use. Use only in well-ventilated areas. Clean the wound prior to application. Apply a maximum of 5 applications per animal with multiple wounds (1 - 2 sprays per wound). To ensure full coverage hold the spray 4 - 8 inches away from the wound when spraying. Spray only the wound site. Repeat once a week, if necessary or as advised by your veterinarian.

This product is water soluble and should be reapplied if animal is bathed or exposed to rain. If the wound persists or worsens, contact your veterinarian.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Information Supporting Emergency Use Authorization

Based on the totality of scientific evidence available to FDA, including information submitted in support of the Minor Species Index File (MIF 900-010) and this EUA, as well as publicly available information, it is reasonable to believe that F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) may be effective for the prevention and treatment of infestations caused by NWS larvae (myiasis) in cattle, horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals, and when used under the conditions described in the authorization, the known and potential benefits of F10 Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) outweigh the known and potential risks.

Information to support the effectiveness of F10 Antiseptic Wound Spray with Insecticide under MIF 900-010 were primarily conducted using blowfly species common in South Africa, including *Lucilia cuprina*, the Australian sheep blowfly. (For further details, see the FOI Summary for MIF 900-010.) Cypermethrin, the insecticide in F10 Antiseptic Wound Spray with Insecticide, has documented activity against several other fly species as well, including fly species which are known to cause myiasis in mammals. There is no direct evidence that F10 Antiseptic Wound Spray with Insecticide is effective against *C. hominivorax*. However, both *C. hominivorax* and *L. cuprina* blowflies belong to the family Calliphoridae and generally, blowflies have similar life cycles and morphology. Additionally, the indexed product is labeled for use as a topical antiseptic for surface wounds; it is reasonable to believe that this product would also function as an antiseptic for wounds in cases of NWS myiasis, reducing the risk of secondary infection. Therefore, it is reasonable to conclude that F10 Antiseptic Wound Spray with Insecticide may be effective for the prevention and treatment of NWS myiasis.

The FOI Summary for MIF 900-010 also contains a summary of the safety information regarding the use of F10 Antiseptic Wound Spray with Insecticide in raptors, pet birds, captive small mammals, captive reptiles, and captive exotic/zoo mammals. The anecdotal evidence provided there shows that F10 Antiseptic Wound Spray with Insecticide has been used safely outside of the U.S. in cattle and numerous minor species of cloven-hoofed (ungulate) mammals such as sheep, goats, deer, buffalo, bison, and antelope. Therefore, it is reasonable to conclude that the benefits of using this product in cattle and minor species of hoof stock outweighs the risks.

Cypermethrin is contained in several products registered by the U.S. Environmental Protection Agency (EPA) for use in horses. Potential adverse reactions include hypersensitivity reactions (including urticaria and pruritus) and hair loss. The benefits in horses of using F10 Antiseptic Wound Spray with Insecticide, which contains cypermethrin, outweighs these risks because myiasis is a potentially deadly disease.

F10 Antiseptic Wound Spray with Insecticide is currently indexed for raptors, pet birds, captive small mammals, captive reptiles, and captive exotic/zoo mammals when there is a reasonable certainty that the treated animal will not be consumed by humans or food-producing animals. It is reasonable to include these animals under the EUA; however, captive reptiles are not susceptible to NWS; therefore, they are not included in the list of species authorized for NWS.

FDA evaluated relevant human food safety information and concluded that the food products obtained from treated animals are safe for human consumption when the conditions of use granted by the EUA are followed, including the withdrawal period and milk discard time listed below.

WARNINGS

Withdrawal Periods and Residue Warnings: Cattle, goats, and sheep must not be slaughtered for human consumption within 30 days of treatment. For non-domesticated minor species hoof stock, use only when there is a reasonable certainty that the treated animal will not be slaughtered or harvested for human consumption within 30 days of treatment. Milk taken from cows, goats, or sheep during treatment and for 10 days after treatment must not be used for human consumption. A withdrawal period has not been established for this product in pre-ruminating calves. Treated calves and calves born to treated cows must not be processed for veal.

User Safety Warnings

Not for use in humans. Keep out of reach of children.

Do not use with soaps or other chemicals.

Use disposable gloves during application and wash hands thoroughly after use. Do not contaminate food, water, eating utensils, or food contact surfaces. Wash hands before eating or drinking. If accidentally ingested, contact a Poison Control Center or a doctor. Do not induce vomiting unless advised by the Poison Control Center or a doctor. If accidental eye contact, hold eye open and rinse with water for 10 minutes. Seek medical help if necessary. Use only in well-ventilated areas. Avoid breathing of spray mist.

To obtain Safety Data Sheets, contact Health and Hygiene (Pty) Ltd at 1-800-517-4705 or <https://f10products.com>.

Animal Safety Warnings

Possible side effects that may occur include the development of hives, itchiness, and hair loss. If any of these signs occur, discontinue use of the product on that animal and contact your veterinarian.

Environmental Warnings

This product is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. Do not apply directly to or near water.

Other Warnings

Do not use in horses intended for human consumption.

Do not use in domestic dogs and cats.

Do not use in food-producing species that have not been assigned a withdrawal period, as listed in the Withdrawal Periods and Residue Warnings section.

Resistance can develop to any drug used to treat and prevent NWS myiasis. Reports in scientific literature provide evidence of the development of resistance to pyrethrins such as

cypermethrin in *C. hominivorax* populations in South America. The effectiveness of F10 Antiseptic Wound Spray with Insecticide for the prevention and treatment of NWS infestations should be closely monitored. If the wound is not healing properly, consult your veterinarian.

Reporting Side Effects

Reporting of side effects potentially related to F10 Antiseptic Wound Spray with Insecticide use under this EUA is strongly encouraged.

Report side effects, lack of effectiveness, and product defects using any of these methods:

1. Contact Health and Hygiene (Pty) Ltd at 1-800-517-4705 or <https://f10products.com>, or
2. Download and submit Form FDA 1932a available at <https://www.fda.gov/reportanimalae>, or
3. Contact FDA at 1-888-FDA-VETS to request this form.

When reporting side effects on Form FDA 1932a, provide the following information when available:

- Age, species and breed, sex, and weight of animal(s)
- Overall health status, number of animals treated, and number of animals affected
- Write "F10 Antiseptic Wound Spray with Insecticide use for NWS under an EUA" in the section labeled "**Adverse Event/Product Problem/Product Use Error.**"
- Describe the signs you observed, when they started in relation to the medication, how long they lasted, any treatment given by you or your veterinarian, and whether/when the animal(s) recovered.
- Note any pre-existing health problems of the animal(s) and any other medications or treatments they are currently receiving.
- Provide details about the use of the product, including dose given, the route of administration, and lot number.

Contact Information

For technical assistance, contact Health and Hygiene (Pty) Ltd at 1-800-517-4705 or <https://f10products.com>.

Justification for Emergency Use of Animal Drugs for NWS

The Secretary of the U.S. Department of Health and Human Services (HHS) has:

- determined that there is a significant potential for a public health emergency that has a significant potential to affect national security or the health and security of United States citizens living abroad and that involves NWS (*Cochliomyia hominivorax*); and
- declared that circumstances exist justifying the authorization of emergency use of animal drugs to treat or prevent NWS myiasis in animals.⁵

An EUA is an FDA authorization for the emergency use of an unapproved product or unapproved use of an approved product (i.e., drug, biological product, or device) in the United States under certain circumstances declared by the Secretary of HHS to justify emergency use authorization, including, among others, a determination that there is a public health emergency or a significant potential for a public health emergency that may affect national security and that involves a biological agent.⁶

Criteria for issuing an EUA include:

- The biological agent(s) can cause a serious or life-threatening disease or condition;
- Based on the totality of available scientific evidence (including data from adequate and well-controlled clinical trials, if available), it is reasonable to believe that:
 - o the product may be effective in diagnosing, preventing, or treating the serious or life-threatening disease or condition; and
 - o the known and potential benefits of the product, when used to diagnose, prevent, or treat such disease or condition, outweigh the known and potential risks of the product; and
- There is no adequate, approved, and available alternative to the product for diagnosing, preventing, or treating the serious or life-threatening disease or condition.⁷

⁵ See U.S. Department of Health and Human Services, Declaration of Emergency Pursuant to the Federal Food, Drug, and Cosmetic Act for New World Screwworm, August 20, 2025:
<https://www.federalregister.gov/documents/2025/08/20/2025-15918/declaration-of-emergency-pursuant-to-the-federal-food-drug-and-cosmetic-act-for-new-world-screwworm>

⁶ Emergency Use Authorization of Medical Products and Related Authorities | FDA (<https://www.fda.gov/regulatory-information/search-fda-guidance-documents/emergency-use-authorization-medical-products-and-related-authorities>)

⁷ <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization>.

How Supplied

3.5 oz and 16.9 oz bottles

Dispensing Status

Over the counter (OTC)

Storage Conditions

Store between 15° to 30°C (59° to 86°F) in dry conditions.

Do not reuse container. If empty, place in trash or recycle if available. If partly filled, contact your local solid waste agency for disposal instructions.

Manufactured by:

Health and Hygiene (Pty) Ltd
Cnr. Leader & Hoefsmid Streets
Stormill Ext. 10
South Africa
1-800-517-4705
<https://f10products.com>

Rev. 06/22/2026

Date of Authorization: March 10, 2026
(As revised June 22, 2026)

FREEDOM OF INFORMATION (FOI) SUMMARY

Emergency Use Authorization (EUA)

EUA 006672

F10[®] Antiseptic Wound Spray with Insecticide

(benzalkonium chloride, polyhexanide and cypermethrin topical solution)

Topical spray

Cattle, horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals

Scope of Authorization:

For the prevention and treatment of infestations caused by New World screwworm (*Cochliomyia hominivorax*) larvae (myiasis) in cattle, horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals.

Sponsored by:

Health and Hygiene (Pty) Ltd

Table of Contents

I. GENERAL INFORMATION	3
II. EFFECTIVENESS	6
<i>A. Dosage Regimen</i>	6
<i>B. Information Supporting Emergency Use Authorization</i>	6
III. TARGET ANIMAL SAFETY	6
IV. HUMAN FOOD SAFETY	7
V. USER SAFETY	8
VI. AGENCY CONCLUSIONS	8
<i>A. Duration of Authorization: Revision and Revocation</i>	9
<i>B. Marketing Status</i>	9

I. GENERAL INFORMATION

A. File Number

EUA 006672

B. Sponsor

Health and Hygiene (Pty) Ltd
PO Box 906
Florida Hills, Gauteng, 1716, South Africa

U.S. Agent Name and Address:

Kristen Khanna, PhD, MBA
CEO
Animal Clinical Investigation, LLC
4445 Willard Ave
Sixth Floor
Chevy Chase, MD 20815

C. Proprietary Name

F10[®] Antiseptic Wound Spray with Insecticide

D. Drug Product Established Name

benzalkonium chloride, polyhexanide and cypermethrin topical solution

E. Pharmacological Category

Ectoparasiticide and antiseptic

F. Dosage Form

Topical spray

G. Amount of Active Ingredient

Benzalkonium chloride 0.405 g/100 mL, polyhexanide 0.03 g/100 mL, and
cypermethrin 0.25 g/100 g

H. How Supplied

3.5 and 16.9 fl. oz. bottles

I. Dispensing Status

Over the counter (OTC)

J. Dosage Regimen

Use disposable gloves during application and wash hands thoroughly after use. Use only in well-ventilated areas. Clean the wound prior to application. Apply a maximum of 5 applications per animal with multiple wounds (1 - 2 sprays per wound). To ensure full coverage hold the spray 4 - 8 inches away from the wound when spraying. Spray only the wound site. Repeat once a week, if necessary or as advised by your veterinarian.

This product is water soluble and should be reapplied if animal is bathed or exposed to rain. If the wound persists or worsens, contact your veterinarian.

K. Route of Administration

Topical spray

L. Species/Classes

Cattle, horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured¹ wild, exotic, and zoo mammals

M. Food and Drug Administration (FDA) Indexed Product Indication

F10[®] Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) is indexed for use as a topical antiseptic for surface wounds, to repel flies and to treat infestations due to fly strike in raptors, pet birds, captive small mammals, captive reptiles and captive exotic/zoo mammals (Minor Species Index File (MIF) 900-010).²

N. Emergency Authorized Uses

For the prevention and treatment of infestations caused by New World screwworm (*Cochliomyia hominivorax*) larvae (myiasis) in cattle, horses, minor species³ of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals.

O. Limitations of Authorized Use

It is a violation of Federal law to use this drug product other than as directed in the fact sheet.

¹ Definitions for "captive wildlife" and "captured wildlife" can be found in the last section of the USDA APHIS New World Screwworm Response Playbook, accessible at <https://www.aphis.usda.gov/animal-emergencies/nws>.

² See FDA's webpage "The Index of Legally Marketed Unapproved New Animal Drugs for Minor Species" at <https://www.fda.gov/animal-veterinary/minor-use/minor-species/index-legally-marketed-unapproved-new-animal-drugs-minor-species>

³ Minor species are all animals, other than humans, that are not one of the major species. They include animals such as zoo animals, parrots, ferrets, and guinea pigs. Some animals of agricultural importance are also minor species. These include sheep, goats, and game birds, among others. The term 'major species' means cattle, horses, swine, chickens, turkeys, dogs, and cats.

Cattle, goats, and sheep must not be slaughtered for human consumption within 30 days of treatment.

For non-domesticated minor species hoof stock,⁴ use only when there is a reasonable certainty that the treated animal will not be slaughtered or harvested for human consumption within 30 days of treatment.

Milk taken from cows, goats, or sheep during treatment and for 10 days after treatment must not be used for human consumption.

A withdrawal period has not been established for this product in pre-ruminating calves. Treated calves and calves born to treated cows must not be processed for veal.

Do not use in horses intended for human consumption.

Do not use in domestic dogs and cats.

Do not use in food-producing species that have not been assigned a withdrawal period, as listed in the Withdrawal Periods and Residue Warnings section of the Fact Sheet.

F10[®] Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) is authorized for this use only for the duration of the declaration that circumstances exist justifying the authorization of the emergency use of F10[®] Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) under Section 564(b)(1) of the Federal Food, Drug, and Cosmetic Act (FD&C Act), 21 U.S.C. § 360bbb-3(b)(1), unless the declaration is terminated or the authorization is revoked sooner.

P. Effect of EUA Revision

Following the authorization of EUA 006672 on March 10, 2026, CVM received several questions about the term “captive” and whether it included target animals that were temporarily held, treated, and then released. To minimize confusion and clarify the intended target animal population, CVM determined that the indication should be revised to clarify that animals that are treated and released are included in the target animal population. This was achieved by adding “and captured” to the “captive wild, exotic, and zoo mammals” part of the indication. FDA adopted the USDA APHIS definitions for “captive wildlife” and “captured wildlife” from their New World Screwworm Response Playbook. Captive wildlife are animals maintained in captivity for exhibition (including zoos), sale, personal use, education, propagation, preservation, protection or hunting purposes with no expectation of returning to, or allowing to move freely in, the wild. Captured wildlife are free-ranging wildlife that are under temporary human control with the intent to be returned to and move freely in the wild.

⁴ For example, deer, elk, antelope, and nilgai.

II. EFFECTIVENESS

A. Dosage Regimen

This EUA does not change the dosage regimen in the previously indexed drug product.

B. Information Supporting Emergency Use Authorization

In accordance with Section 564 of the FD&C Act, the sponsor provided information to support that F10[®] Antiseptic Wound Spray with Insecticide may be effective for the prevention and treatment of infestations caused by New World screwworm (*Cochliomyia hominivorax*) larvae (myiasis) in cattle, horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals. This submission includes information submitted in support of indexing F10[®] Antiseptic Wound Spray with Insecticide under MIF 900-010, information submitted for this EUA, and publicly available information.

Information to support the effectiveness of the addition of F10[®] Antiseptic Wound Spray with Insecticide under MIF 900-010 were primarily conducted using blowfly species common in South Africa, including *Lucilia cuprina*. See the FOI Summary for MIF 900-010 dated February 27, 2015.

Cypermethrin has documented activity against several fly species, including other fly species which are known to cause myiasis in mammals. Additionally, the indexed product is labeled for use as a topical antiseptic for surface wounds; this condition of use would also apply when used in cases of NWS myiasis, reducing the risk of secondary infection. Therefore, F10[®] Antiseptic Wound Spray with Insecticide, which contains cypermethrin as an insecticide ingredient, may be effective for the prevention and treatment of *Cochliomyia hominivorax* myiasis when applied to wounds (prevention) or sites of myiasis (treatment).

III. TARGET ANIMAL SAFETY

FDA did not require target animal safety studies for this authorization. The FOI Summary for F10[®] Antiseptic Wound Spray with Insecticide (MIF 900-010) dated February 27, 2015, contains a summary of the target animal safety conclusions for use of F10[®] Antiseptic Wound Spray with Insecticide as a topical antiseptic for surface wounds, to repel flies, and to treat infestations due to fly strike in raptors, pet birds, captive small mammals, captive reptiles, and captive exotic/zoo mammals.

The indexed product has been used outside the U.S. in cattle and numerous minor species of cloven-hoofed (ungulate) mammals such as sheep, goats, deer, buffalo, bison, and antelope without reports of adverse events. The insecticide component in this spray, cypermethrin, is contained in several products registered by the U.S. Environmental Protection Agency (EPA) for use on horses. These products are widely used on horses in the U.S. without serious safety concerns. Potential adverse reactions include hypersensitivity reactions (including urticaria and pruritus) and hair loss. The benefits in horses of using F10[®] Antiseptic Wound Spray with Insecticide, which contains cypermethrin, outweighs these risks because myiasis is a potentially deadly disease.

The product is currently indexed for raptors, pet birds, captive small mammals, captive reptiles, and captive exotic/zoo mammals when there is a reasonable certainty that the treated animal will not be consumed by humans or food-producing animals. It is reasonable to include these target animals on the fact sheet for this EUA; however, captive reptiles are not susceptible to NWS; therefore, they were not included in the list of target species for NWS.

IV. HUMAN FOOD SAFETY

The human food safety assessment for F10[®] Antiseptic Wound Spray with Insecticide (benzalkonium chloride, polyhexanide and cypermethrin topical solution) was based on publicly available information, with the following as the main sources:

1. U.S. Environmental Protection Agency. (2006, August 10). Toxicology disciplinary chapter for the re-registration eligibility decision (RED) risk assessment: Alkyl dimethyl benzyl ammonium chloride (ADBAC)* [Memorandum].
*Benzalkonium chloride is also called ADBAC.
2. U.S. Environmental Protection Agency. (2008, January 9). Poly (hexamethylene biguanide) hydrochloride (PHMB)*; Exemption from the requirement of a tolerance. Final rule. Federal Register, 73(6), 1512. 40 CFR Part 180. EPA-HQ-OPP-2005-0268; FRL-8345-8.
*Polyhexanide is also called PHMB.
3. World Health Organization. (2004). Cypermethrin and alpha-cypermethrin. WHO Food Additives Series, 53.
4. U.S. Environmental Protection Agency. (2025, July 17). Cypermethrin; Pesticide tolerances. Final rule. Federal Register, 90(135). 40 CFR Part 180. EPA-HQ-OPP-2024-0220; FRL-12817-01-OCSPP.
5. Food and Agriculture Organization (FAO) of the United Nations (1997). Cypermethrin and Alpha-cypermethrin. Residues of some veterinary drugs in animals and foods. FAO Food and Nutrition Paper 41/9.
6. FAO (2002). Cypermethrin and Alpha-cypermethrin. Residues of some veterinary drugs in animals and foods. FAO Food and Nutrition Paper 41/14.
7. FAO (2004). Alpha-cypermethrin and cypermethrin. Residues of some veterinary drugs in animals and foods. FAO Food and Nutrition Paper 41/16.

FDA concluded that the food products obtained from treated animals are safe for human consumption when the conditions of use authorized by the EUA are followed, including the withdrawal period and milk discard time.

For cattle, goats, and sheep, a 30-day withdrawal period and 10-day milk discard time are assigned.

For non-domesticated minor species hoof stock, use only when there is a reasonable certainty that the treated animal will not be slaughtered or harvested for human consumption within 30 days of treatment.

A withdrawal period has not been established for this product in pre-ruminating calves. Treated calves and calves born to treated cows must not be processed for veal.

V. USER SAFETY

The product Fact Sheet contains the following information regarding safety to humans handling, administering, or exposed to F10® Antiseptic Wound Spray with Insecticide:

Not for use in humans. Keep out of reach of children.

Do not use with soaps or other chemicals.

Use disposable gloves during application and wash hands thoroughly after use. Do not contaminate food, water, eating utensils, or food contact surfaces. Wash hands before eating or drinking. If accidentally ingested, contact a Poison Control Center or a doctor. Do not induce vomiting unless advised by the Poison Control Center or a doctor. If accidental eye contact, hold eye open and rinse with water for 10 minutes. Seek medical help if necessary. Use only in well-ventilated areas. Avoid breathing of spray mist.

VI. AGENCY CONCLUSIONS

Based on the totality of scientific evidence available to FDA, including information submitted in support of the Minor Species Index File (MIF 900-010) and this EUA, as well as publicly available information, it is reasonable to believe that F10® Antiseptic Wound Spray with Insecticide when used as authorized, may be effective for the prevention and treatment of infestations caused by New World screwworm (*Cochliomyia hominivorax*) larvae (myiasis) in cattle, horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals; and the known and potential benefits of F10® Antiseptic Wound Spray with Insecticide when used as authorized outweigh the known and potential risks. Additionally, it was concluded that residues in food products derived from cattle, goats, and sheep treated with F10® Antiseptic Wound Spray with Insecticide will not represent a public health concern when the product is used as authorized.

New World screwworm infestations can have significant adverse health consequences and can be fatal if left untreated due to the extensive tissue damage caused by *Cochliomyia hominivorax* larvae. The benefit of preventing or treating this potentially fatal disease outweighs the health risks of using this product in these species.

There is no adequate, approved,⁵ and available alternative to the product for the prevention and treatment of NWS (myiasis) in these species. There are no approved products for horses, minor species of hoof stock, raptors and other wild birds, pet birds, and captive and captured wild, exotic, and zoo mammals. Although there are conditionally approved products for the prevention and treatment of NWS in cattle, F10®

⁵ "Approved" products include conditionally approved products for purposes of EUAs issued under Section 564 of the Federal Food, Drug, and Cosmetic Act, 21 U.S.C. § 360bbb-3.

Antiseptic Wound Spray with Insecticide provides a distinct option for treating and preventing NWS in cattle because it is a topical spray product that is applied locally, directly to wounds. Cattle and other hoof stock are at particular risk of infestation by NWS, and a diverse and sufficient supply of products is needed to adequately address an incursion in the United States.

For additional information on all products authorized or conditionally approved for use to treat and/or prevent New World screwworm, please see FDA's "New World Screwworm: Information for Veterinarians" webpage at <https://www.fda.gov/animal-veterinary/safety-health/new-world-screwworm-information-veterinarians>.

A. Duration of Authorization: Revision and Revocation

This EUA will be effective until revoked under Section 564(g) of the FD&C Act or until the Secretary's declaration of emergency or threat justifying emergency authorized use is terminated (Section 564(f)(1)), with exception for continued use permissible under Section 564(f)(2). FDA may revoke or revise this authorization if emergency use of this animal drug for NWS myiasis is no longer justified, if the product no longer meets the criteria for issuance of an EUA under Section 564(c) of the FD&C Act, or other circumstances make such revocation or revision of the authorization appropriate to protect the public health or safety (Section 564(g)(2) of the FD&C Act).

B. Marketing Status

This product is authorized to be marketed OTC because the authorized labeling contains adequate directions for use by laypersons and the conditions of use prescribed on the label are reasonably certain to be followed in practice.