

Lang's Algebra Chapter 3 Solutions

Amanuel Tewodros

July 14, 2025

- (1) By the second isomorphism theorem, we have

$$\frac{U}{U \cap W} \cong \frac{U + W}{W}.$$

For two vector spaces, $X \supseteq Y$ over a field K , we have $\dim X/Y = \dim X - \dim Y$. Thus $\dim U - \dim U \cap W = \dim U + W - \dim W$.

- (2) Let M be a module over a commutative ring R . Let I be a maximal ideal of R . We first show that for any proper ideal \mathfrak{a} of R and basis set $\{x_1, x_2, \dots\}$, of M ,

Lemma 1.

$$\frac{M}{\mathfrak{a}M} \cong \bigoplus_i \frac{A}{\mathfrak{a}}(x_i + \mathfrak{a}x_i).$$

Proof. $\mathfrak{a}M$ is submodule of M because $\mathfrak{a}M \subseteq M$ by R -closure property of \mathfrak{a} . It immediately follows that $\mathfrak{a}M = \bigoplus_i \mathfrak{a}x_i$. By linear independence of x_i , $(\sum_i r_i x_i) \bmod \mathfrak{a}x_j = (r_j \bmod \mathfrak{a})x_j + \sum_{i \neq j} r_i x_i$. Therefore, $M/\mathfrak{a}M = \bigoplus_i Ax_i/\mathfrak{a}x_i$. By the isomorphism $x_i \mapsto 1_A \mapsto (x_i + \mathfrak{a}x_i)$, $Ax_i/\mathfrak{a}x_i \cong A/\mathfrak{a} \cong A/\mathfrak{a}(x_i + \mathfrak{a}x_i)$. \square

Taking \mathfrak{a} as a maximal ideal of R in the above lemma, we see that $M/\mathfrak{a}M$ is a direct product of vector spaces over the field A/\mathfrak{a} and thus admit a basis of the same cardinality as that of M . Because the dimension of a vector space is independent of the basis choice, M also has a fixed dimension.

- (3) Let $\{x_1, \dots, x_m\}$ form the basis set of R over k and let $1_R = k_1 x_1 + \dots + k_m x_m$ for $k_i \in k$. For any element $a \in R$, define the sequences $\{y_1, \dots, y_m\} \subseteq k$, $\{f_1, f_2, \dots, f_m\} \subseteq R$ as:

$$\begin{aligned} f_1 &= a, & y_1 &= w_{1,1}^{-1} k_1 \\ f_{i+1} &= f_i y_i - k_i x_i, & y_i &= k_i w_{i,i}^{-1}, \end{aligned}$$

,where $f_i = \sum_j w_{i,j} x_j$. By construction, $a^{-1} = \sum_i y_i x_i$. Thus R is a field.

- (4) **Direct Sums**

- (a) First, we show the equivalence of the two statements of the theorem. Suppose there is φ such that $g \circ \varphi = \text{id}$. By the injectivness of the composition, $\text{Im } \varphi \cap \ker g = \{0\}$. But by exactness, $\ker g = \text{Im } f$. We can unambiguously define $\psi(u)$ to be the inverse image of $f^{-1}(u')$ where $u' \equiv u \pmod{\text{Im } \varphi}$ and $u' = f(x)$ for some $x \in M'$ because if $f(x) = f(y) \pmod{\text{Im } \varphi}$, $f(x - y) \in \text{Im } \varphi$ and by injectivity of f , $x = y$. Since $M/\text{Im } f \cong M'' = \text{Im } \varphi$, ψ is defined in all of M . Similarly, if the second statement is true, $\ker \psi \cap \text{Im } f = \{0\}$ because $\psi \circ f$ is injective. By exactness, $\text{Im } f = \ker g$. We can then define $\varphi(u) = u'$ where $u' = y \pmod{\ker \psi}$ and $g(y) = u$ for some y . φ is well-defined because if $g(y_1) = g(y_2)$ for $y_1 \neq y_2$, then $y_1 \neq y_2 \pmod{\ker \varphi}$.

Now suppose $x \in M$. $x - \varphi(u) \in \text{Im } f$ for exactly one u by the argument mentioned previously. Thus we can express $x = r + s$ where $r = \varphi(u) \in \text{Im } \varphi$ and $s = x - \varphi(u) \in \text{Im } f$. This implies $M = \text{Im } f \oplus \text{Im } \varphi$. By bijectivness of $g \circ \varphi$, $\text{Im } \varphi \cong M''$. By contrast, if $M = \text{Im } f \oplus N$ for some N , with isomorphism $t : N \rightarrow M''$. We can define $g : M \rightarrow M''$ as $g(u) = u'$ such that there is $u = y \pmod{N}$ and $t^{-1}(u') = y$. This definition is unambiguous because $N \cap \text{Im } f = \{0\}$. Since $g \circ t^{-1} = \text{id}$, the sequence splits.

Finally, we complete the details of proposition 3.2. We have just shown $M = \text{Im } f \oplus \text{Im } \varphi$. By exactness, $\text{Im } f = \ker g$. Also, $\text{Im } f \cong M'$ and $\text{Im } \varphi \cong M''$ by injectivness of f and φ resp. This proves $M \cong M' \oplus M''$. We can write $x \in M$ as $f(u) + x - f(u)$ where $x - f(u) \in \ker \psi$. u is then uniquely determined by x as $\ker \psi \cap \text{Im } f = \{0\}$ by bijectivness of $\psi \circ f$. This shows $M = \text{Im } f \oplus \ker \psi$.

- (b) First, we note that φ_i is injective because otherwise the composition $\psi_i \circ \varphi_i$ wouldn't be injective, a contradiction. This implies, for every valid i , there is a submodule $E'_i = \text{Im } \varphi_i$ of E that is isomorphic to E_i . Moreover, if $c \in \text{Im } \varphi_i \cap \text{Im } \varphi_j$ for $i \neq j$, then $\psi_i(c) = \psi_j(c) = 0$, forcing c to be 0. These statements prove

$$\bigoplus_{i=1}^n E'_i \subseteq E.$$

The inverse inclusion follows as follows. Let $x \in E$, then $x = \sum_{i=1}^n \varphi_i(\psi_i(x))$, but $\varphi_i(\psi_i(x)) \in E'_i$. Therefore $x \in \bigoplus_i E'_i$.

Let $x = x_1 + \dots + x_m$ where $x_i \in E'_i$. The map defined by $x \mapsto (\psi x_i)_{1 \leq i \leq m}$ is therefore an isomorphism and the inverse map is given by $(\psi x_i)_i \mapsto \sum_i x_i$.

- (5) Let $v'_m = a_1 v_1 + \dots + a_m v_m$. Since $a_m \neq 0$, v'_m , and by the assumption that $\{v_i\}$ is linearly independent over \mathbb{R} , the set $\{v_1, \dots, v_{m-1}, v'_m\}$ is linearly independent over \mathbb{Z} . We also note that, $v'_m - \sum_{i=1}^{m-1} a_i v_i \in A$, thus we can safely assume $a_1 = \dots = a_{m-1} = 0$.

To show, the set spans A , we consider A/A_0 . Suppose, there is $av_m \in A/A_0$ such that $av_m \neq nv'_m$ for all $n \in \mathbb{Z}$. Let r, s be two integers such that $|ra_m + sa| < a_m$. Since contradicts minimality of a_m , it must be the case that $a_m \mid a$. Therefore $A/A_0 = \mathbb{Z}v'_m$.

(6) We induct on the size of S .

First assume that $S = \{w\}$. Then $\mathbb{Z}\langle S \rangle = \{n[w] : n \in \mathbb{Z}\}$. If M is a subgroup of $\mathbb{Z}\langle S \rangle$, then $M = \mathbb{Z}\langle a[w] \rangle$ for some $a \in \mathbb{Z}$. Here we pick $y_w = a[w]$ which is G -linear.

For the induction step, suppose the statement is true for S , $0 \leq |S| \leq m-1$. We shall prove the statement is true for S with m elements. Fix on element $w \in S$, and consider projection map $\pi : \mathbb{Z}\langle S \rangle \rightarrow \mathbb{Z}\langle G \cdot w \rangle$. By correspondence, $\pi(M)$ is a subgroup of $\mathbb{Z}\langle G \cdot w \rangle$ with basis $\{\bar{y}_{gw}\}_{w \in G}$ which satisfy the property for $\sigma \in G$, $\sigma \bar{y}_{gw} = \bar{y}_{\sigma gw}$. We then lift the basis of $\mathbb{Z}\langle \pi(M) \rangle$ to $\mathbb{Z}\langle S \rangle$ by picking a representatives $\mathfrak{R} = \{y_w\}$ in M for \bar{y}_w . The y_w are linearly independent thus form part of the basis for M . Again by hypothesis, $M \cap \mathbb{Z}\langle S - G \cdot w \rangle$ has basis $\mathfrak{B} = \{y_w\}_{w \in S - G \cdot w}$ that satisfy the given property. We finally combine \mathfrak{R} and \mathfrak{B} to get the basis of rank m for M .

(7) For convenience, we identify the properties of a semi-norm as follows

$$\text{SN-1 } |v| \geq 0$$

$$\text{SN-2 } |nv| = |n||v|$$

$$\text{SN-3 } |u + v| \leq |u| + |v|$$

(a) Let $a, b \in M_0$. Then by SN-2 and SN-3, $|u - b| \leq |a| + |b| = 0$. By SN-1, we have $|a - b| \geq 0$, this $a - b \in M_0$. By SN-2, $|0| = |2 \cdot 0| = 2|0|$. This implies $0 \in M_0$. Hence M_0 is a subgroup of M .

(b) If $M_0 \neq \{0\}$, we can make the transformation $x \mapsto x + M_0$ without loss of generality as such map preserves the linear independence of $\{v_i\}$. Thus, we can assume $M_0 = \{0\}$.

Let $N = \langle v_1, \dots, v_r \rangle$. Since M has rank r , the exponent e of M/N is finite and thus eM is a subgroup of N . Moreover, N/eM is torsion group with finite number of elements. Therefore, we can pick the smallest positive integers $n_{i,j}$ such that

$$\sum_{j=1}^i n_{i,j} v_j = dw_i \quad \text{for some } w_i \in M$$

The linear independence follows immediately. Picking $n_{j,k}$ in the range $[0, d-1]$,

$$d|w_i| = |dw_i| \leq \sum_{j=1}^i n_{i,j} |v_j| \leq d \sum_{j=1}^i |v_j|.$$