tends to infinity when k does, for the order of the connectivity of the pair $(F^{k+1}(X), F^k(X))$ and this does not depend on X but only on k.

6.4.4. COROLLARY. If Y is a finite CW-complex of dimension less than 2[k/3] then the natural map

$$[C_0(X), C_0(Y) \otimes M] \rightarrow kk(Y, X)$$

is a bijection.

Proof. Since $F^k(X) = \operatorname{Hom}_1(C(X), M_k)$ is homeomorphic to $F_0^k(X) = \operatorname{Hom}(C_0(X), M_k)$ it follows by Theorem 6.4.2 that the inclusion $F_0^k(X) \hookrightarrow F_0^{k+1}(X)$ is a 2[k/3]-equivalence. Moreover we know that $\lim_{k \to \infty} F_0^k(X)$ is homotopic to F(X) (3.1.2) and so $F_0^k(X) \hookrightarrow F(X)$ is a 2[k/3]-equivalence. Consequently the map $[Y, F_0^k(X)] \hookrightarrow [Y, F(X)]$ is one to one whenever $\dim Y < 2[k/3]$.

6.4.5 REMARK Let X,Y, & as above. Then $\pi_1(F_o^k(X))$ acts trivially on $[Y, F_o^k(X)]$. Therefore $[C_o(X), C_o(Y) \otimes M_k] \cong [C(X), C(Y) \otimes M_k]_1 = k^k(Y,X)$. REFERENCES

- 1. Adams, J.F.: Stable Homotopy and Generalized Homology, Part III, Chicago Lectures in Math., Univ. of Chicago Press, Chicago, 1974.
- 2. Blackadar, B.: Shape theory for C*-algebras, Math. Scand. 56(1985), 249-275.
- 3. Blackadar, B.: K-theory for Operator algebras Springer MSRI Series 1986.
- 4. Blackadar, B.: Comparison theory for simple C*-algebras, preprint.
- 5. Blackadar, B.: Symmetries of the CAR algebra, preprint.
- 6. Borsuk, K.: Theory of shape (Monographie Matematyczne 59), Polska Academia Nauk, Warsaw, 1975.
- 7. Bratteli, O.: Inductive limits of finite dimensional C*-algebras, Trans. Amer. Math. Soc. 171(1972), 195-232.
- 8. Connes, A.: Non-commutative differential geometry, Publ. Math. I.H.E.S. 62(1985), 41-144.
- 9 **Cuntz, J.:** On the homotopy groups of the space of endomorphisms of a C*-algebras (with application to topological Markow Chain) in Operator Algebras and Groups Representation -F. Proc. Internat. Conf. at Neptun, Romania. 1980, Pitman, London, 1984, 124-137

- 26. Kasparov, G.G.: The operator K-functor and extensions of C*-algebras, Izv. Akad. Nauk. SSSR, Ser. Math. 44, (1980), 571-636).
- 27. Loring, T.A.: The K-theory of AF-embeddings of the rational rotation algebras, MSRF-preprint, 1987.
- 28. Loring, T.A.: The noncommutative topology of one dimensional spaces, preprint, 1987.
- 29. Mardesic, S.: Segal, J.: Shape of compacta and ANR-systems, Fund. Math. 72(1971), 41-59.
- 30. Milnor, J.: Construction of universal bundles, II, Ann. of Math. vol.63, No.3, (1956), 430-436.
- 31. Nistor, V.: On the homotopy groups of the automorphism group of AF-algebras, J. Operator Theory, 19(1988), 319-340.
- 32. Nistor, V.: Fields of AF C*-algebras on suspensions, preprint nr. 39, INCREST, 1988.
- 33. Pasnicu, C.: On certain inductive limit C*-algebras, Indiana Univ. Math. J., 35(1986), 269-288.
- 34. Pasnicu, C.: On inductive limits of certain C*-algebras of the form C(X) x F, Trans. Amer. Math. Soc. Vol. 310, No.2, 1988.
- 35. Rieffel, M.A.: The homotopy groups of the unitary groups of noncommutative tori, J. Operator Theory 17(1987), 237-254.
- 36. Rosenberg, J.: The role of K-theory in non-commutative algebraic topology, Contemporary Math. AMS, vol. 10, Providence, 1982.
- 37. Rosenberg, J.; Schochet, C.: The Kunneth Theorem and the Universal Coefficient Theorem for Kasparov's generalized K-functor, Duke Math. J. 55

 (1987), 431-474. + Mermers 348/1886
- 38. Schochet, C.: Topological methods for C*-algebra III: Axiomatic homology, Pacific J. Math. 114(1984), 399-445.
- 39. Segal, G.: K-homology theory and algebraic K-theory, in K-theory and Operator Algebras, Athens, Georgia, 1975, Lecture Notes in Math, no.575, Springer-Verlag, 1977, pp. 113-127.
- 40. Smith, L.: On the relation of conective K-theory to homology Proc. Comb. Phil. Soc. (1970), 68, 637-639.