

Note on Commutative Algebra

JIAWEN HUANG

20 January 2026

Contents

1	January 17, 2026	4
2	January 20, 2026	5

Acknowledgement

This note is basically a copy of Eisenbud's Commutative Algebra with a View Toward Algebraic Geometry, with some additional explanations and examples from Eisenbud's lectures and my own understanding.

This latex style is made by Evan Chen.

Corollary 1.1 (Summand of polynomial rings over fields is finitely generated)

Let k be a field, and let R be a k -subalgebra of $S = k[x_1, \dots, x_n]$. R is a direct summand of S in the sense that there is a surjective R -algebra homomorphism $\phi : S \rightarrow R$ such that $\phi|_R = \text{id}_R$ and ϕ preserves degrees. Then R is finitely generated as a k -algebra.

Proof. Intuition: R itself is a graded k -algebra, with grading inherited from S . As a graded algebra, R can be decomposed into its homogeneous components. R_0 is easy to get, just include 1 in the generators. Therefore, to generate R , we really need to generate the positive degree part. That motivates us to look at the ideal R_+ .

But R_+ may not be finitely generated as an ideal. However, we know that S is a Noetherian ring, so we can look at the ideal R_+S . It must be finitely generated, say by $f_1, \dots, f_m \in R_+$.

Now, we claim that $R = k[1, f_1, \dots, f_m]$. Certainly, the right hand side is contained in the left hand side. For the other direction, it suffices to show that any homogeneous element of R can be generated by $f_1, \dots, f_m$ with coefficients in k . We prove this by induction on degree d . $d = 0$ is trivial. Now suppose we have shown this for degree less than d . Take any homogeneous element $r \in R$ of degree d . Since $r \in R_+S$, we can write $r = \sum g_i f_i$ for some homogeneous element $g_i \in S$. Apply ϕ to both sides, we get

$$r = \phi(r) = \sum \phi(g_i)\phi(f_i) = \sum \phi(g_i)f_i.$$

Note that $\deg(\phi(g_i)) = \deg(g_i) < d$, so by the inductive hypothesis, each $\phi(g_i)$ can be generated by $f_1, \dots, f_m$ over k . Therefore, r can also be generated by $f_1, \dots, f_m$ over k . This finishes the proof. $\square$

Remark 1.2. This shows that the invariant ring of a linearly reductive group acting on a polynomial ring over a field is finitely generated. If the group G is finite, the surjective homomorphism ϕ can be taken as $\phi(f) = \sum_{g \in G} gf$. (Weyl's averaging trick)

Remark 1.3. Graded rings are great because it allows us to use induction on degree. So we only need to worry about homogeneous elements.

2 January 20, 2026

First day of Commutative Algebra course!

We discuss the history of Commutative Algebra and three of the four big theorems.

Theorem 2.1 (Hilbert's Basis Theorem)

If R is a Noetherian ring, then the polynomial ring $R[x]$ is also Noetherian.

Proof. Let I be an ideal of $R[x]$. Our goal is to show that I is finitely generated. We first pick the smallest degree nonzero polynomial f_1 in I . Next, we take f_2 to be the smallest degree polynomial in I which is not in the ideal generated by f_1 , that is $f_2 \in I \setminus (f_1)$. Continuing this process, we get a sequence of polynomials $f_1, f_2, \dots$ in I such that f_n is the smallest degree polynomial in $I \setminus (f_1, \dots, f_{n-1})$. Clearly we have $\deg(f_{i+1}) \leq \deg(f_i)$ for all i . Since R is Noetherian, the leading coefficients of f_i generate a finitely generated ideal in R . This means that there exists j such that the leading coefficient of f_1 to f_j generate all the leading coefficients of f_i for $i > j$.

Now, we look at f_{j+1} . Its leading coefficient can be generated by the leading coefficients of f_1 to f_j , that is $a_{j+1} = \sum_{i=1}^j r_i a_i$ for some $r_i \in R$. Consider the polynomial

$$g = f_{j+1} - \sum_{i=1}^j r_i x^{\deg(f_{j+1}) - \deg(f_i)} f_i$$

Since $\deg(g) < \deg(f_{j+1})$ so $g \in (f_1, \dots, f_j)$. Thus, $f_{j+1} \in (f_1, \dots, f_j)$, contradicting the choice of f_{j+1} . Therefore, $I = (f_1, \dots, f_j)$ is finitely generated. $\square$

Theorem 2.2 (Hilbert's syzygy theorem)

If $R = k[x_1, \dots, x_n]$ is a graded polynomial ring over a field k , then every finitely generated graded R -module M has a finite graded free resolution of length at most n .

Proof is omitted.

Theorem 2.3 (Hilbert's function theorem)

Let M be a finitely generated graded module over the polynomial ring $R = k[x_1, \dots, x_n]$. Then there exists a polynomial $P(t) \in \mathbb{Q}[t]$ such that for all sufficiently large d , we have $\dim_k M_d = H_M(d) = P(d)$.

We will present two proofs of this theorem.

Proof. We will use the Hilbert's syzygy theorem to prove this. By the theorem, we have a finite graded free resolution of M :

$$0 \rightarrow F_n \rightarrow F_{n-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0.$$

Thus, we have $H_M(d) = \sum_{i=0}^n (-1)^i H_{F_i}(d)$. So we only need to study the dimension of free modules F_i , which is a direct sum of $k[x_1, \dots, x_n]$. Note that $H_R(d) = \dim_k R_d = \binom{n+d-1}{n-1}$ (star and bar counting trick), which is a polynomial in d when d is sufficiently large. Therefore, $H_M(d)$ is also a polynomial in d for sufficiently large d . $\square$

Second one is via induction on n .

Proof. $n = 0$ is trivial. Suppose it's true for number less than n . Consider the short exact sequence

$$0 \rightarrow (0 : x_n) \rightarrow M(-1) \xrightarrow{x_n} M \rightarrow M/x_n M \rightarrow 0.$$

Notice that $M/x_n M$ is a finitely generated graded module over $k[x_1, \dots, x_{n-1}]$ (x_n acts as zero) and $(0 : x_n)$ is also a finitely generated graded module over $k[x_1, \dots, x_{n-1}]$ (the generators of $(0 : x_n)$ over $k[x_1, \dots, x_{n-1}, x_n]$ also generate $(0 : x_n)$ over $k[x_1, \dots, x_{n-1}]$ since x_n would act as zero). By the inductive hypothesis, both $M/x_n M$ and $(0 : x_n)$ have Hilbert dimensions agree with polynomials when d is large enough. Thus, by the exactness of the sequence,

$$H_M(d) - H_M(d-1) = H_{M/x_n M}(d) - H_{(0:x_n)}(d-1) = \lambda(d).$$

for some polynomial $\lambda(d)$ when d is large enough. Thus,

$$H_M(d) - H_M(0) = \sum_{i=1}^d (H_M(i) - H_M(i-1)) = \sum_{i=1}^d \lambda(i).$$

Notice that $\sum_{i=1}^d i^k$ is a polynomial in d of degree $k+1$. Therefore, $\sum_{i=1}^d \lambda(i)$ is also a polynomial in d when d is large enough. Therefore, $H_M(d)$ also agrees with a polynomial when d is large enough. $\square$