

Math Daily Log

JIAWEN HUANG

7 November 2025

Acknowledgment

This is Evan Style LaTeX template. Thanks to Evan Chen for creating such a beautiful style file.

October 28, 2025

1. Weyl Algebra $\frac{k\langle x,y \rangle}{xy-yx-1}$.
2. Center of Weyl Algebra is k in characteristic 0.
3. In characteristic p , the center of Weyl Algebra is $k[x^p, y^p]$.
4. Weyl Algebra is simple.
5. x and y can be seen as operators on $k[t]$ by $x : f \mapsto tf$ and $y : f \mapsto \frac{df}{dt}$. This is amazing. I think the structure $xy - yx = 1$ can be used to make some algebra olympiad problems. Very rich and interesting structure. When you interpret as operators on polynomials, everything becomes very concrete and easy to understand. Perfect for a math olympiad problem.

October 29, 2025

1. Splitting field of a polynomial is unique up to isomorphism.
2. Separable. An inseparable polynomial: $x^p - t$ over $\mathbb{F}_p(t)$. $x^p - t = (x - u)^p$ in the algebraic closure.
3. A polynomial is separable iff its derivative and f are coprime. In particular, over characteristic 0, all irreducible polynomials are separable.
4. A field is perfect if all irreducible polynomials are separable iff every algebraic extension is separable.

-
5. A finite extension is separable iff $[F : k]_s = [F : k]$ iff $F = k(\alpha_1, \dots, \alpha_n)$ where α_i are separable.
 6. A finite Normal extension is a splitting field of a polynomial. This is amazing.
 7. Riemann integral is equivalent to Darboux integral.

October 30, 2025

1. Riemann integrable iff discontinuities has measure 0.
2. A lot of exercises on separable extensions from Aluffi chapter 7.
3. Formal power series $k[[x]]$ on math 250 lecture.

Nov. 1, 2025

1. Cyclotomic polynomial is in $\mathbb{Z}[x]$, and it's always irreducible. Proof used Dirichlet's Theorem and the fact that finite fields are perfect. Too smart.
2. Finite Field of order p^r is unique up to isomorphism, it's the splitting field of $x^{p^r} - x$.
3. $x^4 + 1$ is reducible in all finite field. But irreducible in $\mathbb{Z}[x]$. Proof use the fact that $x^4 + 1 \mid x^8 - 1 \mid x^{p^2} - x$. So the root β of $x^4 + 1$ in F_{p^2} has degree of 1 or 2. Thus, its minimal polynomial has degree of 1 or 2.
4. Let $F = \mathbb{F}_q$, $x^{p^n} - x$ is the product of all the irreducible polynomial of degree d for all $d \mid n$.
5. The automorphism $\text{Aut}_{\mathbb{F}_p}(\mathbb{F}_{p^d}) \cong C_d$. The generator is Frobenius function $x \mapsto x^p$.

Nov. 2, 2025

1. There are countably many jump discontinuities.
2. Set of Critical Value is zero set.
3. The automorphism $\phi \in \text{Aut}_k(k(x))$ must be $\phi : x \mapsto \frac{ax+b}{cx+d}$ where $ac-bd \neq 0$. A surprising result but hard to prove. Is there a natural but easy way to see this?
4. Review H104.

Nov. 3, 2025

1. $k \subseteq F$ iff there are finitely many fields E such that $k \subseteq E \subseteq F$. This is highly nontrivial to me. My intuition is that if $k(a, b)$ is not a simple extension of k , then it may have a lot of possible combination of a and b to form an intermediate field E . This turns out to be helpful for one direction.
2. All finite separable extensions are simple. This is very surprising and makes our life substantially easier. Consequently, it's very rare to see a nonsimple extension. The proof is very hard, and I find it hard to come up with.

Nov. 4, 2025

1. Set up Wolfram.
2. Wedderburn Little Theorem: all finite division rings are fields. The proof by Ernst Witt is incredibly clever. It uses class formula of group and the cyclotomic polynomial.
3. A Riemann integrable function has integral 0 iff it's zero almost everywhere (set of roots is a zero set).
4. Antiderivative Theorem: The antiderivative of a Riemann integrable function, if it exists, differs from the indefinite integral by a constant. This can be proved using Mean Value Theorem.

Nov. 5, 2025

1. Zsigmondy's Theorem: For $a > b > 0$ coprime integers, $a^n - b^n$ has a prime factor which does not divide $a^k - b^k$ for any $k < n$, with exceptions $(a, b, n) = (2, 1, 6)$ or $n = 2$ and $a + b$ is a power of 2. The proof used LET lemma and properties of cyclotomic polynomial. Very clever proof.
2. There are amazing interactions with cyclotomic polynomials and order of an element a in the multiplicative group mod p when $p \nmid n$: $p \mid \Phi_n(a)$ iff the order of $a \bmod p$ is n . Thus, $n \mid p - 1$ by Fermat's Little Theorem. With this, we can show that there are infinitely many primes congruent to 1 mod n by considering $\Phi_n(k)$ for some integer k .

Nov. 6, 2025

1. Took 250 midterm. All questions are trivial.

2. Went to an AYCE Sushi at San Francisco.

3. Learn Series in H104, easy.

Nov. 7, 2025

1. Fundamental Theorem of Galois Theory: Let $k \subseteq F$ be a finite Galois extension. The lattice of field extensions between k and F is anti-isomorphic to the lattice of subgroups of $Gal(F/k)$. This is astonishing. The proof is quite straightforward, but I don't know how Galois came up with it. Too deep.
2. Mrs. Xinyang Qian gives a guest lecture on the examples of a derivative of a function not being Riemann integrable using Cantor fat set and topologist sine curve. Very interesting examples.
3. Prove that if f_n is a sequence of continuous functions converging uniformly to f , then f is continuous.