DRUG 1	DRUG 2	EFFECT	OUTCOME	TYPE
atropine	Levodopa	↓GI motility ↑ levodopa absorption	↑ levodopa effect	
Adrenaline	Lignocaine	↓ systemic absorption of lignocaine	↑ prolonged action	
Diuretics	NSAIDS	NSAID ↓ PG → ↓ Renal Blood Flow	↓ Diuretic effect	
Allopurinol	Azathioprine	Allopurinol ⊖ Xanthine oxidase	↑ Azathioprine toxicity	
Theophylline	Phenytoin	Phenytoin induces liver enzymes	↓ Theophylline levels	
Proton Pump inhibitors	L – thyroxine	↑ gastric pH → ↓ thyroxine absorption	↓ Thyroxine effect	
Digitalis	Thiazides	Thiazides cause hypokalemia	↑ Digitalis Toxicity	
Captopril	Potassium sparing diuretics	Both ↑ potassium	Hyperkalemia	
Warfarin	NSAIDs	NSAIDs inhibit platelets + gastric mucosal damage	↑ Bleeding risk	

Insulin — Probenecid — Probenecid may ↓ renal clearance of insulin — ↑ Hypoglycemia risk.

DRUG 1	DRUG 2	EFFECT	OUTCOME	TYPE
Tertacyclin	Iron preparation	Chelation in GIT ↓ absorpt of Tetracycline	↓ Antibiotic efficacy	
Levothyroxine	Iron preparation	Iron Binds thyroxine - ↓ absorpt	Hypothyroidism ↓ effects	
levodopa	Pyridoxine (Vitamin B_6)	↑ peripheral decarboxylation of levodopa	↓ Antiparkinson effect	
Levodopa	Antipsychotics	Dopamine receptor Blocker	Antagonizes levodopa effect	
Sildenafil	Nitrates	↑ cGMP → marked vasodilation	severe hypotension	
Metronidazole	Alcohol	Disulfiram like reaction (↓ ALDH)	flushing, vomiting, tachycardia	
Grapefruit juice	Simvastatin	CYP3A4 inhibition → ↓ metabolism	↑ statin toxicity (myopathy)	
Insulin	Beta blockers	Mask hypoglycemic + inhibits glycogenolysis	severe undetected hypoglycemia.	