1

DISCRETE

EE23BTECH11006 - Ameen Aazam*

Question: Show that the products of the corresponding terms of the sequences $a, ar, ar^2, \dots ar^{n-1}$ and $A, AR, AR^2, \dots AR^{n-1}$ form a G.P., and find the common ratio.

Solution: General term of the n^{th} term of the 1^{st}

Input Parameters	Values	Description
а		First term of 1 st G.P.
r		Common ratio of 1 st G.P.
$x_1(n)$	$x_1(n) = ar^n u(n)$	General term of 1 st G.P.
$X_1(z)$		z-Transform of 1 st G.P.
A		First term of 2 nd G.P.
R		Common ratio of 2^{nd} G.P.
$x_2(n)$	$x_1(n) = AR^n u(n)$	General term of 2^{nd} G.P.
$X_{2}\left(z\right)$		z-Transform of 2 nd G.P.

TABLE 0 Parameters

G.P.,

$$x_1(n) = ar^n u(n) \tag{1}$$

Now the sequence in the z domain would be,

$$X_{1}(z) = \sum_{n=-\infty}^{\infty} ar^{n} u(n) z^{-n}$$
 (2)

$$= \frac{a}{1 - rz^{-1}}, \quad |z| > r \tag{3}$$

General for the 2^{nd} G.P. is given as,

$$x_2(n) = AR^n u(n) \tag{4}$$

And the z-Transform,

$$X_{2}(z) = \sum_{n=-\infty}^{\infty} AR^{n}u(n)z^{-n}$$
 (5)

$$= \frac{A}{1 - Rz^{-1}}, \quad |z| > R \tag{6}$$

Now taking the product will result in a sequence as,

$$y(n) = x_1(n) x_2(n) (7)$$

$$= aA (rR)^n u(n)$$
 (8)

z-Transform of the resulting sequence,

$$Y(z) = \sum_{n=-\infty}^{\infty} aA (rR)^n u(n) z^{-n}$$
 (9)

$$=\frac{aA}{1-rRz^{-1}}, \quad |z| > rR \tag{10}$$

So, form 8, taking the ratio of two consecutive terms,

$$\frac{y(n)}{y(n-1)} = \frac{aA(rR)^n u(n)}{aA(rR)^{n-1} u(n-1)}$$
(11)

$$= rR \tag{12}$$

As we can see the ratio of any two consecutive terms, rR, is a constant. Which means the product of the corresponding terms of the two G.P.s results in another G.P. And the common ratio is rR.