

SETSQUARE ACADEMY

Degree Engineering (MU)

COMPUTER ENGINEERING

S.E. SEMESTER III

Question Paper Set

(December 2013 - May 2018)

CONTENTS

Sr. No.	Subject	Page No.
1.	Applied Mathematics III	2 – 15
2.	Digital Logic Design And Analysis	
3.	Discrete Mathematics (Discrete Structures)	
4.	Electronic Circuits & Communication Fundamentals	
5.	Data Structures	
6.	Object Oriented Programming Methodology (OLD)	

As per the Revised Syllabus effective from Academic Year 2017–18

To Download Original Question Paper Set Visit:

www.setsquareacademy.com

APPLIED MATHEMATICS - III

DECEMBER 2013

(REVISED COURSE)

(3 Hours)

QP Code : GX-12040

[Total Marks : 80]

Con. 7854-13

- N.B.: (1) Question No. 1 is compulsory.
 (2) Answer any three questions from Q. 2 to Q. 6.
 (3) Each question carry equal marks.
 (4) Non-programmable calculator is allowed.

1. (a) Find $L^{-1} \left\{ \frac{e^{4-3s}}{(s+4)^{5/2}} \right\}$ 5
 (b) Find the constant a, b, c, d and e If $f(z) = (ax^4 + bx^2y^2 + cy^4 + dx^2 - 2y^2) + i(4x^3y - exy^3 + 4xy)$ is analytic. 5
 (c) Obtain half range Fourier cosine series for $f(x) = \sin x$, $x \in (0, \pi)$. 5
 (d) If \mathbf{r} and $\bar{\mathbf{r}}$ have their usual meaning and a is constant vector, prove that $\nabla \times \left[\frac{\mathbf{a} \times \bar{\mathbf{r}}}{r^n} \right] = \frac{(2-n)}{r^n} \mathbf{a} + \frac{n(\mathbf{a} \cdot \bar{\mathbf{r}})\bar{\mathbf{r}}}{r^{n+2}}$ 5
2. (a) Find the analytic function $f(c) = u + iv$, If $3u + 2v = y^2 - x^2 + 16xy$. 6
 (b) Find the z - transform of $\{a^{|k|}\}$ and hence find the z-transform of $\left\{ \left(\frac{1}{2} \right)^{|k|} \right\}$ 6
 (c) Obtain Fourier series expansion for $f(x) = \sqrt{1 - \cos x}$, $x \in (0, 2\pi)$ and hence deduce that $\sum_{n=1}^{\infty} \frac{1}{4n^2 - 1} = \frac{1}{2}$ 8
3. (a) Find :- (i) $L^{-1} \left\{ \frac{s}{(2s+1)^2} \right\}$ (ii) $L^{-1} \left\{ \log \frac{s^2 + a^2}{\sqrt{s+b}} \right\}$ 6
 (b) Find the orthogonal trajectories of the family of curves $e^x \cos y + xy = \infty$ where ∞ is the real constant in xy - plane. 6
 (c) Show that $\bar{\mathbf{F}} = (y e^{xy} \cos z)\mathbf{i} + (x e^{xy} \cos z)\mathbf{j} - (e^{xy} \sin z)\mathbf{k}$ is irrotational and find the scalar potential for $\bar{\mathbf{F}}$ and evaluate $\int_c \bar{\mathbf{F}} \cdot d\mathbf{r}$ along the curve joining the points (0, 0, 0) and $(-1, 2, \pi)$. 8
4. (a) Evaluate by Green's theorem. $\int e^{-x} \sin y \, dx + e^{-x} \cos y \, dy$ where c is the rectangle whose vertices are $(0, 0)$, $(\pi, 0)$, $\left(\pi, \frac{\pi}{2}\right)$ and $\left(0, \frac{\pi}{2}\right)$. 6
 (b) Find the half range sine series for the function. $f(x) = \frac{2kx}{\ell}$, $0 \leq x \leq \frac{\ell}{2}$ 6

$$= \frac{2k}{\ell}(\ell - x), \quad \frac{\ell}{2} \leq x \leq \ell$$

- (c) Find the inverse z-transform of $\frac{1}{(z-3)(z-2)}$ 8
- (i) $|z| < 2$ (ii) $2 < |z| < 3$ (iii) $|z| > 3$.
5. (a) Solve using Laplace transform. $\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 3y = e^{-x}$, $y(0) = 1$, $y'(0) = 1$. 6
- (b) Express $f(x) = \frac{\pi}{2} e^{-x} \cos x$ for $x > 0$ as Fourier sine integral and show that 6
- $$\int_0^{\infty} \frac{w^3 \sin wx}{w^4 + 4} dw = \frac{\pi}{2} e^{-x} \cos x$$
- (c) Evaluate $\iint_s \vec{F} \cdot \vec{n} ds$, where $\vec{F} = x\vec{i} - y\vec{j} + (z^2 - 1)\vec{k}$ and s is the cylinder formed by the surface $z = 0$, $z = 1$, $x^2 + y^2 = 4$, using the Gauss - Divergence theorem. 8
6. (a) Find the inverse Laplace transform by using convolution theorem $L^{-1} \left\{ \frac{s^2 + 2s + 3}{(s^2 + 2s + 5)(s^2 + 2s + 2)} \right\}$ 6
- (b) Find the directional derivative of $\phi = 4e^{2x-y+z}$ at the point $(1, 1, -1)$ in the direction towards the point $(-3, 5, 6)$. 6
- (c) Find the image of the circle $x^2 + y^2 = 1$, under the transformation $w = \frac{5-4z}{4z-2}$ 8

MAY 2014

(REVISED COURSE)

(3 Hours)

Con.9833-14

QP Code : NP-18619

[Total Marks : 80]

- N.B.: (1) Question No. 1 is compulsory.
 (2) Attempt any three questions from Question No.2 to Question No.6
 (3) Non-programmable calculator is allowed.

1. (a) Find $L^{-1} \left[\frac{se^{-\pi s}}{s^2 + 2s + 2} \right]$ 5
- (b) State true or false with proper justification "There does not exist an analytic function whose real part is $x^3 - 3x^2y - y^3$ ". 5
- (c) Prove that $f_1(x) = 1$, $f_2(x) = x$, $f_3(x) = \frac{(3x^2 - 1)}{2}$ are orthogonal over $(-1, 1)$. 5
- (d) Using Green's theorem in the plane, evaluate $\int_C (x^2 - y) dx + (2y^2 + x) dy$ around the boundary of the region defined by $y = x^2$ and $y = 4$. 5
2. (a) Find the fourier cosine integral representation of the function $f(x) = e^{-ax}$, $x > 0$ and hence show that 6
- $$\int_0^{\infty} \frac{\cos ws}{1 + w^2} dw = \frac{\pi}{2} e^{-x}, x \geq 0.$$

- (b) Verify Laplace's equation for $U = \left(r + \frac{a^2}{r}\right) \cos \theta$. Also find V and $f(z)$ 6
- (c) Solve the following eqn. by using Laplace transform $\frac{dy}{dt} + 2y + \int_0^t y dt = \sin t$ given that $y(0) = 1$. 8
3. (a) Expand $f(x) = \begin{cases} \pi x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$ with period 2 into a Fourier series. 6
- (b) A vector field is given by $\vec{F} = (x^2 + xy^2)\mathbf{i} + (y^2 + x^2y)\mathbf{j}$ show that \vec{F} is irrotational and find its scalar potential 6
- (c) Find the inverse z -transform of $f(z) = \frac{z+2}{z^2-2z+1}$, $|z| > 1$ 8
4. (a) Find the constants 'a' and 'b' so that the surface $ax^2 - byz = (a+2)x$ will be orthogonal to the surface $4x^2y + z^3 = 4$ at $(1, -1, 2)$ 6
- (b) Given $L(\operatorname{erf} \sqrt{t}) = \frac{1}{S\sqrt{S+1}}$, evaluate $\int_0^\infty t e^{-t} \operatorname{erf}(\sqrt{t}) dt$ 6
- (c) Obtain the expansion of $f(x) = x(\pi - x)$, $0 < x < \pi$ as a half-range cosine series. 8
- Hence show that - (i) $\sum_{n=1}^\infty \frac{(-1)^{n+1}}{n^2} = \frac{\pi^2}{12}$ (ii) $\sum_{n=1}^\infty \frac{1}{n^4} = \frac{\pi^4}{90}$
5. (a) If the imaginary part of the analytic function $W = f(z)$ is $V = x^2 - y^2 + \frac{x}{x^2 + y^2}$ find the real part U . 6
- (b) If $f(k) = 4^k U(k)$ and $g(k) = 5^k U(k)$, then find the z -transform of $f(k) \cdot g(k)$ 6
- (c) Use Gauss's Divergence theorem to evaluate $\iint_S \vec{N} \cdot \vec{F} ds$ where $\vec{F} = 4x\mathbf{i} + 3y\mathbf{j} - 2z\mathbf{k}$ and S is the surface bounded by $x = 0$, $y = 0$, $z = 0$ and $2x + 2y + z = 4$. 8
6. (a) Obtain complex form of Fourier series for $f(x) = \cos h 3x + \sin h 3x$ in $(-3, 3)$. 6
- (b) Find the inverse Laplace transform of $L^{-1} \frac{(s-1)^2}{(s^2-2s+5)^2}$ 6
- (c) Find the bilinear transformation under which $1, i, -1$ from the z -plane are mapped onto $0, 1, \infty$ of w -plane. Also show that under this transformation the unit circle in the w -plane is mapped onto a straight line in the z -plane. Write the name of this line. 8

DECEMBER 2014

(REVISED COURSE)

(3 Hours)

GN.Con.6452-14

QP Code : 14544

[Total Marks : 80]

- N.B.: (1) Question no. 1 is compulsory.
 (2) Attempt any three from the remaining.
 (3) Figures to the right indicate full marks.

1. (a) Find the Laplace Transform of $\sin t \cos 2t \cos t$. 5
 (b) Find the Fourier series expansion of $f(x) = x^2$ $(-\pi, \pi)$ 5

- (c) Find the z-transform of $\left(\frac{1}{3}\right)^{|k|}$ 5
- (d) Find the directional derivative of $4xz^2 + x^2yz$ at $(1, -2, -1)$ in the direction of $2\bar{i} - \bar{j} - 2\bar{k}$ 5
2. (a) Find an analytic function $f(z)$ whose real part is $e^x(x \cos y - y \sin y)$ 6
- (b) Find inverse Laplace Transform by using convolution theorem, $\frac{1}{(s-3)(s+4)^2}$ 6
- (c) Prove that $\bar{F} = (6xy^2 - 2z^3)\bar{i} + (6x^2y + 2yz)\bar{j} + (y^2 - 6z^2x)\bar{k}$ is a conservative field. Find the scalar potential Φ such that $\nabla\Phi = \bar{F}$. Hence find the work done by \bar{F} in displacing a particle from $A(1, 0, 2)$ to $B(0, 1, 1)$ along AB . 8
3. (a) Find the inverse z-transform of $f(z) = \frac{z^3}{(z-3)(z-2)^2}$ 6
- (i) $2 < |z| < 3$ (ii) $|z| > 3$
- (b) Find the image of the real axis under the transformation $w = \frac{2}{z+i}$ 6
- (c) Obtain the Fourier series expansion of $f(x) = \pi x; 0 \leq x \leq 1$ 8
- $= \pi(2-x); 1 \leq x \leq 2$
- Here deduce That $\frac{1}{1^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{8}$
4. (a) Find the Laplace Transform of $f(t) = E; 0 \leq t \leq p/2$ 6
- $f(t+p) = f(t)$
 $= -E; p/2 \leq t \leq p,$
- (b) Using Green's theorem evaluate $\int_c \frac{1}{y} dx + \frac{1}{x} dy$ where c is the boundary of the region bounded by 6
- $x=1, x=4, y=1, y=\sqrt{x}$
- (c) Find the Fourier integral for $f(x) = 1 - x^2, 0 \leq x \leq 1$ 8
- $= 0 \quad x > 1$
- Hence evaluate $\int_0^\infty \frac{\lambda \cos \lambda - \sin \lambda}{\lambda^3} \cos\left(\frac{\lambda}{2}\right) d\lambda$
5. (a) If $\bar{F} = x^2\bar{i} + (x-y)\bar{j} + (y+z)\bar{k}$ moves a particle from $A(1, 0, 1)$ to $B(2, 1, 2)$ along line AB . Find the work done. 6
- (b) Find the complex form of fourier series $f(x) = \sinh ax (-\ell, \ell)$ 6
- (c) Solve the differential equation using Laplace Transform. $(D^2 + 2D + 5)y = e^{-t} \sin t, y(0) = 0, y'(0) = 1$ 8
6. (a) If $\int_0^\infty e^{-2t} \sin(t+\alpha) \cos(t-\alpha) dt = \frac{3}{8}$ find the value of α . 6

- (b) Evaluate $\iint_s (y^2 z^2 \bar{i} + z^2 x^2 \bar{j} + z^2 y^2 \bar{k}) \cdot \bar{n} \, ds$ where s is the hemisphere $x^2 + y^2 + z^2 = 1$ above xy -plane and bounded by this plane. 6
- (c) Find Half range sine series for $f(x) = \ell x - x^2$ ($0, \ell$). Hence prove that $\frac{1}{1^6} + \frac{1}{3^6} + \dots = \frac{\pi^6}{960}$ 8

MAY 2015

(REVISED COURSE)

(3 Hours)

JP-Con.8899-15

QP Code : 4827

[Total Marks : 80]

- N.B.: (1) Question No.1 is compulsory.
 (2) Attempt any three from the remaining six questions.
 (3) Figures to the right indicate full marks.

1. (a) Find Laplace Transform of $\frac{\sin t}{t}$ 20
 (b) Prove that $f(z) = \sinh z$ is analytic and find its derivative
 (c) Find Fourier Series for $f(x) = 9 - x^2$ over $(-3, 3)$
 (d) Find $Z[f(k)*g(k)]$ if $f(k) = \frac{1}{3^k}$, $g(k) = \frac{1}{5^k}$
2. (a) Prove that $\bar{F} = ye^{xy} \cos z \bar{i} + xe^{xy} \cos z \bar{j} - e^{xy} \sin z \bar{k}$ is irrotational. Find Scalar Potential for \bar{F} 6
 Hence evaluate $\int_C \bar{F} \cdot d\bar{r}$ along the curve C joining the points $(0, 0, 0)$ and $(-1, 2, \pi)$
 (b) Find the Fourier series for $f(x) = \frac{\pi - x}{2}$, $0 \leq x \leq 2\pi$ 6
 (c) Find Inverse Laplace Transform of (i) $\frac{s+29}{(s+4)(s^2+9)}$ (ii) $\frac{e^{-2s}}{s^2+8s+25}$ 8
3. (a) Find the Analytic function $f(z) = u + iv$ if $u + v = \frac{x}{x^2 + y^2}$ 6
 (b) Find Inverse Z transform of $\frac{1}{(z-1/2)(z-1/3)}$, $1/3 < |z| < 1/2$ 6
 (c) Solve the Differential Equation $\frac{d^2 y}{dt^2} + y = t$, $y(0) = 1$, $y'(0) = 0$, using Laplace Transform 8
4. (a) Find the Orthogonal Trajectory of $3x^2y - y^3 = k$ 6
 (b) Using Greens theorem evaluate $\int_C (xy + y^2) dx + x^2 dy$, C is closed path formed by $y = x$, $y = x^2$ 6
 (c) Find Fourier Integral of $f(x) = \begin{cases} \sin x & 0 \leq x \leq \pi \\ 0 & x > \pi \end{cases}$. Hence show that $\int_0^\infty \frac{\cos(\lambda \pi / 2)}{1 - \lambda^2} d\lambda = \frac{\pi}{2}$ 8

5. (a) Find Inverse Laplace Transform using Convolution theorem $\frac{s}{(s^4 + 8s^2 + 16)}$ 6
- (b) Find the Bilinear Transformation that maps the points $z = 1, i, -1$ into $w = i, 0, -i$. 6
- (c) Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where C is the boundary of the plane $2x + y + z = 2$ cut off by co-ordinate planes 8
and $\vec{F} = (x + y)\mathbf{i} + (y + z)\mathbf{j} - x\mathbf{k}$.
6. (a) Find the Directional derivative of $\phi = x^2 + y^2 + z^2$ in the direction of the line $\frac{x}{3} = \frac{y}{4} = \frac{z}{5}$ at $(1, 2, 3)$ 6
- (b) Find Complex Form of Fourier Series for $e^{2x}; 0 < x < 2$ 6
- (c) Find Half Range Cosine Series for $f(x) = \begin{cases} kx; & 0 \leq x \leq \ell/2 \\ k(\ell - x); & \ell/2 \leq x \leq \ell \end{cases}$ hence find $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$ 8

DECEMBER 2015

MD-Con.7526-15

(REVISED COURSE)

QP Code : 5067

(3 Hours)

[Total Marks : 80]

- Instructions: (1) Question No. 1 is compulsory.
(2) Attempt any THREE of the remaining.
(3) Figures to the right indicate full marks.

1. (a) Find Laplace of $\{t^5 \cosht\}$ 5
- (b) Find Fourier series for $f(x) = 1 - x^2$ in $(-1, 1)$ 5
- (c) Find a, b, c, d, e if, $f(z) = (ax^4 + bx^2y^2 + cy^4 + dx^2 - 2y^2) + i(4x^3y - exy^3 + 4xy)$ is analytic 5
- (d) Prove that $\nabla \left(\frac{1}{r} \right) = -\frac{\mathbf{r}}{r^3}$ 5
2. (a) If $f(z) = u + iv$ is analytic and $u + v = \frac{3 \sin 2x}{e^{2y} + e^{-2y} - 2 \cos 2x}$, find $f(z)$ 6
- (b) Find inverse Z-transform of $f(z) = \frac{z + 2}{z^2 - 2z + 1}$ for $|z| > 1$ 6
- (c) Find Fourier series for $f(x) = \sqrt{1 - \cos x}$ in $(0, 2\pi)$ Hence, deduce that $\frac{1}{2} = \sum_{n=1}^{\infty} \frac{1}{4n^2 - 1}$ 8
3. (a) Find $L^{-1} \left\{ \frac{1}{(s-2) + (s+3)} \right\}$ Using Convolution theorem 6
- (b) Prove that $f_1(x) = 1, f_2(x) = x, f_3(x) = (3x^2 - 1)/2$ are orthogonal over $(-1, 1)$ 6
- (c) Verify Green's theorem for $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F} = (x^2 - y^2)\mathbf{i} + (x + y)\mathbf{j}$ and c is the triangle with vertices $(0, 0), (1, 1), (2, 1)$ 8
4. (a) Find Laplace Transform of $f(t) = |\sin t|, t \geq 0$ 6

(b) Show that $\vec{F} = (y \sin z - \sin x) \mathbf{i} + (x \sin z + 2yz) \mathbf{j} + (xy \cos z + y^2) \mathbf{k}$ is irrotational
Hence, find its scalar potential. 6

(c) Obtain Fourier expansion of $f(x) = x + \frac{\pi}{2}$ where $-\pi < x < 0$ 8
 $= \frac{\pi}{2} - x$ where $0 < x < \pi$

Hence, deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$ (ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

5. (a) Using Gauss Divergence theorem to evaluate $\iint_S \vec{N} \cdot \vec{F} ds$ where $\vec{F} = 4x\mathbf{i} - 2y^2\mathbf{j} + z^2\mathbf{k}$ and S is the 6

region bounded by $x^2 + y^2 = 4$, $z = 0$, $z = 3$.

(b) Find $Z[2^k \cos(3k+2)]$, $k \geq 0$ 6

(c) Solve $(D^2 + 2D + 5)y = e^{-t} \sin t$, with $y(0) = 0$ and $y'(0) = 1$ 8

6. (a) Find $L^{-1} \left\{ \tan^{-1} \left(\frac{z}{s^2} \right) \right\}$ 6

(b) Find the bilinear transformation which maps the points 2, i, -2 onto points 1, i, -1 by using cross-ratio property. 6

(c) Find Fourier Sine integral representation for $f(x) = \frac{e^{-ax}}{x}$ 8

MAY 2016

(REVISED COURSE)

(3 Hours)

FW-Con.9413-16

QP Code : 30557

[Total Marks : 80]

- N.B.: (1) Question No. 1 is compulsory.
 (2) Attempt any THREE of the remaining.
 (3) Figures to the right indicate full marks.

1. (a) If $\int_0^\infty e^{-2t} \sin(t+\alpha) \cos(t-\alpha) dt = \frac{1}{4}$ find α . 5

(b) Find half range Fourier cosine series for $f(x) = x$, $0 < x < 2$ 5

(c) If $u(x, y)$ is a harmonic function then prove that $f(z) = u_x - iu_y$ is an analytic function. 5

(d) Prove that $\nabla f(r) = f'(r) \frac{\vec{r}}{r}$ 5

2. (a) If $v = e^x \sin y$, prove that v is a harmonic function. Also find the corresponding analytic function. 6

(b) Find Z-transform of $f(k) = b^k$, $k \geq 0$ 6

(c) Obtain Fourier series for $f(x) = \frac{3x^2 - 6x\pi + 2\pi^2}{12}$ in $(0, 2\pi)$, where $f(x + 2\pi) = f(x)$. 8

Hence deduce that $\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots$

3. (a) Find inverse Laplace of $\frac{(s+3)^2}{(s^2+6s+5)^2}$ using Convolution theorem 6
- (b) Show that the set of functions $\{\sin x, \sin 3x, \sin 5x, \dots\}$ is orthogonal over $[0, \pi/2]$. Hence construct orthonormal set of functions. 6
- (c) Verify Green's theorem for $\int_C \frac{1}{y} dx + \frac{1}{x} dy$ where C is the boundary of region defined by $x=1, x=4, y=1$ and $y=\sqrt{x}$ 8
4. (a) Find $Z\{k^2 a^{k-1} U(k-1)\}$ 6
- (b) Show that the map of the real axis of the z -plane is a circle under the transformation $w = \frac{2}{z+i}$. Find its centre and the radius. 6
- (c) Express the function $f(x) = \begin{cases} \sin x & |x| < \pi \\ 0 & |x| > \pi \end{cases}$ as Fourier sine Integral. 8
5. (a) Using Gauss Divergence theorem evaluate $\iiint_S \vec{N} \cdot \vec{F} ds$ where $\vec{F} = x^2 \vec{i} + z \vec{j} + yz \vec{k}$ and S is the cube bounded by $x=0, x=1, y=0, y=1, z=0, z=1$ 6
- (b) Find inverse Z-transform of $F(z) = \frac{z}{(z-1)(z-2)}, |z| > 2$ 6
- (c) Solve $(D^2 + 3D + 2)y = e^{-2t} \sin t$, with $y(0) = 0$ and $y'(0) = 0$ 8
6. (a) Find Fourier expansion of $f(x) = 4 - x^2$ in the interval $(0, 2)$ 6
- (b) A vector field is given by $\vec{F} = (x^2 + xy^2) \vec{i} + (y^2 + x^2y) \vec{j}$. Show that \vec{F} is irrotational and find its scalar potential. 6
- (c) Find (i) $L^{-1} \left\{ \tan^{-1} \left(\frac{a}{s} \right) \right\}$ (ii) $L^{-1} \left(\frac{e^{-\pi s}}{s^2 - 2s + 2} \right)$ 8

DECEMBER 2016

(REVISED COURSE)

(3 Hours)

QP Code : 540701

[Total Marks : 80]

- N.B.: (1) Question No. 1 is compulsory.
 (2) Attempt any three of the remaining.
 (3) Figures to the right indicate full marks.

1. (a) Find the Laplace transform of $te^{3t} \sin 4t$. 5
- (b) Find half-range cosine series for $f(x) = e^x, 0 < x < 1$. 5
- (c) Is $f(z) = \frac{z}{z}$ analytic? 5
- (d) Prove that $\nabla \times (\vec{a} \times \nabla \log r) = 2 \frac{(\vec{a} \cdot \vec{r}) \vec{r}}{r^4}$, where \vec{a} is a constant vector. 5

2. (a) Find the inverse Z-transform of $\frac{1}{(z-5)^3}$ if $|z| < 5$ 6
- (b) If $V = 3x^2y + 6xy - y^3$, show that V is harmonic & find the corresponding analytic function. 6
- (c) Obtain Fourier series for the function $f(x) = \begin{cases} 1 + \frac{2x}{\pi}, & -\pi \leq x \leq 0 \\ 1 - \frac{2x}{\pi}, & 0 \leq x \leq \pi \end{cases}$ 8
- hence deduce that $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$
3. (a) Find $L^{-1} \left[\frac{(s+2)^2}{(s^2+4s+8)^2} \right]$ using convolution theorem. 6
- (b) Show that the set of functions $1, \sin\left(\frac{\pi x}{L}\right), \cos\left(\frac{\pi x}{L}\right), \sin\left(\frac{2\pi x}{L}\right), \cos\left(\frac{2\pi x}{L}\right), \dots$ 6
- Form an orthogonal set in $(-L, L)$ and construct an orthonormal set.
- (c) Verify Green's theorem for $\int_C \{e^{2x} - xy^2\} dx + (ye^x + y^2) dy$ Where C is the closed curve bounded by 8
- $y^2 = x$ & $x^2 = y$.
4. (a) Find Laplace transform of $f(x) = K \frac{t}{T}$ for $0 < t < T$ & $f(t) = f(t+T)$. 6
- (b) Show that the vector, $\vec{F} = (x^2 - yz)\mathbf{i} + (y^2 - zx)\mathbf{j} + (z^2 - xy)\mathbf{k}$ is irrotational and hence, find ϕ such 6
- that $\vec{F} = \nabla \phi$
- (c) Find Fourier series for $f(x)$ in $(0, 2\pi)$, $f(x) = \begin{cases} x, & 0 \leq x \leq \pi \\ 2\pi - x, & \pi \leq x \leq 2\pi \end{cases}$ hence deduce that 8
- $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$
5. (a) Use Gauss's Divergence theorem to evaluate $\iint_s \vec{N} \cdot \vec{F} ds$ where $\vec{F} = 2x\mathbf{i} + xy\mathbf{j} + z\mathbf{k}$ over the region 6
- bounded by the cylinder $x^2 + y^2 = 4$, $z = 0$, $z = 6$.
- (b) Find inverse Z - transform of $f(x) = \frac{z}{(z-1)(z-2)}$, $|z| > 2$ 6
- (c) (i) Find $L^{-1} \left[\log \left(\frac{s+1}{s-1} \right) \right]$ (ii) $L^{-1} \left[\frac{s+2}{s^2-4s+13} \right]$ 8
6. (a) Solve $(D^2 + 3D + 2)y = 2(t^2 + t + 1)$ with $y(0) = 2$ & $y'(0) = 0$. 6
- (b) Find the bilinear transformation which maps the points $0, i, -2i$ of z-plane onto the points $-4i, \infty, 0$ 6
- respectively of w-plane. Also obtain fixed points of the transformation.

- (c) Find Fourier sine integral of $f(x) = \begin{cases} x, & 0 < x < 1 \\ 2-x, & 1 < x < 2 \\ 0, & x > 2 \end{cases}$ 8

MAY 2017
(REVISED COURSE)
(3 Hours)

QP Code : 540702
[Total Marks : 80]

- N.B.: (1) Question No. 1 is compulsory
(2) Attempt any three from the remaining six questions
(3) Figures to the right indicate full marks

1. (a) Find the Laplace Transform of $e^{-t} \int_0^t u \cos 2u \, du$ 20
(b) Prove that $f(z) = \sinh z$ is analytic and find its derivative
(c) Obtain Half range Sine Series for $f(x) = x + 1$ in $(0, \pi)$
(d) Find a unit vector normal to the surface $x^2y + 2xz = 4$ at $(2, -2, 3)$
2. (a) Prove that $\vec{F} = (2xy^2 + yz)\mathbf{i} + (2x^2y + xz + 2yz^2)\mathbf{j} - (2y^2z + xy)\mathbf{k}$ is Irrotational. Find Scalar Potential for \vec{F} 6
(b) Find the inverse Laplace Transform using Convolution theorem $\frac{(s-1)^2}{(s^2-2s+5)^2}$ 6
(c) Find Fourier Series of $f(x) = \begin{cases} \pi x; & 0 \leq x \leq 1 \\ \pi(2-x); & 1 \leq x \leq 2 \end{cases}$ 8
3. (a) Find the Analytic function $f(z) = u + iv$ if $v = \frac{x}{x^2+y^2} + \cosh x \cos y$ 6
(b) Find Inverse Z transform of $\frac{(3z^2 - 18z + 26)}{(z-2)(z-3)(z-4)}$, $3 < |z| < 4$ 6
(c) Solve the Differential Equation $\frac{d^2y}{dt^2} + 2\frac{dy}{dx} + 2y = 5 \sin t$, $y(0) = 0$, $y'(0) = 0$ using Laplace Transform 8
4. (a) Find the Orthogonal Trajectory of $3x^2y - y^3 = k$ 6
(b) Find the z-transform of $2^k \sinh 3K$, $K \geq 0$ 6
(c) Express the function $f(x) = \begin{cases} 1 : & |x| < 1 \\ 0 : & |x| > 1 \end{cases}$ as Fourier Integral. Hence evaluate $\int_0^\infty \frac{\sin \lambda}{\lambda} \cdot \cos(\lambda x) d\lambda$ 8
5. (a) Evaluate using Stoke's theorem $\int_C (2x - y)dx - yz^2dy - y^2zdz$ where C is the circle $x^2 + y^2 = 1$ 6
corresponding to the sphere $x^2 + y^2 + z^2 = 1$ above the XY plane

- (b) Show that $w = \frac{2z+3}{z-4}$ maps the circle $x^2 + y^2 - 4x = 0$ into straight line $4u + 3 = 0$ 6
- (c) Find Inverse Laplace Transform (i) $e^{-s} \tanh^{-1}s$ (ii) $\frac{6}{(2s+1)^3}$ 8
6. (a) Find the Laplace transform of $f(t) = \frac{2t}{3}, 0 \leq t \leq 3, f(t+3) = f(t)$ 6
- (b) Find Complex Form of Fourier Series for $\sin(\alpha x); (-\pi, \pi), \alpha$ is not an integer 6
- (c) Verify Green's theorem for $\int_C (2x^2 - y^2)dx + (x^2 + y^2)dy$ where C is the boundary of the surface 8
enclosed by lines $x = 0, y = 0, x = 2, y = 2$.

APPLIED MATHEMATICS - III

DECEMBER 2017

(REVISED COURSE)

(3 Hours)

QP Code : 24510

[Total Marks : 80]

- N.B.: (1) Question no.1 is compulsory.
 (2) Attempt any three questions from Q.2 to Q.6
 (3) Figures to the right indicate full marks.

1. (a) Find the Laplace transform of $\frac{1}{t} e^{-t} \sin t$ 5
- (b) Find the inverse Laplace transform of $\frac{1}{\sqrt{2s+1}}$ 5
- (c) Show that the function $f(z) = \sinh z$ is analytic and find $f'(z)$ in terms of z . 5
- (d) Find the Fourier series for $f(x) = x$ in $(0, 2\pi)$
2. (a) Use Laplace transform to prove $\int_0^\infty e^{-t} \frac{\sin^2 t}{t} dt = \frac{1}{4} \log 5$ 6
- (b) If $\{f(k)\} = \begin{cases} 4^k, & k < 0 \\ 3^k, & k \geq 0 \end{cases}$ find $Z\{f(k)\}$ 6
- (c) Show that the function $u = \cos x \cosh y$ is a harmonic function. Find its harmonic conjugate and corresponding analytic function 8
3. (a) Find the equation of the line of regression of Y on X for the following data : 6
- | | | | | | | | |
|---|----|----|----|----|----|----|----|
| X | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| Y | 11 | 14 | 14 | 15 | 12 | 17 | 16 |
- (b) Find the bilinear transformation which maps the points $1, -i, 2$ on z -plane onto $0, 2, -i$ respectively of w -plane 6
- (c) Find half range sine series for $f(x) = \begin{cases} x, & 0 < x < \frac{\pi}{2} \\ \pi - x, & \frac{\pi}{2} < x < \pi \end{cases}$. Hence find the sum of $\sum_{(2n-1)}^\infty \frac{1}{n^4}$ 8
4. (a) Find the inverse Laplace transform by using convolution theorem $\frac{1}{(s-a)(s+a)^2}$ 6
- (b) Calculate the coefficient of correlation between X and Y from the following data 6
- | | | | | | | |
|---|---|---|----|----|----|---|
| X | 8 | 8 | 7 | 5 | 6 | 2 |
| Y | 3 | 4 | 10 | 13 | 22 | 8 |

- (c) Find the inverse Z - transform of 8

(i) $\frac{1}{(z-a)^2} \mid z < a$ (ii) $\frac{1}{(z-3)(z-2)} \mid z > 3$

5. (a) Using Laplace transform evaluate $\int_0^\infty e^{-t} (1+2t-t^2+t^3) H(t-1) dt$ 6
 (b) Show that set of functions $\cos x, \cos 2x, \cos 3x \dots$ Is a set orthogonal functions over $[-\pi, \pi]$. Hence construct a set of orthonormal functions. 6
 (c) Solve using Laplace transform
 $(D^3 - 2D^2 + 5D)y = 0$, with $y(0) = 0, y'(0) = 0, y''(0) = 1$

6. (a) Find the complex form of Fourier series for $f(x) = 2x$ in $(0, 2\pi)$ 6
 (b) If $f(z)$ and $\overline{f(z)}$ are both analytic, prove that $f(z)$ is constant 6
 (c) Fit a curve of the form $y = ab^x$ to the following data 8

X	1	2	3	4	5	6
Y	151	100	61	50	20	8

MAY 2018
 (REVISED COURSE)
 (3 Hours)

QP Code : 21236
 [Total Marks : 80]

- N.B.: (1) Q.1 is COMPULSORY.
 (2) Attempt ANY 3 questions from Q.2 to Q.6
 (3) Use of scientific calculators allowed.
 (4) Figures to right indicate marks.

1. (a) Find the Laplace transform of $e^{-2t} t \cos t$ 5
 (b) Find the inverse Laplace transform of $\frac{3s+7}{s^2-2s-3}$ 5
 (c) Determine whether the function $f(z) = (x^3 + 3xy^2 - 3x) + i(3x^2y - y^3 + 3y)$ is analytic and if so find its derivative. 5
 (d) Find the Fourier series for $f(x) = x^2$ in the interval $(-\pi, \pi)$. 5
2. (a) Evaluate $\int_0^\infty \left(\frac{\sin 2t + \sin 3t}{te^t} \right) dt = \frac{3\pi}{4}$ 6
 (b) Find the Z-Transform of $\left\{ \left(\frac{1}{4} \right)^k \right\}$ 6
 (c) Show that the function $v = e^x(x \sin y + y \cos y)$ is a harmonic function. Find its harmonic conjugate and corresponding analytic function. 8
3. (a) From 8 observations the following results were obtained. 6
 $\Sigma x = 59$; $\Sigma y = 40$; $\Sigma x^2 = 524$; $\Sigma y^2 = 256$; $\Sigma xy = 364$.
 Find the equation of the line of regression of x on y and the coefficient of correlation.
 (b) Find the bilinear transformation which maps the points $z = -1, 0, 1$ onto the points $w = -1, -i, 1$. 6

- (c) Obtain half-range sine series for $f(x) = (x-1)^2$ in $0 < x < 1$. Hence find $\sum_{n=1}^{\infty} \frac{1}{n^2}$ 8

4. (a) Find the inverse Laplace Transform by using convolution theorem $\frac{1}{(s^2 + a^2)(s^2 + b^2)}$ 6

- (b) Compute Spearman's Rank correlation coefficient for the following data: 6

X	85	74	85	50	65	78	74	60	74	90
Y	78	91	78	58	60	72	80	55	68	70

- (c) Find the inverse Z-transform for the following: 8

(i) $\frac{1}{(z-5)^2}, |z| < 5$ (ii) $\frac{\pi}{(z-2)(z-3)}, |z| > 3$

5. (a) Using Laplace Transform evaluate $\int_0^{\infty} e^{-t} (1+3t+t^2) H(t-2) dt$ 6

- (b) Prove that $f_1(x) = 1$; $f_2(x) = x$; $f_3(x) = \left(\frac{3x^2-1}{2}\right)$ are orthogonal over $(-1, 1)$. 6

- (c) Solve using Laplace transform $\frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y = 2e^{3x}$, $y = 2$, $y' = 3$ at $x = 0$ 8

6. (a) Find the complex form of Fourier series for $f(x) = e^x$, $(-\pi, \pi)$ 6

- (b) If u, v are harmonic conjugate functions, show that uv is a harmonic function. 6

- (c) Fit a straight line of the form $y = a + bx$ to the following data and estimate the value of y for $x = 3.5$ 8

x	0	1	2	3	4
Y	1	1.8	3.3	4.5	6.3



Procedure to get remaining all subject's printable typed Question Sets

Those students who have already subscribed last semester need not follow 1st and 2nd step but they have to follow 3rd and 4th step in order to get the required question sets.

- 1) First open the link <https://goo.gl/im6SrJ> & register yourself or login if already registered. After log in you will be redirected to the subscription page.
- 2) Pay the one time **all semesters** subscription charges of just **Rs. 100/-** by selecting the convenient payment method on or before **20th July 2018**. This is the last chance for one time subscription after this deadline it will be valid for just current academic year.
- 3) After successful payment you have to E-mail us a clear photocopy of your "Current College ID" from you registered Email ID to qpsets@setsquareacademy.com for verification of your details as we have observed many fake and wrong registrations.
- 4) Along with college ID also specify the semesters for which you want the printable typed question sets. You can get maximum four semester's question sets within a year.

After receiving your Email we will first verify your subscription and registration details. After successful verification we will send you requested printable question paper set PDFs as soon as we finish with updation of latest May 18 question papers.

The printable question paper set PDFs will contain watermark of your details like Name, Email ID and contact number in the same format as that of all pages in this sample PDF. If we find the circulation of same PDFs in either softcopy or printed copy with watermark of your exact details then you have to face a legal action along with permanent cancellation of your registration and subscription. Also you will not receive any notification for updated question paper sets from next semester onwards. In case of circulation we will not entertain any reasons like misplacing or stealing of the documents by someone else without your notice. It's totally your responsibility to keep the documents secured from sharing.

Every semester you have to send us a new requirement for updated printable question paper set PDFs through Email. So accordingly we will send you the printable PDFs well in advance as soon as we finish with updation work.

Please do not reply to this message. For any queries feel free to contact on 9920848746.

Thanking you!

Regards,
Managing Director,
Setsquare Academy Pvt. Ltd.

Address : 2nd Floor, Govind Smruti, Near Jagdish Book Depot, Lohar Ali, Thane (West), 400602.

Web : www.setsquareacademy.com, **Contact :** 9920848746, **E mail :** info@setsquareacademy.com.