

IPS9 in R: Inference for Proportions (Chapter 8)

Bonnie Lin and Nicholas Horton (nhorton@amherst.edu)

July 13, 2018

Introduction and background

These documents are intended to help describe how to undertake analyses introduced as examples in the Ninth Edition of *Introduction to the Practice of Statistics* (2017) by Moore, McCabe, and Craig.

More information about the book can be found [here](#). The data used in these documents can be found under Data Sets in the Student Site. This file as well as the associated R Markdown reproducible analysis source file used to create it can be found at <https://nhorton.people.amherst.edu/ips9/>.

This work leverages initiatives undertaken by Project MOSAIC (<http://www.mosaic-web.org>), an NSF-funded effort to improve the teaching of statistics, calculus, science and computing in the undergraduate curriculum. In particular, we utilize the `mosaic` package, which was written to simplify the use of R for introductory statistics courses. A short summary of the R needed to teach introductory statistics can be found in the mosaic package vignettes (<http://cran.r-project.org/web/packages/mosaic>). A paper describing the mosaic approach was published in the *R Journal*: <https://journal.r-project.org/archive/2017/RJ-2017-024>.

Chapter 8: Inference for Proportions

This file replicates the analyses from Chapter 8: Inference for Proportions.

First, load the packages that will be needed for this document:

```
library(mosaic)
library(readr)
```

Section 8.1: Inference for a single proportion

```
ROBOT <- read_csv("https://nhorton.people.amherst.edu/ips9/data/chapter08/EG08-01ROBOT.csv")
ROBOT
```

```
## # A tibble: 2 x 2
##   Decrease Count
##   <chr>      <int>
## 1 Yes         910
## 2 No         986
```

```
### Example 8.2, page 487
binom.test(910, 910 + 986,
            alternative="two.sided",
            conf.level=0.95)
```

```
##
##
##
## data:  910 out of 910 + 986
## number of successes = 910, number of trials = 1896, p-value =
## 0.08496
## alternative hypothesis: true probability of success is not equal to 0.5
```

```
## 95 percent confidence interval:
## 0.4572519 0.5027258
## sample estimates:
## probability of success
## 0.4799578
```

By default, the `read_csv()` function will output the types of columns, as we see above. To improve readability for future coding, we will suppress the “Parsed with column specification” message by adding `message = FALSE` at the top of the code chunks.

```
SUNBL <- read_csv("https://nhorton.people.amherst.edu/ips9/data/chapter08/EG08-05SUNBL.csv")
### Example 8.5 and 8.6, page 492-493
binom.test(13, 20, alternative = "two.sided")
```

```
##
##
##
## data: 13 out of 20
## number of successes = 13, number of trials = 20, p-value = 0.2632
## alternative hypothesis: true probability of success is not equal to 0.5
## 95 percent confidence interval:
## 0.4078115 0.8460908
## sample estimates:
## probability of success
## 0.65
## XX The p value is not 0.1802 as the book reports
```

```
INSTAG <- read_csv("https://nhorton.people.amherst.edu/ips9/data/chapter08/EG08-11INSTAG.csv")
### Example 8.11, page 507
INSTAG_prop <- INSTAG %>%
  group_by(Sex) %>%
  mutate(Prop_by_Sex = Count/sum(Count)) %>%
  ungroup() %>%
  filter(User == "Yes")
INSTAG_prop
```

```
## # A tibble: 2 x 4
##   Sex    User Count Prop_by_Sex
##   <chr> <chr> <int>      <dbl>
## 1 1Women Yes    328      0.611
## 2 2Men   Yes    234      0.440
```

```
### Example 8.15, page 513
prop.test(x = c(328, 234), n = c(537, 532), correct = FALSE)
```

```
##
## 2-sample test for equality of proportions without continuity
## correction
##
## data: c(328, 234) out of c(537, 532)
## X-squared = 31.323, df = 1, p-value = 2.185e-08
## alternative hypothesis: two.sided
## 95 percent confidence interval:
## 0.1119627 0.2299396
## sample estimates:
## prop 1 prop 2
```

```
## 0.6108007 0.4398496
### Example 8.16, page 515
zstar = qnorm(.975) # for 95% confidence interval
p = 0.5 # planned proportion estimate
moe = 0.1 # margin of error
p*(zstar/moe)^2

## [1] 192.0729
### Example 8.17, page 516
## XX unknown power plot
```

Section 8.2: Comparing two proportions