

IS5 in R: Relationships Between Categorical Variables–Contingency Tables (Chapter 3)

Margaret Chien and Nicholas Horton (nhorton@amherst.edu)

June 16, 2018

Introduction and background

This document is intended to help describe how to undertake analyses introduced as examples in the Fifth Edition of *Intro Stats* (2018) by De Veaux, Velleman, and Bock. More information about the book can be found at http://wps.aw.com/aw_deveaux_stats_series. This file as well as the associated R Markdown reproducible analysis source file used to create it can be found at <http://nhorton.people.amherst.edu/is5>.

This work leverages initiatives undertaken by Project MOSAIC (<http://www.mosaic-web.org>), an NSF-funded effort to improve the teaching of statistics, calculus, science and computing in the undergraduate curriculum. In particular, we utilize the `mosaic` package, which was written to simplify the use of R for introductory statistics courses. A short summary of the R needed to teach introductory statistics can be found in the `mosaic` package vignettes (<http://cran.r-project.org/web/packages/mosaic>). A paper describing the `mosaic` approach was published in the *R Journal*: <https://journal.r-project.org/archive/2017/RJ-2017-024>.

Chapter 3: Relationships Between Categorical Variables–Contingency Tables

Section 3.1: Contingency Tables

```
library(mosaic)
library(readr)
library(janitor)
OKCupid <- read_csv("http://nhorton.people.amherst.edu/is5/data/OKCupid_CatsDogs.csv", skip = 1) %>%
  clean_names()
```

```
## Parsed with column specification:
## cols(
##   CatsDogsBoth = col_character(),
##   Gender = col_character(),
##   `drugsY/N` = col_character(),
##   `smokesY/N` = col_character()
## )
```

```
# If you look at the data set, you'll see that the first line is "Col1", "Col2", etc
tally(~ cats_dogs_both + gender, margin = TRUE, data = OKCupid)
```

```
##           gender
## cats_dogs_both  F    M Total
##   Has Both    897   577 1474
##   Has cats   3412  2388 5800
##   Has dogs   3431  3587 7018
##   <NA>      16377 29274 45651
##   Total     24117 35826 59943
```

```
tally(~ cats_dogs_both + gender, format = "percent", margin = TRUE, data = OKCupid)
```

```
##                gender
## cats_dogs_both      F      M      Total
##      Has Both  1.4964216  0.9625811  2.4590027
##      Has cats  5.6920741  3.9837846  9.6758587
##      Has dogs  5.7237709  5.9840182 11.7077891
##      <NA>      27.3209549 48.8363946 76.1573495
##      Total    40.2332216 59.7667784 100.0000000
```

```
tally(~ cats_dogs_both | gender, format = "percent", margin = TRUE, data = OKCupid)
```

```
##                gender
## cats_dogs_both      F      M
##      Has Both  3.719368  1.610562
##      Has cats 14.147697  6.665550
##      Has dogs 14.226479 10.012282
##      <NA>      67.906456 81.711606
##      Total    100.000000 100.000000
```

```
tally(~ gender | cats_dogs_both, format = "percent", margin = TRUE, data = OKCupid)
```

```
##          cats_dogs_both
## gender  Has Both  Has cats  Has dogs  <NA>
## F      60.85482  58.82759  48.88857  35.87435
## M      39.14518  41.17241  51.11143  64.12565
## Total 100.00000 100.00000 100.00000 100.00000
```

Example 3.1: Exploring Marginal Distributions

```
SuperBowl <- read_csv("http://nhorton.people.amherst.edu/is5/data/Watch_the_Super_bowl.csv", skip = 1)
```

```
## Parsed with column specification:
## cols(
##   Plan = col_character(),
##   Sex = col_character()
## )
```

```
tally(~ Plan + Sex, data = SuperBowl)
```

```
##                Sex
## Plan      Female Male
## Commercials  156   81
## Game         200  279
## Wont Watch   160  132
```

Example 3.2: Exploring Percentages: Children and First-Class Ticket Holders First?

```
Titanic <- read_csv("http://nhorton.people.amherst.edu/is5/data/Titanic.csv")
```

```
## Parsed with column specification:
## cols(
##   Name = col_character(),
##   Survived = col_character(),
##   Boarded = col_character(),
##   Class = col_character(),
##   MWC = col_character(),
##   Age = col_double(),
##   Adut_or_Chld = col_character(),
##   Sex = col_character(),
##   Paid = col_double(),
##   Ticket_No = col_character(),
##   Boat_or_Body = col_character(),
##   Job = col_character(),
##   Class_Dept = col_character(),
##   Class_Full = col_character()
## )
```

```
tally(~ Class + Survived, format = "percent", margin = TRUE, data = Titanic)
```

```
##           Survived
## Class      Alive      Dead      Total
## 1      9.103261  5.570652 14.673913
## 2      5.389493  7.518116 12.907609
## 3      8.152174 24.003623 32.155797
## Crew    9.601449 30.661232 40.262681
## Total 32.246377 67.753623 100.000000
```

```
tally(~ Survived | Class, format = "percent", margin = TRUE, data = Titanic)
```

```
##           Class
## Survived      1      2      3      Crew
##   Alive 62.03704 41.75439 25.35211 23.84702
##   Dead 37.96296 58.24561 74.64789 76.15298
##   Total 100.00000 100.00000 100.00000 100.00000
```


```
tally(~ Class | Survived, format = "percent", margin = TRUE, data = Titanic)
```

```
##           Survived
## Class      Alive      Dead
## 1      28.230337  8.221925
## 2      16.713483 11.096257
## 3      25.280899 35.427807
## Crew  29.775281 45.254011
## Total 100.000000 100.000000
```

Section 3.2: Conditional Distributions


See displays on 68-69.

```
barplot(tally(~ cats_dogs_both + gender, data = OKCupid), beside = TRUE)
```



#There are many who don't own either

```
barplot(tally(~ cats_dogs_both + gender, data = filter(OKCupid, cats_dogs_both != "NA")), beside = TRUE)
```



Example 3.3: Finding Conditional Distributions: Watching the Super Bowl

```
tally(~ Plan + Sex, margin = TRUE, data = SuperBowl)
```


```
##           Sex
## Plan      Female Male Total
##  Commercials   156   81  237
##    Game       200  279  479
##  Wont Watch    160  132  292
##    Total       516  492 1008
```

```
tally(~ Plan | Sex, format = "percent", data = SuperBowl)
```

```
##           Sex
## Plan      Female    Male
##   Commercials 30.23256 16.46341
##   Game        38.75969 56.70732
##   Wont Watch  31.00775 26.82927
```

Example 3.4: Looking for Associations Between Variables: Still Watching the Super Bowl

```
barplot(tally(~ Plan | Sex, format = "percent", data = SuperBowl), beside = TRUE)
```



Examining Contingency Tables

See displays on page 72.

```
FishDiet <- read_csv("http://nhorton.people.amherst.edu/is5/data/Fish_diet.csv", skip = 1) %>%
  clean_names()
```

```
## Parsed with column specification:
## cols(
##   `Diet:Counts` = col_character(),
##   `Cancer:Counts` = col_character()
## )
```

```
tally(~ diet_counts + cancer_counts, data = FishDiet)
```

```
##           cancer_counts
## diet_counts    No    Yes
##   Large        507    42
##   Moderate    2769   209
##   Never        110    14
##   Small       2420   201
```

Random Matters

See display on page 74.

```
Nightmares <- read_csv("http://nhorton.people.amherst.edu/is5/data/Nightmares.csv", skip = 1)
```

```
## Parsed with column specification:
## cols(
##   Side = col_character(),
##   Dream = col_character()
## )
```

```
Nightmares$Dream <- ifelse(Nightmares$Dream == "N", "Nightmare", "SweetDreams")
tally(~ Side + Dream, data = Nightmares)
```

```
##      Dream
## Side Nightmare SweetDreams
##   L           9           13
##   R           6           35
```

Section 3.3: Displaying Contingency Tables


```
tally(~ Class + Survived, format = "count", data = Titanic)
```

```
##      Survived
## Class  Alive Dead
##   1      201  123
##   2      119  166
##   3      180  530
## Crew    212  677
```


```
tally(~ Class + Survived, format = "percent", data = Titanic)
```

```
##      Survived
## Class  Alive    Dead
##   1    9.103261 5.570652
##   2    5.389493 7.518116
##   3    8.152174 24.003623
## Crew  9.601449 30.661232
```

```
barplot(tally(~ Survived | Class, format = "percent", data = Titanic), beside = TRUE)
```




```
barplot(tally(~ Class | Survived, format = "percent", data = Titanic))
```



```
vcd::mosaic(tally(~ Survived + Class, data = Titanic),
  main = "Mosaic plot of Class by Survival",
  shade = TRUE)
```

Mosaic plot of Class by Survival



See the mosaic plots on page 77.


Section 3.4: Three Categorical Variables

```
tally(~ gender + cats_dogs_both + drugs_y_n, format = "percent", data = OKCupid)
```

```
## , , drugs_y_n = No
##
##      cats_dogs_both
## gender  Has Both  Has cats  Has dogs    <NA>
##      F  1.0243064  3.4199156  3.9437466 18.0187845
##      M  0.5922293  2.0819779  3.7769214 30.0719016
##
## , , drugs_y_n = Yes
##
##      cats_dogs_both
## gender  Has Both  Has cats  Has dogs    <NA>
##      F  0.2085314  0.8941828  0.6272626  2.9794972
##      M  0.1901807  0.8658225  0.9041923  6.9132342
##
## , , drugs_y_n = NA
##
##      cats_dogs_both
## gender  Has Both  Has cats  Has dogs    <NA>
##      F  0.2635837  1.3779757  1.1527618  6.3226732
##      M  0.1801712  1.0359842  1.3029044 11.8512587
```

Example 3.7: Looking for Associations Among Three Variables at Once


```
vcd::mosaic(tally(~ Sex + Survived + Class, data = Titanic), shade = TRUE)
```



Example 3.8: Simpson's Paradox: Gender Discrimination?

See tables on page 80.

```
# Making the data frame
school <- c("A", "B", "C", "D", "E", "F")
male_admits <- c(512, 313, 120, 138, 53, 22)
female_admits <- c(89, 17, 202, 131, 94, 24)
male_percent <- c(.621, .602, .369, .331, .277, .059)
female_percent <- c(.824, .680, .341, .349, .239, .070)
Berk <- data.frame(school, male_admits, female_admits, male_percent, female_percent)
Berk
```

```
##   school male_admits female_admits male_percent female_percent
## 1     A         512           89         0.621         0.824
## 2     B         313           17         0.602         0.680
## 3     C         120          202         0.369         0.341
## 4     D         138          131         0.331         0.349
## 5     E          53           94         0.277         0.239
## 6     F          22           24         0.059         0.070
```