

HW2_Fall2020

Rong Fan

9/20/2020

Variogram function and its applications

```
# Define the variogram function
variogram_func <- function(x, lag) {
  x <- as.matrix(x) # Make sure the x is a vector. It represents the observations of y_t.
  Lag <- NULL
  var_k <- NULL
  vario <- NULL
  for (k in 1:lag) {
    Lag[k] <- k
    var_k[k] <- sd(diff(x, k))^2
    vario[k] <- var_k[k] / var_k[1]
  }
  return(as.data.frame(cbind(Lag, vario)))
}
```

Variograms of chemical process viscosity data (Text: Fig 2.17, page 45)

```
library(readxl)
viscosity <- read_excel("AppendixB_datafile.xls", skip = 2, sheet = "B.3-VISC")

## New names:
## * `Time Period` -> `Time Period...1`
## * Reading -> Reading...2
## * `Time Period` -> `Time Period...3`
## * Reading -> Reading...4
## * `Time Period` -> `Time Period...5`
## * ...

readings <- na.omit(c(viscosity$Reading...2, viscosity$Reading...4,
                     viscosity$Reading...6, viscosity$Reading...8))
head(readings)


## [1] 86.7418 85.3195 84.7355 85.1113 85.1487 84.4775
class(readings)

## [1] "numeric"
x <- readings
lag_length <- 25
lag_readings <- 1:lag_length
```

```
z <- variogram_func(x, lag_length)
variogram_readings <- z$vario
variogram_readings
```

```
## [1] 1.000000 1.723942 2.376856 3.033604 3.348690 3.862875 4.129339 4.248845
## [9] 4.461848 4.569809 4.860165 5.274703 5.119000 4.715869 4.366605 4.298101
## [17] 4.605405 4.866596 5.201105 5.505501 5.436292 5.350848 4.968960 4.537469
## [25] 4.010016
```

```
plot(lag_readings, variogram_readings, type = "o", col = "dark red",
     main = "Variogram of chemical process viscosity")
```



Variograms of Whole Foods Market stock Price (Text: Fig 2.19, page 46)

```
library(readxl)
stock <- read_excel("AppendixB_datafile.xls", skip = 3, sheet = "B.7-WFMI")
```

```
## New names:
## * Date -> Date...1
## * Dollars -> Dollars...2
## * Date -> Date...3
## * Dollars -> Dollars...4
## * Date -> Date...5
## * ...
```

```
price <- na.omit(c(stock$Dollars...2, stock$Dollars...4,
                  stock$Dollars...6, stock$Dollars...8, stock$Dollars...10))
head(price)
```

```
## [1] 28.05 28.23 26.25 25.41 26.25 26.03
```

```
class(price)
```

```
## [1] "numeric"
```

```
x <- price
lag_length <- 25
lag_stock <- 1:lag_length
z <- variogram_func(x, lag_length)
variogram_stock <- z$vario
variogram_stock
```


```
## [1] 1.000000 2.139660 3.219850 4.411881 5.583359 6.784479 7.974550
```

```
## [8] 9.102236 10.109781 11.240052 12.378525 13.431489 14.627292 15.821995
```

```
## [15] 16.972685 18.123384 19.159940 20.130744 21.074192 21.999525 23.088225
```

```
## [22] 24.249945 25.582925 26.749122 27.881126
```

```
plot(lag_stock, variogram_stock, type = "o", col = "dark red",
     main = "Variogram of stock")
```




HW2

R version

```
library(readxl)
Blue <- read_excel("AppendixB_datafile.xls",
  sheet = "B.4-BLUE"
)
```

```
x <- acf(Blue[, 2], lag.max = 25, type = "correlation",
  main = "ACF of Blue and Gorgonzola cheese") # ACF plot
```



2.2 a)

```
x$acf[1:26] # values of ACFs for the first 25 lags
```

```
## [1] 1.00000000 0.92255797 0.85024468 0.79968986 0.74328987 0.69263229
## [7] 0.64664768 0.59209915 0.52647671 0.46521790 0.39867153 0.34387841
## [13] 0.28989096 0.22336880 0.15486809 0.09742316 0.05132157 0.01216789
## [19] -0.04327517 -0.10876777 -0.17264515 -0.22894671 -0.27937815 -0.30055353
## [25] -0.31694514 -0.35312179
```

```
x <- Blue$Production
lag_length <- length(x) / 4
lag_cheese <- 1:lag_length
z <- variogram_func(x, lag_length)
variogram_cheese <- z$vario
variogram_cheese
```

```
plot(lag_cheese, variogram_cheese, type = "o", col = "dark red",
     main = "Variogram of Blue and Gorgonzola cheese")
```

More general, let's get all graphs related to the Bule and Gorgonzola cheese data. You can ignore the code below if you don't want to get all the outputs in one function.

```
time.series.graphics <- function(data, title = "", xlab = "", ylab = "") {
  # Use the variogram_func() defined above
  # or include the definition of the variogram function within the chunk so that the function
  # time.series.graphics() can be used independently.
  # Define the variogram function
  variogram <- function(x, lag) {
    x <- as.matrix(x)
    Lag <- NULL
    var_k <- NULL
    vario <- NULL
    for (k in 1:lag) {
      Lag[k] <- k
      var_k[k] <- sd(diff(x, k))^2
      vario[k] <- var_k[k] / var_k[1]
    }
    return(as.data.frame(cbind(Lag, vario)))
  }
  # End of variogram function

  layout(mat = matrix(c(0, 1, 1, 0, 2, 2, 3, 3), nrow = 2, byrow = T))

  fit.data <- lm(data[, 2] ~ data[, 1])
  plot(data, type = "l", ylab = ylab, xlab = xlab, main = title)
  points(data, pch = 16, cex = .5)
  lines(data[, 1], fit.data$fitted.values, col = "red", lty = 2)
  legend(1990, 12000, c("Actual", "Fit"), pch = (c(16, NA)), lwd = c(.5, .5),
        lty = c(1, 2), cex = .55, col = c("black", "red"))




  acf.data <- acf(data[, 2], lag.max = 17, type = "correlation", main = "")


  data.vario <- variogram(data[, 2], length(data[, 2]) / 4)

  plot(data.vario, type = "l")
  points(data.vario)

  par(mfrow = c(2, 2), oma = c(0, 0, 0, 0))
  qqnorm(fit.data$residuals, datax = T, pch = 16, xlab = "Residual", main = "")
  qqline(fit.data$residuals, data = T)
  plot(fit.data$fitted.values, fit.data$residuals, pch = 16,
       xlab = "Fitted Value", ylab = "Residual")
  abline(h = 0)
  hist(fit.data$residuals, col = "grey", xlab = "Residual", main = "")
  plot(fit.data$residuals, type = "l", xlab = "Observation Order", ylab = "Residual")
  points(fit.data$residuals, pch = 16, cex = .5)
  abline(h = 0)
}
```

```
newBlue <- as.data.frame(Blue)
time.series.graphics(newBlue)
```





The time series plot, ACF and variogram show that the US production of blue and gorgonzola cheeses data is non-stationary.


```
library(readxl)
Anomaly_C02 <- read_excel("AppendixB_datafile.xls", skip = 0, sheet = "B.6-GSAT-C02")
head(Anomaly_C02)
```

2.4.1 ACF and variogram of Anomaly

```
## # A tibble: 6 x 3
##   Year Anomaly  C02
##   <dbl> <dbl> <dbl>
## 1  1880  -0.11  291.
## 2  1881  -0.13  291.
## 3  1882  -0.01  292.
## 4  1883  -0.04  292.
## 5  1884  -0.42  293.
## 6  1885  -0.23  293
```


```
plot(Anomaly_C02$Anomaly, type = "l", main = "Time series of Anomaly")
```

Time series of Anomaly



```
x <- acf(Anomaly_CO2$Anomaly, lag.max = 25, type = "correlation",  
        main = "ACF of Anomaly") # ACF plot
```


ACF of Anomaly



```
x$acf[1:26] # values of ACFs for the first 25 lags
```


```
## [1] 1.0000000 0.8071948 0.6942248 0.6670964 0.6787528 0.6235193 0.6184973
## [8] 0.5817489 0.5538216 0.4980878 0.4995987 0.4732532 0.4447530 0.4187977
## [15] 0.3996724 0.3696274 0.3434193 0.3114057 0.2871918 0.2427561 0.2549324
## [22] 0.2498699 0.1944899 0.1524225 0.1064507 0.1255333
```

```
x <- Anomaly_CO2$Anomaly
lag_length <- 25
lag_Anomaly <- 1:lag_length
z <- variogram_func(x, lag_length)
variogram_Anomaly <- z$vario
variogram_Anomaly
```

```
## [1] 1.000000 1.508014 1.491944 1.305226 1.507647 1.443786 1.436555 1.452578
## [9] 1.719642 1.614282 1.711940 1.737889 1.813212 1.756689 1.767938 1.833040
## [17] 1.897329 1.966501 2.212713 2.069221 2.082261 2.391755 2.618794 2.748224
## [25] 2.420406
```

```
plot(lag_Anomaly, variogram_Anomaly, type = "o", col = "dark red",
     main = "Variogram of Anomaly")
```

Variogram of Anomaly



The time series plot, ACF and variogram show that the anomaly data is non-stationary.


```
x <- Anomaly_C02$C02
lag_length <- 25
lag_C02 <- 1:lag_length
z <- variogram_func(x, lag_length)
variogram_C02 <- z$vario
variogram_C02
```

2.4.2 ACF and variogram of CO2


```
## [1] 1.000000 3.684466 7.751564 13.189532 20.099805 28.440840
## [7] 38.038272 48.450856 60.223840 72.931205 86.374042 100.487887
## [13] 115.864015 132.391615 149.733282 168.023418 186.988846 205.541761
## [19] 224.244712 243.576099 263.123240 282.636155 302.050303 321.739180
## [25] 341.597948
```

```
plot(lag_C02, variogram_C02, type = "o", col = "dark red",
     main = "Variogram of CO2")
```

Variogram of CO2



```
plot(x, type = "l", main = "Time series of CO2")
```



The time series plot, ACF and variogram show that the CO2 data is non-stationary.

JMP version

2.2 a) ACF and variogram of Blue and Gorgonzola cheese

2.4.1 ACF and variogram of Anomaly

2.4.2 ACF and variogram of CO2




Figure 1: ACF and Variogram of blue and gorgonzola cheese




Figure 2: ACF and Variogram of Anomaly




Figure 3: ACF and Variogram of CO2