

Quantum Field Theory in Curved Spacetime

*Study Notes on Divergences, Functional Methods,
and Zeta Function Regularisation*

Mohamed Amine Ezzine

KTH Royal Institute of Technology · ISAE-SUPAERO

ezzine@kth.se · amine-physics.github.io

First version: May 2026

Updated regularly as the notes progress.

Introduction

These notes document my personal study of quantum field theory in curved spacetime, following primarily the treatment in [6]. They are not a substitute for the textbook but rather a record of the calculations I worked through, the points where I got stuck, and the understanding I eventually reached.

The central theme running through these notes is the problem of *ultraviolet divergences* in quantum field theories formulated on curved backgrounds. In flat spacetime, divergences are handled by standard renormalisation; in curved spacetime, new divergences appear that are sensitive to the local geometry, and their treatment requires more sophisticated tools. The natural language for this is the *heat kernel*, a Green's function for the heat equation on the Manifold, whose short-time asymptotic expansion encodes precisely the geometric divergences of the theory. The connection between heat kernels and spectral zeta functions then provides a clean and mathematically rigorous regularisation scheme.

This connects directly to the mathematics notes on this site, where the heat kernel and its asymptotic expansion are developed from the differential-geometric side. The present notes approach the same object from the physics side: as a tool for computing one-loop effective actions and understanding the renormalisation of quantum fields in gravitational backgrounds.

Conventions. Throughout these notes we use natural units $\hbar = c = 1$. The spacetime metric has signature $(-, +, +, +)$. Einstein's summation convention is always in force. The Riemann tensor is defined by

$$[\nabla_\mu, \nabla_\nu]V^\rho = R^\rho{}_{\sigma\mu\nu}V^\sigma,$$

and the Ricci tensor by $R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}$.

A note on updates. These notes are updated regularly as my studies progress. The current version covers:

- Divergences in curved spacetime
- Functional determinants and one-loop effective actions
- Zeta function regularisation

Forthcoming sections will treat the heat kernel expansion in detail, the effective potential, and applications to Yang-Mills theory.

Contents

Introduction	I
1 Divergences in Curved Spacetime	4
1.1 Introduction	4
1.2 Euclidean Functional Integral	4
1.3 Functional Determinant	6
1.4 Effective Action	9
1.5 Zeta Function regularization	12
2 Heat Kernel Expansion	13
3 Effective Potential	13

I Divergences in Curved Spacetime

I.1 Introduction

Quantum field theories in Minkowski spacetime generically exhibit ultraviolet divergences. In ϕ^4 theory for example, the vacuum energy density acquires a quartic divergence $\sim \Lambda^4$, the mass parameter receives a quadratic divergence $\sim \Lambda^2$, and the self-coupling is renormalised by a logarithmic divergence $\sim \log \Lambda$, where Λ is the UV cutoff.

Since spacetime is locally Minkowski and UV divergences arise from short-distance behaviour, they are governed by the local structure of the theory. One therefore expects every divergence present in flat spacetime to persist in the curved case. Beyond this, the curvature tensor introduces a new dimensionful scale into the theory, and divergences proportional to powers of curvature invariants are expected to appear. For instance, in ϕ^4 theory the scalar field has mass dimension $[\phi] = 1$ in four dimensions, and the mass counterterm $\sim \Lambda^2 \phi^2$ has dimension four. The Ricci scalar has dimension $[R] = 2$ (inverse length squared), so the combination $R\Lambda^2$ has the correct dimension four to appear as a new counterterm in the curved-spacetime theory, with no flat-space analogue. This is the first sign that renormalisation in curved spacetime requires extending the gravitational action beyond the Einstein–Hilbert term.

I.2 Euclidean Functional Integral

Consider a real scalar field ϕ propagating on a Manifold M with a fixed external metric $g_{\mu\nu}$. The action of the field is given by

$$S(\phi, g) = -\frac{1}{2} \int dx^d \sqrt{|g|} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi. \quad (1)$$

If we consider a source j coupled to the field, then the functional integral is

$$\mathcal{Z} = \int \mathcal{D}\phi e^{i(S(\phi, g) + \int dx^d \sqrt{|g|} \phi j)} \quad (2)$$

Physical Remark. The path integral defined in equation 2 is not well defined in Minkowski signature. To see this consider the case without a source term. Field configurations with large and small kinetic energy, (action), contribute equally in amplitude to the integral, hence the oscillating behavior is intense and the integral will generically won't converge.

To ensure that the path integral formulation is better defined, One performs a Wick rotation. Consider that the metric is of the form

Derivation I.1 (Eucliden Signature).

$$g_{\mu\nu} = \begin{pmatrix} g_{00} & 0 \\ 0 & g_{ij} \end{pmatrix}, \quad (3)$$

with $g_{00} < 0$. Let $t = it_E$, then the action of the theory is

$$iS(\phi, g) = i \frac{1}{2} \int d^{d-1}x dt \sqrt{|g|} \left[-g^{00} (\partial_t \phi)^2 - g^{ij} \partial_i \phi \partial_j \phi \right], \quad (4a)$$

$$= \frac{1}{2} \int d^{d-1}x dt_E \sqrt{|g|} \left[g^{00} (\partial_{t_E} \phi)^2 - g^{ij} \partial_i \phi \partial_j \phi \right], \quad (4b)$$

$$= -S_E(\phi, g_E). \quad (4c)$$

$S_E(\phi, g)$ is the action with a Euclidean signature,

$$S_E(\phi, g) = \frac{1}{2} \int d^d x_E \sqrt{g_E} g_E^{\mu\nu} \partial_\mu \phi \partial_\nu \phi. \quad (5)$$

g_E is a positive definite metric defined on an analytically continued Manifolds M_E .

$$(g_E)_{\mu\nu} \begin{pmatrix} -g_{00} & 0 \\ 0 & g_{ij} \end{pmatrix} \quad (6)$$

Similarly we get for the source term

$$i \int d^d x \sqrt{|g|} j \phi = \int d^d x_E \sqrt{g_E} j \phi \quad (7)$$

Henceforward we drop the subscript E , unless explicitly indicated, we work in Euclidean signature.

Physical Remark. Working in the Euclidean signature does not entirely solve the problem of convergence of the action. However now field configuration with higher energy contribute smaller to the functional integral. Hence, in Euclidean signature there is better chance that the integral is well defined.

If the Manifold M is compact without boundary, than when performing integration by parts the boundary terms cancel.

Derivation 1.2 (Simplified action and Laplacian operator).

$$S(\phi, g) = \frac{1}{2} \int d^d x \sqrt{g} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi, \quad (8a)$$

$$= \frac{1}{2} \int d^d x \sqrt{g} g^{\mu\nu} \partial_\mu (\phi \partial_\nu \phi) - \frac{1}{2} \int d^d x \sqrt{g} g^{\mu\nu} \phi \partial_\mu \partial_\nu \phi, \quad (8b)$$

$$= -\frac{1}{2} \int d^d x \sqrt{g} g^{\mu\nu} \phi \partial_\mu \partial_\nu \phi. \quad (8c)$$

The first term in equation 8b is a boundary term, hence it is equal to zero. The action can then be expressed as

$$S(\phi, g) = \frac{1}{2} \int d^d x \sqrt{g} \phi \Delta \phi. \quad (9)$$

With $\Delta = -\nabla_\mu \nabla^\mu$ is the covariant Laplacian operator.

In Euclidean Signature the functional integral is given by

$$\mathcal{Z}(j, g) = \int \mathcal{D}\phi e^{-S(\phi, g) + \int d^d x \sqrt{g} j \phi}.$$

Let $\mathcal{W}(j, g)$ be the functional defined by

$$\mathcal{Z}(j, g) = e^{\mathcal{W}(j, g)}.$$

$\mathcal{W}(j, g)$ is called the generating functional of connected Greens functions.

1.3 Functional Determinant

In this subsection we set the source to $j = 0$. In curved space time the family of exponential functions $\{e^{ik_\mu x^\mu}\}$ are no longer eigenfunction of the Laplacian operator. Instead the spectral decomposition of the operator is used. When M is compact without boundary the spectrum of the operator Δ is discrete. Let $\{\varphi_n\}_{n \in \mathbb{N}}$ be the eigenfunction of Δ ,

$$\Delta \varphi_n = \lambda_n \varphi_n, \quad (10)$$

with $\{\lambda_n\}_{n \in \mathbb{N}}$ is the corresponding family of eigenvalues.

Physical Remark. The operator $\Delta = -\nabla^2$ has a zero eigenvalue corresponding to the eigenfunction $\varphi_n = \text{const}$. This leads to an indumped term in the functional integral leading to divergences. Hence this case needs to be treated separately.

The eigenfunctions $\{\varphi_n\}_{n \in \mathbb{N}}$ are taking to be dimensionless and orthonormal with respect to the inner product on $C^\infty(M)$.

$$\langle \varphi_n | \varphi_m \rangle = \mu^d \int_M d^d x \sqrt{g} \varphi_n(x) \varphi_m(x) \quad (11a)$$

$$= \delta_{nm} \quad (11b)$$

with μ is a constant with dimension of mass.

The eigenfunctions of the operator Δ form a basis on the space of function on M . Hence the field ϕ can be decomposed as

$$\phi(x) = \sum_{n \in \mathbb{N}} a_n \varphi_n(x), \quad (12)$$

a_n are constant with the same mass dimension as ϕ . The Euclidean action is then

$$S(\phi, g) = \frac{1}{2\mu^d} \sum_{n \in \mathbb{N}} \lambda_n a_n^2 \quad (13a)$$

$$= \frac{1}{2} \sum_{n \in \mathbb{N}} \tilde{\lambda}_n \tilde{a}_n^2 \quad (13b)$$

with

$$\tilde{\lambda}_n = \frac{\lambda_n}{\mu^2} \quad , \quad \tilde{a}_n = \frac{a_n}{\mu^{\frac{d}{2}-1}}.$$

The path integral measure is then $\mathcal{D}\phi = \mathcal{N} \prod_{n \in \mathbb{N}} \frac{d a_n}{\mu^{\frac{d}{2}-1}} = \mathcal{N} \prod_{n \in \mathbb{N}} d \tilde{a}_n$. $\mathcal{N}$ is a renormalization constant such that

$$\int \mathcal{D}\phi \exp \left[\frac{-\mu^2}{2} \int d^d x \sqrt{g} \phi^2 \right] = 1 \quad (14)$$

Derivation 1.3 (Formal expression of $\mathcal{N}$). Consider first the integral

$$\int d^d x \sqrt{g} \phi^2 = \mu^{-d} \sum_{n,m \in \mathbb{N}} a_n a_m \mu^d \int d^d x \sqrt{g} \varphi_n \varphi_m, \quad (15a)$$

$$= \mu^{-d} \sum_{n,m \in \mathbb{N}} a_n a_m \langle \varphi_n | \varphi_m \rangle, \quad (15b)$$

$$= \mu^{-2} \sum_{n \in \mathbb{N}} \tilde{a}_n^2. \quad (15c)$$

Injecting this result in equations 14, we get

$$1 = \int \mathcal{D}\phi \exp \left[\frac{-\mu^2}{2} \int d^d x \sqrt{g} \phi^2 \right], \quad (16a)$$

$$= \mathcal{N} \prod_{n \in \mathbb{N}} \int d \tilde{a}_n \exp \left[-\frac{1}{2} \sum_{n \in \mathbb{N}} \tilde{a}_n^2 \right], \quad (16b)$$

$$= \mathcal{N} \prod_{n \in \mathbb{N}} \left[\int d \tilde{a}_n e^{-\frac{1}{2} \tilde{a}_n^2} \right], \quad (16c)$$

$$= \mathcal{N} \prod_{n \in \mathbb{N}} \sqrt{2\pi}. \quad (16d)$$

In going from 16c to 16d we used the Gaussian integral formula. Equation 16d implicitly defines the value of the normalization constant $\mathcal{N}$.

The path integral with zero source can then be expressed as a formal determinant of the scaled operator $\tilde{\Delta} = \frac{1}{\mu^2} \Delta$.

$$\mathcal{Z}(0, g) = e^{\mathcal{W}(0, g)} = \mathcal{N} \prod_{n \in \mathbb{N}} \left[\int d \tilde{a}_n e^{-\frac{1}{2} \tilde{\lambda}_n \tilde{a}_n^2} \right], \quad (17a)$$

$$= \mathcal{N} \prod_{n \in \mathbb{N}} \sqrt{\frac{2\pi}{\tilde{\lambda}_n}} \quad (17b)$$

$$= \det(\tilde{\Delta})^{-\frac{1}{2}} \quad (17c)$$

Consider next the case of non trivial source term j . We get

$$\mathcal{Z}(j, g) = (\det \tilde{\Delta})^{-\frac{1}{2}} e^{\frac{1}{2} \int d^d x \sqrt{g} j \Delta^{-1} j} \quad (18)$$

Derivation 1.4 (Proof of equation 18). Consider the spectral decomposition of the source term j

$$j(x) = \sum_{n \in \mathbb{N}} b_n \varphi_n(x). \quad (19)$$

Then the source term in the functional integral gives

$$\int d^d x \sqrt{g} j \phi = \mu^{-d} \sum_{n, m \in \mathbb{N}} a_n b_m \mu^d \int d^d x \sqrt{g} \varphi_m \varphi_n, \quad (20a)$$

$$= \frac{1}{\mu^2} \sum_{n \in \mathbb{N}} \tilde{a}_n \tilde{b}_n. \quad (20b)$$

Consider now the sum of the action and the source term

$$-S(\phi, j) + \int d^d x \sqrt{g} j \phi = \sum_{n \in \mathbb{N}} -\frac{1}{2} \tilde{\lambda}_n \tilde{a}_n^2 + \frac{\tilde{b}_n}{\mu^2} \tilde{a}_n, \quad (21a)$$

$$= \sum_{n \in \mathbb{N}} -\frac{1}{2} \tilde{\lambda}_n \left(\tilde{a}_n^2 - 2 \frac{\tilde{b}_n}{\tilde{\lambda}_n} \tilde{a}_n \right), \quad (21b)$$

$$= \sum_{n \in \mathbb{N}} -\frac{1}{2} \tilde{\lambda}_n \left(\tilde{a}_n - \frac{\tilde{b}_n}{\tilde{\lambda}_n} \right)^2 + \frac{1}{2} \frac{1}{\mu^d} \frac{b_n^2}{\lambda_n}. \quad (21c)$$

Plugging 21c into the expression of the functional integral we get

$$\mathcal{Z}(j, g) = e^{\mathcal{W}(j, g)} = \mathcal{N} \prod_{n \in \mathbb{N}} \int d\tilde{a}_n e^{-\frac{1}{2} \tilde{\lambda}_n \left(\tilde{a}_n - \frac{\tilde{b}_n}{\tilde{\lambda}_n} \right)^2 + \frac{1}{2} \frac{1}{\mu^d} \frac{b_n^2}{\lambda_n}}, \quad (22a)$$

$$= \left[\mathcal{N} \prod_{n \in \mathbb{N}} \int d\tilde{a}_n e^{-\frac{1}{2} \tilde{\lambda}_n \left(\tilde{a}_n - \frac{\tilde{b}_n}{\tilde{\lambda}_n} \right)^2} \right] e^{\frac{1}{2} \frac{1}{\mu^d} \sum_{n \in \mathbb{N}} \frac{b_n^2}{\lambda_n}}. \quad (22b)$$

The first term in 22b is, after redefining the integration variable, the same term appearing in the case with zero source treated earlier. The second term in the product of 22b can be expressed in terms of the sources again as follows

$$\frac{1}{\mu^d} \sum_{n \in \mathbb{N}} \frac{b_n^2}{\lambda_n} = \frac{1}{\mu^d} \sum_{n, m \in \mathbb{N}} b_n b_m \frac{1}{\lambda_n} \mu^d \int d^d x \sqrt{g} \varphi_n \varphi_m, \quad (23a)$$

$$= \int d^d x \left(\sum_{n \in \mathbb{N}} b_n \varphi_n \right) \left(\Delta^{-1} \sum_{m \in \mathbb{N}} b_m \varphi_m \right), \quad (23b)$$

$$= \int d^d x j \Delta^{-1} j. \quad (23c)$$

Putting everything together we get

$$\mathcal{Z}(j, g) = (\det \tilde{\Delta})^{-\frac{1}{2}} e^{\frac{1}{2} \int d^d x \sqrt{g} j \Delta^{-1} j} \quad (24)$$

Physical Remark. The functional $\mathcal{W}(j, g)$, is given by

$$\mathcal{W}(j, g) = -\frac{i}{2} \ln(\det \tilde{\Delta}) + \frac{i}{2} \int d^d x \sqrt{g} j \Delta^{-1} j. \quad (25)$$

The first term in this expression reflect the vacuum contribution while the source term corresponds to the propagator.

1.4 Effective Action

The expectation value of the field ϕ is derived from the generating function $\mathcal{W}$ by taking its functional derivative with respect to the source j . Hence,

$$\varphi = \frac{\delta \mathcal{W}}{\delta j} \quad (26)$$

φ_j is a classical C-number field. The generating function is also used to obtain the connected Green's functions, for example the two point correlator is given by

$$G(x, y)_j = \langle \phi(x) \phi(y) \rangle - \langle \phi(x) \rangle \langle \phi(y) \rangle \quad (27a)$$

$$= \frac{\delta^2 \mathcal{W}}{\delta j(x) \delta j(y)} \quad (27b)$$

Derivation 1.5 (Expectation Value). Starting for the relation between $\mathcal{Z}$ and $m\mathcal{W}$ and taking the functional derivative with respect to the source j we get

$$\frac{\delta \mathcal{Z}}{\delta j} = \frac{\delta}{\delta j} e^{\mathcal{W}}, \quad (28a)$$

$$= \mathcal{Z} \frac{\delta \mathcal{W}}{\delta j}. \quad (28b)$$

Using the expression of $\mathcal{Z}$

$$\mathcal{Z}(j, g) = \int \mathcal{D}\phi e^{-S(\phi, g) + \int d^d x \sqrt{g} j \phi}, \quad (29)$$

we get

$$\frac{\delta \mathcal{Z}}{\delta j} = \int \mathcal{D}\phi \phi e^{-S(\phi, g) + \int d^d x \sqrt{g} j \phi}, \quad (30a)$$

$$= \mathcal{Z} \langle \phi \rangle_j. \quad (30b)$$

Equating 30b and 28b we get

$$\langle \phi \rangle_j = \frac{\delta \mathcal{W}}{\delta j} \quad (31)$$

The relationship between the field expectation value and the source depicted in equation 26 can be inverted using a Legendre transformation. Let j_φ the source needed to produce an expectation value of the field φ . Then the effective action is the Legendre transform of the generating functional $\mathcal{W}$.

$$\Gamma(\varphi, g) = -\mathcal{W}(j_\varphi, g) + \int d^d x \sqrt{g} j_\varphi \varphi. \quad (32)$$

Derivation 1.6 (φ in terms of j). Given a certain expectation value of the field ϕ , we want to find the expression of the corresponding source. We use equation 26. To compute the functional derivative, we consider a perturbation δj of the source, then $\mathcal{W} \rightarrow \mathcal{W} + \delta \mathcal{W}$, to first order in the perturbation we get

$$\delta \mathcal{W} = \frac{1}{2} \int d^d x \sqrt{g} (\delta j \Delta^{-1} j + j \Delta^{-1} \delta j) \quad (33)$$

At this stage, it is useful to discuss the self-adjointness properties of the operator Δ^{-1} . For a compact Manifold M without Boundaries, the boundary terms in integration by part conceal out. Hence taking two function f and g , we get

$$\langle f | \Delta g \rangle = \int d^d x \sqrt{g} f \Delta g, \quad (34a)$$

$$= - \int d^d x \sqrt{g} f \nabla_\mu \nabla^\mu g \quad (34b)$$

$$= - \int d^d x \sqrt{g} \nabla_\mu (f \nabla^\mu g) + \int d^d x \sqrt{g} \nabla_\mu f \nabla^\mu g. \quad (34c)$$

The first term being a boundary term it cancels out. The second term can be reworked in the opposite step so the Laplacian operator is now acting on f .

$$\int d^d x \sqrt{g} \nabla_\mu f \nabla^\mu g = \int d^d x \sqrt{g} \nabla^\mu (\nabla_\mu f g) - \int d^d x \sqrt{g} \nabla_\mu \nabla^\mu f g, \quad (35a)$$

$$= \int d^d x \sqrt{g} \Delta f g, \quad (35b)$$

$$= \langle \Delta f | g \rangle \quad (35c)$$

Hence the operator Δ is self-adjoint. Consider next the inverse operator

$$\langle f | \Delta^{-1} g \rangle = \langle \Delta (\Delta^{-1} f) | \Delta^{-1} g \rangle, \quad (36a)$$

$$= \langle \Delta^{-1} f | \Delta (\Delta^{-1} g) \rangle, \quad (36b)$$

$$= \langle \Delta^{-1} f | g \rangle. \quad (36c)$$

Thus the inverse Laplacian is also self-adjoint. Going back to the variation of the generating

functional 33. Using the self-adjoinance of the operator we get

$$\delta \mathcal{W} = \int d^d x \sqrt{g} \delta j \Delta^{-1} j. \quad (37)$$

Thus, we finally get

$$\varphi = \frac{\delta \mathcal{W}}{\delta j}, \quad (38a)$$

$$= \Delta^{-1} j. \quad (38b)$$

Hence given an expectation value φ , the corresponding source is given by $\Delta\varphi$. The effective action is therefore given by

$$\Gamma(\varphi, g) = -\mathcal{W}(j_\varphi, g) + \int d^d x \sqrt{g} j_\varphi \varphi, \quad (39a)$$

$$= \frac{1}{2} \ln(\det \tilde{\Delta}) + \frac{1}{2} \int d^d x \sqrt{g} \varphi \Delta \varphi, \quad (39b)$$

$$= \frac{1}{2} \text{Tr}(\ln(\tilde{\Delta})) + \frac{1}{2} \int d^d x \sqrt{g} \varphi \Delta \varphi. \quad (39c)$$

In going from 39b to 39c we used the identity

$$\ln(\det(\tilde{\Delta})) = \text{Tr}(\ln(\tilde{\Delta})). \quad (40)$$

The second part in equation 39c corresponds to the action of the classical field φ .

Physical Remark (On the physical meaning of the effective action). As I encountered this material for the first time, a question that kept bothering me was why one defines the effective action at all, since it initially appeared to be a mere formal transformation of the generating functional $\mathcal{W}$. It was only through further study that the importance of this object became clear.

The discomfort starts earlier, with the introduction of the source j . This seems at first to be an artificial addition to the theory, a parameter with no clear physical meaning, especially in the case of a free theory. The correlators are obtained from $\mathcal{W}$ by differentiating with respect to j , but j itself is not a physical quantity. The real physics should be expressible without it.

The effective action Γ resolves this by reformulating the dynamics entirely in terms of φ , the expectation value of the field, which is the physically observable quantity. The source j is eliminated via the Legendre transform, and what remains is a functional whose stationary points are the true quantum-corrected field configurations.

More precisely, $\Gamma[\varphi]$ plays the role of the classical action $S[\phi]$, but with all quantum corrections already incorporated. The equation of motion $\delta\Gamma/\delta\varphi = 0$ at $j = 0$ looks identical in form to the classical variational principle, but now encodes the full quantum dynamics. In this sense, Γ is the answer to the question: if one integrates out all quantum fluctuations, what effective

classical theory remains?

1.5 Zeta Function regularization

In the expression of the effective action 39c, one encounters the quantity

$$\frac{1}{2} \text{Tr} (\ln (\tilde{\Delta})) = \frac{1}{2} \sum_{n \in \mathbb{N}} \ln (\tilde{\lambda}_n) . \quad (41)$$

One way to make sense of this expression is through the Zeta function regularization scheme. In an analogous manner to the Riemann Zeta function

$$\zeta_R(s) = \sum_{n \in \mathbb{N}^*} n^{-s} , \quad s \in \mathbb{C} , \quad (42)$$

one defines for an operator O with a discrete spectrum $\{\gamma_n\}_{n \in \mathbb{N}}$, the Zeta function $\zeta_O(s)$ given by

$$\zeta_O(s) = \sum_{n \in \mathbb{N}^*} \gamma_n^{-s} , \quad s \in \mathbb{C} . \quad (43)$$

Derivation 1.7 (Zeta function of the Laplacian operator). Using the previous definition, the Zeta function of the Laplacian operator is

$$\zeta_{\tilde{\Delta}}(s) = \sum_{n \in \mathbb{N}} \tilde{\lambda}_n^{-s} . \quad (44)$$

In this expression degenerate eigenvalues are counted separately otherwise a multiplicity coefficient is added. Using

$$\ln(z) = -\frac{d}{ds} z^{-s} \Big|_{s=0} , \quad (45)$$

we get

$$\frac{1}{2} \text{Tr} (\ln (\tilde{\Delta})) = -\frac{1}{2} \sum_{n \in \mathbb{N}} \frac{d}{ds} \tilde{\lambda}_n^{-s} \Big|_{s=0} , \quad (46a)$$

$$= -\frac{1}{2} \frac{d}{ds} \zeta_{\tilde{\Delta}}(s) \Big|_{s=0} . \quad (46b)$$

Equation 46b, provides an alternative expression of the trace part in the effective action. However to use this new formulation, one still needs to explicitly compute an expression for the Zeta function. This can be done approximately by using the following result on the density of eigenvalues of the Laplacian:

Let M is a compact d dimensional Manifold with a metric tensor $g_{\mu\nu}$. Let $\{\lambda_n\}_{n \in \mathbb{N}}$ be the set of eigenvalues of Δ , in ascending order, i.e $\lambda_1 \leq \lambda_2 \leq \dots$. For $x \in \mathbb{R}$, let $N(x) = \# \{\lambda_i \leq x, i \in \mathbb{N}\}$ be the counting function. It can be shown (Weyl's theorem [6]) that for large values of x , the counting function $N(x) \simeq C x^{\frac{d}{2}}$, where C is a constant that depends on the geometry.

Using this result, the density of eigenvalues in a small interval $[\lambda, \lambda + d\lambda]$ is $\frac{dC}{2} \lambda^{\frac{d}{2}-1} d\lambda$. The sum in the zeta function can then be approximated by

$$\zeta_{\hat{\Delta}}(s) \approx \frac{C d}{2} \int d\lambda \lambda^{\frac{d}{2}-1-s}. \quad (47)$$

This integral is well defined for $\Re(s) > \frac{d}{2}$. Hence, for $\Re(s) > \frac{d}{2}$, the Zeta function is convergent, then through analytical continuity it is extended to the whole complex plane allowing for the computation of its derivative at $s = 0$. Therefore obtaining a finite value for the effective action Γ .

2 Heat Kernel Expansion

Remark. This section is in preparation. It will cover the Seeley–DeWitt asymptotic expansion

$$\text{Tr } e^{-tA} \sim (4\pi t)^{-n/2} \sum_{k=0}^{\infty} a_k(A) t^k \quad \text{as } t \rightarrow 0^+,$$

the computation of the first few coefficients a_0, a_1, a_2 in terms of geometric invariants, and their role as counterterms in curved spacetime QFT. See the mathematics notes for the purely geometric treatment.

3 Effective Potential

Remark. This section is in preparation. It will derive the one-loop effective potential from the zeta-regularised determinant, discussing its renormalisation and the Coleman-Weinberg mechanism in curved spacetime.

References

- [1] L. Parker and D. Toms, *Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity*, Cambridge University Press, 2009.
- [2] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press, 1982.
- [3] D. V. Vassilevich, *Heat kernel expansion: user's manual*, Physics Reports **388** (2003) 279–360.
- [4] S. A. Fulling, *Aspects of Quantum Field Theory in Curved Spacetime*, Cambridge University Press, 1989.
- [5] S. W. Hawking, Zeta function regularization of path integrals in curved spacetime, *Communications in Mathematical Physics* **55** (1977) 133–148.
- [6] R. Percacci, *An Introduction to Covariant Quantum Gravity and Asymptotic Safety*, World Scientific, 2017.