



Linear Regression with One Variable

Quiz, 5 questions

1
point

1. Consider the problem of predicting how well a student does in her second year of college/university, given how well she did in her first year.

Specifically, let x be equal to the number of "A" grades (including A-, A and A+ grades) that a student receives in their first year of college (freshmen year). We would like to predict the value of y , which we define as the number of "A" grades they get in their second year (sophomore year).

Here each row is one training example. Recall that in linear regression, our hypothesis is $h_{\theta}(x) = \theta_0 + \theta_1 x$, and we use m to denote the number of training examples.

x	y
5	4
3	4
0	1
4	3

For the training set given above (note that this training set may also be referenced in other questions in this quiz), what is the value of m ? In the box below, please enter your answer (which should be a number between 0 and 10).

Enter answer here

1
point

2. Consider the following training set of $m = 4$ training examples:

x	y
1	0.5
2	1
4	2
0	0

Consider the linear regression model $h_{\theta}(x) = \theta_0 + \theta_1 x$. What are the values of θ_0 and θ_1 that you would expect to obtain upon running gradient descent on this model? (Linear regression will be able to fit this data perfectly.)

- ☐ $\theta_0 = 0.5, \theta_1 = 0$
- ☐ $\theta_0 = 0.5, \theta_1 = 0.5$

☐ $\theta_0 = 0, \theta_1 = 0.5$

☐ $\theta_0 = 1, \theta_1 = 0.5$

☐ $\theta_0 = 1, \theta_1 = 1$

1
point

3. Suppose we set $\theta_0 = -1, \theta_1 = 0.5$. What is $h_\theta(4)$?

Enter answer here

1
point

4. Let f be some function so that

$f(\theta_0, \theta_1)$ outputs a number. For this problem,

f is some arbitrary/unknown smooth function (not necessarily the cost function of linear regression, so f may have local optima).

Suppose we use gradient descent to try to minimize $f(\theta_0, \theta_1)$

as a function of θ_0 and θ_1 . Which of the

following statements are true? (Check all that apply.)

☐

Setting the learning rate α to be very small is not harmful, and can only speed up the convergence of gradient descent.

☐

If θ_0 and θ_1 are initialized at the global minimum, then one iteration will not change their values.

☐

If the first few iterations of gradient descent cause $f(\theta_0, \theta_1)$ to **increase** rather than decrease, then the most likely cause is that we have set the learning rate α to too large a value.

☐

No matter how θ_0 and θ_1 are initialized, so long as α is sufficiently small, we can safely expect gradient descent to converge to the same solution.

1
point

5. Suppose that for some linear regression problem (say, predicting housing prices as in the lecture), we have some training set, and for our training set we managed to find some θ_0, θ_1 such that $J(\theta_0, \theta_1) = 0$.

Which of the statements below must then be true? (Check all that apply.)



This is not possible: By the definition of $J(\theta_0, \theta_1)$, it is not possible for there to exist

θ_0 and θ_1 so that $J(\theta_0, \theta_1) = 0$



For this to be true, we must have $\theta_0 = 0$ and $\theta_1 = 0$

so that $h_\theta(x) = 0$



For these values of θ_0 and θ_1 that satisfy $J(\theta_0, \theta_1) = 0$,

we have that $h_\theta(x^{(i)}) = y^{(i)}$ for every training example $(x^{(i)}, y^{(i)})$



We can perfectly predict the value of y even for new examples that we have not yet seen.

(e.g., we can perfectly predict prices of even new houses that we have not yet seen.)

☐ I understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account.

Learn more about Coursera's Honor Code

Enter your legal name

Submit Quiz

