

L2: Review of probability and statistics

Probability

- Definition of probability
- Axioms and properties
- Conditional probability
- Bayes theorem

Random variables

- Definition of a random variable
- Cumulative distribution function
- Probability density function
- Statistical characterization of random variables

Random vectors

- Mean vector
- Covariance matrix

The Gaussian random variable


Review of probability theory

Definitions (informal)

- Probabilities are numbers assigned to events that indicate “*how likely*” it is that the event will occur when a random experiment is performed
- A probability law for a random experiment is a rule that assigns probabilities to the events in the experiment
- The sample space S of a random experiment is the set of all possible outcomes


Axioms of probability

- Axiom I: $P[A_i] \geq 0$
- Axiom II: $P[S] = 1$
- Axiom III: $A_i \cap A_j = \emptyset \Rightarrow P[A_i \cup A_j] = P[A_i] + P[A_j]$



Warm-up exercise

- I show you three colored cards
 - One BLUE on both sides
 - One RED on both sides
 - One BLUE on one side, RED on the other



- I shuffle the three cards, then pick one and show you one side only. The side visible to you is **RED**
 - Obviously, the card has to be either A or C, *right?*
- I am willing to bet \$1 that the other side of the card has the same color, and need someone in class to bet another \$1 that it is the other color
 - On the average we will end up even, *right?*
 - Let's try it!

More properties of probability


- $P[A^C] = 1 - P[A]$
- $P[A] \leq 1$
- $P[\emptyset] = 0$
- *given* $\{A_1 \dots A_N\}, \{A_i \cap A_j = \emptyset, \forall ij\} \Rightarrow P[\cup_{k=1}^N A_k] = \sum_{k=1}^N P[A_k]$
- $P[A_1 \cup A_2] = P[A_1] + P[A_2] - P[A_1 \cap A_2]$
- $P[\cup_{k=1}^N A_k] = \sum_{k=1}^N P[A_k] - \sum_{j < k}^N P[A_j \cap A_k] + \dots + (-1)^{N+1} P[A_1 \cap A_2 \dots \cap A_N]$
- $A_1 \subset A_2 \Rightarrow P[A_1] \leq P[A_2]$

Conditional probability

- If A and B are two events, the probability of event A when we already know that event B has occurred is

$$P[A|B] = \frac{P[A \cap B]}{P[B]} \quad \text{if } P[B] > 0$$

- This conditional probability $P[A|B]$ is read:
 - the “conditional probability of A conditioned on B”, or simply
 - the “probability of A given B”
- Interpretation
 - The new evidence “B has occurred” has the following effects
 - The original sample space S (the square) becomes B (the rightmost circle)
 - The event A becomes $A \cap B$
 - $P[B]$ simply re-normalizes the probability of events that occur jointly with B



Theorem of total probability

- Let $B_1, B_2 \dots B_N$ be a partition of S (mutually exclusive that add to S)
- Any event A can be represented as


$$A = A \cap S = A \cap (B_1 \cup B_2 \dots B_N) = (A \cap B_1) \cup (A \cap B_2) \dots (A \cap B_N)$$

- Since $B_1, B_2 \dots B_N$ are mutually exclusive, then

$$P[A] = P[A \cap B_1] + P[A \cap B_2] + \dots + P[A \cap B_N]$$

- and, therefore

$$P[A] = P[A|B_1]P[B_1] + \dots P[A|B_N]P[B_N] = \sum_{k=1}^N P[A|B_k]P[B_k]$$



Bayes theorem

- Assume $\{B_1, B_2 \dots B_N\}$ is a partition of S
- Suppose that event A occurs
- What is the probability of event B_j ?
- Using the definition of conditional probability and the Theorem of total probability we obtain

$$P[B_j|A] = \frac{P[A \cap B_j]}{P[A]} = \frac{P[A|B_j]P[B_j]}{\sum_{k=1}^N P[A|B_k]P[B_k]}$$

- This is known as Bayes Theorem or Bayes Rule, and is (one of) the most useful relations in probability and statistics

Bayes theorem and statistical pattern recognition

- When used for pattern classification, BT is generally expressed as

$$P[\omega_j|x] = \frac{p[x|\omega_j]P[\omega_j]}{\sum_{k=1}^N p[x|\omega_k]P[\omega_k]} = \frac{p[x|\omega_j]P[\omega_j]}{p[x]}$$

- where ω_j is the j -th class (e.g., phoneme) and x is the feature/observation vector (e.g., vector of MFCCs)
- A typical decision rule is to choose class ω_j with highest $P[\omega_j|x]$
 - Intuitively, we choose the class that is more “likely” given observation x
- Each term in the Bayes Theorem has a special name
 - $P[\omega_j]$ prior probability (of class ω_j)
 - $P[\omega_j|x]$ posterior probability (of class ω_j given the observation x)
 - $p[x|\omega_j]$ likelihood (probability of observation x given class ω_j)
 - $p[x]$ normalization constant (does not affect the decision)

Example

- Consider a clinical problem where we need to decide if a patient has a particular medical condition on the basis of an imperfect test
 - Someone with the condition may go undetected (false-negative)
 - Someone free of the condition may yield a positive result (false-positive)
- Nomenclature
 - The true-negative rate $P(\text{NEG} | \neg\text{COND})$ of a test is called its SPECIFICITY
 - The true-positive rate $P(\text{POS} | \text{COND})$ of a test is called its SENSITIVITY
- Problem
 - Assume a population of 10,000 with a 1% prevalence for the condition
 - Assume that we design a test with 98% specificity and 90% sensitivity
 - Assume you take the test, and the result comes out POSITIVE
 - What is the probability that you have the condition?
- Solution
 - Fill in the joint frequency table next slide, or
 - Apply Bayes rule

	TEST IS POSITIVE	TEST IS NEGATIVE	ROW TOTAL
HAS CONDITION	<i>True-positive</i> $P(POS COND)$	<i>False-negative</i> $P(NEG COND)$	
FREE OF CONDITION	<i>False-positive</i> $P(POS \neg COND)$	<i>True-negative</i> $P(NEG \neg COND)$	
COLUMN TOTAL			


	TEST IS POSITIVE	TEST IS NEGATIVE	ROW TOTAL
HAS CONDITION	<i>True-positive</i> $P(POS COND)$ 100×0.90	<i>False-negative</i> $P(NEG COND)$ $100 \times (1 - 0.90)$	100
FREE OF CONDITION	<i>False-positive</i> $P(POS \neg COND)$ $9,900 \times (1 - 0.98)$	<i>True-negative</i> $P(NEG \neg COND)$ $9,900 \times 0.98$	9,900
COLUMN TOTAL	288	9,712	10,000

- Applying Bayes rule

$$\begin{aligned} P[cond|+] &= \\ &= \frac{P[+|cond]P[cond]}{P[+]} = \\ &= \frac{P[+|cond]P[cond]}{P[+|cond]P[cond] + P[+|\neg cond]P[\neg cond]} = \\ &= \frac{0.90 \times 0.01}{0.90 \times 0.01 + (1 - 0.98) \times 0.99} = \\ &= 0.3125 \end{aligned}$$

Random variables

- When we perform a random experiment we are usually interested in some measurement or numerical attribute of the outcome
 - e.g., weights in a population of subjects, execution times when benchmarking CPUs, shape parameters when performing ATR
- These examples lead to the concept of random variable
 - A random variable X is a function that assigns a real number $X(\xi)$ to each outcome ξ in the sample space of a random experiment
 - $X(\xi)$ maps from all possible outcomes in sample space onto the real line
- The function that assigns values to each outcome is fixed and deterministic, i.e., as in the rule “count the number of heads in three coin tosses”
 - Randomness in X is due to the underlying randomness of the outcome ξ of the experiment
- Random variables can be
 - Discrete, e.g., the resulting number after rolling a dice
 - Continuous, e.g., the weight of a sampled individual



Cumulative distribution function (cdf)



- The cumulative distribution function $F_X(x)$ of a random variable X is defined as the probability of the event $\{X \leq x\}$

$$F_X(x) = P[X \leq x] \quad -\infty < x < \infty$$

- Intuitively, $F_X(b)$ is the long-term proportion of times when $X(\xi) \leq b$

- Properties of the cdf

- $0 \leq F_X(x) \leq 1$
- $\lim_{x \rightarrow \infty} F_X(x) = 1$
- $\lim_{x \rightarrow -\infty} F_X(x) = 0$
- $F_X(a) \leq F_X(b)$ if $a \leq b$
- $F_X(b) = \lim_{h \rightarrow 0} F_X(b + h) = F_X(b^+)$



Probability density function (pdf)

- The probability density function $f_X(x)$ of a continuous random variable X , if it exists, is defined as the derivative of $F_X(x)$



$$f_X(x) = \frac{dF_X(x)}{dx}$$


- For discrete random variables, the equivalent to the pdf is the probability mass function

$$f_X(x) = \frac{\Delta F_X(x)}{\Delta x}$$


- Properties

- $f_X(x) > 0$
- $P[a < x < b] = \int_a^b f_X(x) dx$
- $F_X(x) = \int_{-\infty}^x f_X(x) dx$
- $1 = \int_{-\infty}^{\infty} f_X(x) dx$
- $f_X(x|A) = \frac{d}{dx} F_X(x|A)$ where $F_X(x|A) = \frac{P[\{X < x\} \cap A]}{P[A]}$ if $P[A] > 0$





- **What is the probability of somebody weighting 200 lb?**
 - According to the pdf, this is about 0.62
 - This number seems reasonable, right?
- **Now, what is the probability of somebody weighting 124.876 lb?**
 - According to the pdf, this is about 0.43
 - But, intuitively, we know that the probability should be zero (or very, very small)
- **How do we explain this paradox?**
 - The pdf DOES NOT define a probability, but a probability DENSITY!
 - To obtain the actual probability we must integrate the pdf in an interval
 - So we should have asked the question: what is the probability of somebody weighting 124.876 lb plus or minus 2 lb?



- **The probability mass function is a 'true' probability (reason why we call it a 'mass' as opposed to a 'density')**
 - The pmf is indicating that the probability of any number when rolling a fair dice is the same for all numbers, and equal to $1/6$, a very legitimate answer
 - The pmf DOES NOT need to be integrated to obtain the probability (it cannot be integrated in the first place)

Statistical characterization of random variables

- The cdf or the pdf are SUFFICIENT to fully characterize a r.v.
- However, a r.v. can be PARTIALLY characterized with other measures
- Expectation (center of mass of a density)

$$E[X] = \mu = \int_{-\infty}^{\infty} x f_X(x) dx$$

- Variance (spread about the mean)

$$var[X] = \sigma^2 = E[(X - E[X])^2] = \int_{-\infty}^{\infty} (x - \mu)^2 f_X(x) dx$$

- Standard deviation

$$std[X] = \sigma = var[X]^{1/2}$$

- N-th moment

$$E[X^N] = \int_{-\infty}^{\infty} x^N f_X(x) dx$$

Random vectors

- An extension of the concept of a random variable
 - A random vector \underline{X} is a function that assigns a vector of real numbers to each outcome ξ in sample space S
 - We generally denote a random vector by a column vector
- The notions of cdf and pdf are replaced by ‘joint cdf’ and ‘joint pdf’
 - Given random vector $\underline{X} = [x_1, x_2 \dots x_N]^T$ we define the joint cdf as
$$F_{\underline{X}}(\underline{x}) = P_{\underline{X}}[\{X_1 \leq x_1\} \cap \{X_2 \leq x_2\} \dots \{X_N \leq x_N\}]$$
 - and the joint pdf as

$$f_{\underline{X}}(\underline{x}) = \frac{\partial^N F_{\underline{X}}(\underline{x})}{\partial x_1 \partial x_2 \dots \partial x_N}$$

- The term marginal pdf is used to represent the pdf of a subset of all the random vector dimensions
 - A marginal pdf is obtained by integrating out variables that are of no interest
 - e.g., for a 2D random vector $\underline{X} = [x_1, x_2]^T$, the marginal pdf of x_1 is

$$f_{X_1}(x_1) = \int_{x_2=-\infty}^{x_2=+\infty} f_{X_1 X_2}(x_1 x_2) dx_2$$

Statistical characterization of random vectors


- A random vector is also fully characterized by its joint cdf or joint pdf
- Alternatively, we can (partially) describe a random vector with measures similar to those defined for scalar random variables
- Mean vector

$$E[X] = \underline{\mu} = [E[X_1], E[X_2] \dots E[X_N]]^T = [\mu_1, \mu_2, \dots \mu_N]^T$$

- Covariance matrix

$$\begin{aligned} cov[X] &= \Sigma = E \left[\left(\underline{X} - \underline{\mu} \right) \left(\underline{X} - \underline{\mu} \right)^T \right] = \\ &= \begin{bmatrix} E[(x_1 - \mu_1)^2] & \dots & E[(x_1 - \mu_1)(x_N - \mu_N)] \\ \vdots & \ddots & \vdots \\ E[(x_1 - \mu_1)(x_N - \mu_N)] & \dots & E[(x_N - \mu_N)^2] \end{bmatrix} = \\ &= \begin{bmatrix} \sigma_1^2 & \dots & c_{1N} \\ \vdots & \ddots & \vdots \\ c_{1N} & \dots & \sigma_N^2 \end{bmatrix} \end{aligned}$$

- The covariance matrix indicates the tendency of each pair of features (dimensions in a random vector) to vary together, i.e., to co-vary*
- The covariance has several important properties
 - If x_i and x_k tend to increase together, then $c_{ik} > 0$
 - If x_i tends to decrease when x_k increases, then $c_{ik} < 0$
 - If x_i and x_k are uncorrelated, then $c_{ik} = 0$
 - $|c_{ik}| \leq \sigma_i \sigma_k$, where σ_i is the standard deviation of x_i
 - $c_{ii} = \sigma_i^2 = \text{var}[x_i]$
- The covariance terms can be expressed as $c_{ii} = \sigma_i^2$ and $c_{ik} = \rho_{ik} \sigma_i \sigma_k$
 - where ρ_{ik} is called the correlation coefficient




A numerical example

Given the following samples from a 3D distribution

- Compute the covariance matrix
- Generate scatter plots for every pair of vars.
- Can you observe any relationships between the covariance and the scatter plots?
- You may work your solution in the templates below

	Variables (or features)		
Examples	x_1	x_2	x_3
1	2	2	4
2	3	4	6
3	5	4	2
4	6	6	4

Example	x_1	x_2	x_3	$x_1 - \mu_1$	$x_2 - \mu_2$	$x_3 - \mu_3$	$(x_1 - \mu_1)^2$	$(x_2 - \mu_2)^2$	$(x_3 - \mu_3)^2$	$(x_1 - \mu_1)(x_2 - \mu_2)$	$(x_1 - \mu_1)(x_3 - \mu_3)$	$(x_2 - \mu_2)(x_3 - \mu_3)$
1												
2												
3												
4												
Average												





The Normal or Gaussian distribution

- The multivariate Normal distribution $N(\mu, \Sigma)$ is defined as


$$f_X(x) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{1/2}} e^{-\frac{1}{2}(x-\mu)^T \Sigma^{-1} (x-\mu)}$$

- For a single dimension, this expression is reduced to


$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$



The Gaussian Distribution



$$\mathcal{N}(x|\mu, \sigma^2) = \frac{1}{(2\pi\sigma^2)^{1/2}} \exp \left\{ -\frac{1}{2\sigma^2} (x - \mu)^2 \right\}$$






$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\boldsymbol{\Sigma}|^{1/2}} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right\}$$

Example

- Three-class 2D problem with equal priors




$$\begin{aligned}\mu_1 &= [3 \ 2]^T & \mu_2 &= [7 \ 4]^T & \mu_3 &= [2 \ 5]^T \\ \Sigma_1 &= \begin{bmatrix} 2 & \\ & 2 \end{bmatrix} & \Sigma_2 &= \begin{bmatrix} 2 & \\ & 2 \end{bmatrix} & \Sigma_3 &= \begin{bmatrix} 2 & \\ & 2 \end{bmatrix}\end{aligned}$$



Example

- Three-class 2D problem with equal priors




$$\begin{aligned}\mu_1 &= [3 \ 2]^T & \mu_2 &= [5 \ 4]^T & \mu_3 &= [2 \ 5]^T \\ \Sigma_1 &= \begin{bmatrix} 1 & \\ & 2 \end{bmatrix} & \Sigma_2 &= \begin{bmatrix} 1 & \\ & 2 \end{bmatrix} & \Sigma_3 &= \begin{bmatrix} 1 & \\ & 2 \end{bmatrix}\end{aligned}$$



Example

- Three-class 2D problem with equal priors


$$\begin{array}{lll} \mu_1 = [3 \ 2]^T & \mu_2 = [5 \ 4]^T & \mu_3 = [2 \ 5]^T \\ \Sigma_1 = \begin{bmatrix} 1 & .7 \\ .7 & 2 \end{bmatrix} & \Sigma_2 = \begin{bmatrix} 1 & .7 \\ .7 & 2 \end{bmatrix} & \Sigma_3 = \begin{bmatrix} 1 & .7 \\ .7 & 2 \end{bmatrix} \end{array}$$






Example

- Three-class 2D problem with equal priors

$$\begin{aligned}\mu_1 &= [3 \ 2]^T & \mu_2 &= [5 \ 4]^T & \mu_3 &= [2 \ 5]^T \\ \Sigma_1 &= \begin{bmatrix} .5 & \\ & .5 \end{bmatrix} & \Sigma_2 &= \begin{bmatrix} 1 & \\ & 1 \end{bmatrix} & \Sigma_3 &= \begin{bmatrix} 2 & \\ & 2 \end{bmatrix}\end{aligned}$$






Zoom out




Example

- Three-class 2D problem with equal priors

$$\begin{aligned}\mu_1 &= [3 \ 2]^T & \mu_2 &= [5 \ 4]^T & \mu_3 &= [3 \ 4]^T \\ \Sigma_1 &= \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} & \Sigma_2 &= \begin{bmatrix} 1 & -1 \\ -1 & 7 \end{bmatrix} & \Sigma_3 &= \begin{bmatrix} .5 & .5 \\ .5 & 3 \end{bmatrix}\end{aligned}$$



Zoom
out



- Gaussian distributions are very popular since
 - Parameters (μ, Σ) uniquely characterize the normal distribution
 - If all variables x_i are uncorrelated ($E[x_i x_k] = E[x_i]E[x_k]$), then
 - Variables are also independent ($P[x_i x_k] = P[x_i]P[x_k]$), and
 - Σ is diagonal, with the individual variances in the main diagonal
 - Central Limit Theorem (next slide)
 - The marginal and conditional densities are also Gaussian
 - Any linear transformation of any N jointly Gaussian rv's results in N rv's that are also Gaussian
 - For $X = [X_1 X_2 \dots X_N]^T$ jointly Gaussian, and $A_{N \times N}$ invertible, then $Y = AX$ is also jointly Gaussian

$$f_Y(y) = \frac{f_X(A^{-1}y)}{|A|}$$

Central Limit Theorem

- Given any distribution with a mean μ and variance σ^2 , the sampling distribution of the mean approaches a normal distribution with mean μ and variance σ^2/N as the sample size N increases
 - No matter what the shape of the original distribution is, the sampling distribution of the mean approaches a normal distribution
 - N is the sample size used to compute the mean, not the overall number of samples in the data
- Example: 500 experiments are performed using a uniform distribution
 - $N = 1$
 - One sample is drawn from the distribution and its mean is recorded (500 times)
 - The histogram resembles a uniform distribution, as one would expect
 - $N = 4$
 - Four samples are drawn and the mean of the four samples is recorded (500 times)
 - The histogram starts to look more Gaussian
 - As N grows, the shape of the histograms resembles a Normal distribution more closely

