| Parameter | Norm | Range | Patient value | Diagnosis |
|---|---|---|---|---|
| FH to SN(SN validity) | 7 | 6  8 | 13.2 | |
| Saddle Angle | 129 | 124  134 | 123.4 | |
| Articular Angle | 143 | 137  149 | 146.9 | |
| Gonial Angle | 123 | 128.5  117.5 | 132.4 | |
| Sum Of posterior Angles | 395 | | 402.7 | Long face |
| Y axis | 66 | 63  71 | 73 | |
| Inclination angle | 85 | | 88 | Anteinclined Maxilla |
| Basal angle(PP-GoMe) | 25 | | 42.9 | |
| SN-GoMe | 34 | 28.5  39.5 | 42.7 | |
| S-Go:N-Me (Jaraback Index) | | 62% - 65% | 58 | Long face |
| Occlusal Plane-SN | 14 | | 19.5 | |
| SNA | 82 | 80-84 | 76 | Maxillary Retrognathism |
| SNB | 80 | 78-82 | 81.7 | |
| ANB | 2 (3 for Iranian) | | -5.6 | Class III |
| SN-Pog | 82 (86 for iranian) | | 81.7 | Retruded Chin |
| U1 to SN | 102 | 100  104 | 115.5 | Proclined maxillary Incisors |
| U1 to NA distance | 4 mm | | -12.4 | Retrusion Incisors |
| U1-PP | 110 | | 115.4 | Proclined maxillary Incisors |
| L1-MeGo | 90 | 87  93 | 86.7 | Retroclined Mandibular Incisors |
| L1 to NB distance | 4 mm | | -7.3 | Retrusion mandibular Incisors |
| Interincisal angle | 135 | 130  150.5 | 115 | |
| Witts | | -1  0 | 16.8 | Class II |
| Ramus Height | 43 mm | | 46.2 mm | |
| Se-N (mm) | | | 49.5 | |
| Mandibular Length | 52.5 | | 63.9 | Increased Mandibular Length |
| Maxillary Length | 35 | | 29.3 | Decreased Maxillary Length |
| Ramus Length (A.M.Schwarz) | 37.5 | | 46.2 | Increased Ramus Length |