| Parameter | Norm | Range | Patient value | Diagnosis |
|---|---|---|---|---|
| FH to SN(SN validity) | 7 | 6  8 | 15.1 | |
| Saddle Angle | 129 | 124  134 | 115.3 | |
| Articular Angle | 143 | 137  149 | 145.7 | |
| Gonial Angle | 123 | 128.5  117.5 | 113.9 | |
| Sum Of posterior Angles | 395 | | 374.9 | Short face |
| Y axis | 66 | 63  71 | 54.4 | |
| Inclination angle | 85 | | 84 | Retroinclined Maxilla |
| Basal angle(PP-GoMe) | 25 | | 13.3 | |
| SN-GoMe | 34 | 28.5  39.5 | 14.9 | |
| S-Go:N-Me (Jaraback Index) | | 62% - 65% | 75 | Short face |
| Occlusal Plane-SN | 14 | | 2.6 | |
| SNA | 82 | 80-84 | 87.7 | Maxillary Retrognathism |
| SNB | 80 | 78-82 | 89.2 | Mandibular Prognathism |
| ANB | 2 (3 for Iranian) | | -1.5 | Class III |
| SN-Pog | 82 (86 for iranian) | | 90.8 | Prominent Chin |
| U1 to SN | 102 | 100  104 | 125.8 | Proclined maxillary Incisors |
| U1 to NA distance | 4 mm | | 7.2 | Protrusion Incisors |
| U1-PP | 110 | | 127.4 | Proclined maxillary Incisors |
| L1-MeGo | 90 | 87  93 | 95.6 | Proclined Mandibular Incisors |
| L1 to NB distance | 4 mm | | 2.6 | Retrusion mandibular Incisors |
| Interincisal angle | 135 | 130  150.5 | 123.7 | |
| Witts | | -1  0 | -5 | Class III |
| Ramus Height | 43 mm | | 64.4 mm | |
| Se-N (mm) | | | 67.1 | |
| Mandibular Length | 70.1 | | 73.2 | Increased Mandibular Length |
| Maxillary Length | 46.7 | | 44.6 | Decreased Maxillary Length |
| Ramus Length (A.M.Schwarz) | 50.1 | | 64.4 | Increased Ramus Length |