| Parameter | Norm | Range | Patient value | Diagnosis |
|-------------------------------|---------------------|-------------|---------------|----------------------------------|
| FH to SN(SN validity) | 7 | 6 8 | 4.3 | · · |
| Saddle Angle | 129 | 124 134 | 121.3 | |
| Articular Angle | 143 | 137 149 | 154 | |
| Gonial Angle | 123 | 128.5 117.5 | 118.4 | |
| Sum Of posterior Angles | 395 | | 393.7 | Short face |
| Y axis | 66 | 63 71 | 69.1 | |
| Inclination angle | 85 | | 83 | Retroinclined Maxilla |
| Basal angle(PP- GoMe) | 25 | | 22.3 | |
| SN-GoMe | 34 | 28.5 39.5 | 33.8 | |
| S-Go:N-Me (Jaraback Index) | | 62% - 65% | 67 | Short face |
| Occlusal Plane-SN | 14 | | 20.6 | |
| SNA | 82 | 80-84 | 81.9 | Maxillary Retrognathism |
| SNB | 80 | 78-82 | 79 | |
| ANB | 2 (3 for Iranian) | | -2.9 | Class III |
| SN-Pog | 82 (86 for iranian) | | 79.7 | Retruded Chin |
| U1 to SN | 102 | 100 104 | 103.3 | |
| U1 to NA distance | 4 mm | | -6.1 | Retrusion Incisors |
| U1-PP | 110 | | 114.7 | Proclined maxillary Incisors |
| L1-MeGo | 90 | 87 93 | 110.6 | Proclined Mandibular Incisors |
| L1 to NB distance | 4 mm | | -9 | Retrusion mandibular Incisors |
| Interincisal angle | 135 | 130 150.5 | 112.3 | |
| Witts | | -1 0 | 7.2 | Class II |
| Ramus Height | 43 mm | | 61.5 mm | |
| Se-N (mm) | | | 68.7 | |
| Mandibular Length | 71.7 | | 82.1 | Increased Mandibular Length |
| Maxillary Length | 47.8 | | 50.7 | Increased Maxillary Length |
| Ramus Length (A.M.Schwarz) | 51.2 | | 61.5 | Increased Ramus Length |