| Parameter | Norm | Range | Patient value | Diagnosis |
|-------------------------------|---------------------|-------------|---------------|--------------------------------|
| FH to SN(SN validity) | 7 | 6 8 | 0.6 | |
| Saddle Angle | 129 | 124 134 | 115.8 | |
| Articular Angle | 143 | 137 149 | 147.3 | |
| Gonial Angle | 123 | 128.5 117.5 | 121.7 | |
| Sum Of posterior Angles | 395 | | 384.8 | Short face |
| Y axis | 66 | 63 71 | 61.7 | |
| Inclination angle | 85 | | 89 | Anteinclined Maxilla |
| Basal angle(PP-GoMe) | 25 | | 16.1 | |
| SN-GoMe | 34 | 28.5 39.5 | 24.8 | |
| S-Go:N-Me (Jaraback Index) | | 62% - 65% | 78 | Short face |
| Occlusal Plane-SN | 14 | | 13.8 | |
| SNA | 82 | 80-84 | 80.6 | Maxillary Retrognathism |
| SNB | 80 | 78-82 | 84.9 | Mandibular Prognathism |
| ANB | 2 (3 for Iranian) | | -4.3 | Class III |
| SN-Pog | 82 (86 for iranian) | | 86.1 | Prominent Chin |
| U1 to SN | 102 | 100 104 | 108 | Proclined maxillary Incisors |
| U1 to NA distance | 4 mm | | 6.1 | Protrusion Incisors |
| U1-PP | 110 | | 116.7 | Proclined maxillary Incisors |
| L1-MeGo | 90 | 87 93 | 91.2 | |
| L1 to NB distance | 4 mm | | 2 | Retrusion mandibular Incisors |
| Interincisal angle | 135 | 130 150.5 | 136 | |
| Witts | | -1 0 | -14.3 | Class III |
| Ramus Height | 43 mm | | 66.9 mm | |
| Se-N (mm) | | | 71.7 | |
| Mandibular Length | 74.7 | | 79.8 | Increased Mandibular Length |
| Maxillary Length | 49.8 | | 49.1 | Decreased Maxillary Length |
| Ramus Length (A.M.Schwarz) | 53.4 | | 66.9 | Increased Ramus Length |