| Parameter                  | Norm                | Range       | Patient value | Diagnosis                          |
|----------------------------|---------------------|-------------|---------------|------------------------------------|
| FH to SN(SN validity)      | 7                   | 6 8         | 5.4           |                                    |
| Saddle Angle               | 129                 | 124 134     | 118           |                                    |
| Articular Angle            | 143                 | 137 149     | 141.9         |                                    |
| Gonial Angle               | 123                 | 128.5 117.5 | 140           |                                    |
| Sum Of posterior<br>Angles | 395                 |             | 399.9         | Long face                          |
| Y axis                     | 66                  | 63 71       | 66.6          |                                    |
| Inclination angle          | 85                  |             | 95            | Anteinclined Maxilla               |
| Basal angle(PP-<br>GoMe)   | 25                  |             | 40.5          |                                    |
| SN-GoMe                    | 34                  | 28.5 39.5   | 40            |                                    |
| Occlusal Plane-SN          | 14                  |             | 18.8          |                                    |
| SNA                        | 82                  | 80-84       | 84.9          | Maxillary<br>Retrognathism         |
| SNB                        | 80                  | 78-82       | 86.7          | Mandibular<br>Prognathism          |
| ANB                        | 2 (3 for Iranian)   |             | 1.8           | Class III                          |
| SN-Pog                     | 82 (86 for iranian) |             | 86.6          | Prominent Chin                     |
| U1 to SN                   | 102                 | 100 104     | 112.3         | Proclined maxillary Incisors       |
| U1 to NA distance          | 4 mm                |             | -3.4          | Retrusion Incisors                 |
| U1-PP                      | 110                 |             | 111.8         | Proclined maxillary Incisors       |
| L1-MeGo                    | 90                  | 87 93       | 70            | Retroclined<br>Mandibular Incisors |
| L1 to NB distance          | 4 mm                |             | 2.3           | Retrusion mandibular Incisors      |
| Interincisal angle         | 135                 | 130 150.5   | 137.8         |                                    |
| Witts                      |                     | -1 0        | 17.8          | Class II                           |
| Ramus Height               | 43 mm               |             | 59.8 mm       |                                    |
| Se-N (mm)                  |                     |             | 61.4          |                                    |
| Mandibular Length          | 64.4                |             | 75.7          | Increased<br>Mandibular Length     |
| Maxillary Length           | 42.9                |             | 43.7          | Increased Maxillary<br>Length      |
| Ramus Length (A.M.Schwarz) | 46                  |             | 59.8          | Increased Ramus<br>Length          |