


Probability, Independence, Bayes Rule

Hayk Aprikyan, Hayk Tarkhanyan

Preliminaries

Recall the set operations:


$A \cup B$



Union $A \cup B$:

All elements that belong to A or B or both


$A \cap B$



Intersection $A \cap B$:

All elements that belong to *both* A and B

A^c



Complement A^c :

All elements that do not belong to A

What is probability?

- Oftentimes, we encounter a situation where we do not know the precise value or outcome of a particular event or circumstance.

What is probability?

- Oftentimes, we encounter a situation where we do not know the precise value or outcome of a particular event or circumstance.
- E.g. predicting the score of a football match, or the outcome of a tossed coin.

What is probability?

- Oftentimes, we encounter a situation where we do not know the precise value or outcome of a particular event or circumstance.
- E.g. predicting the score of a football match, or the outcome of a tossed coin.
- While we cannot tell the exact outcome of such an event, we can still speculate about the likely outcomes (e.g. which outcome is more likely than the others).

What is probability?

- Oftentimes, we encounter a situation where we do not know the precise value or outcome of a particular event or circumstance.
- E.g. predicting the score of a football match, or the outcome of a tossed coin.
- While we cannot tell the exact outcome of such an event, we can still speculate about the likely outcomes (e.g. which outcome is more likely than the others).
- The mathematical notion associated with the likeliness of a particular output to happen is called **probability**.

- A **random** (or **probabilistic**) **experiment** is a situation, where we are uncertain about the result.

Outcomes

- A **random** (or **probabilistic**) **experiment** is a situation, where we are uncertain about the result.
- The possible results of a random experiment are called the **outcomes**.

Outcomes

- A **random** (or **probabilistic**) **experiment** is a situation, where we are uncertain about the result.
- The possible results of a random experiment are called the **outcomes**.
- The set of all outcomes is called the **sample space** of the experiment.

- A **random** (or **probabilistic**) **experiment** is a situation, where we are uncertain about the result.
- The possible results of a random experiment are called the **outcomes**.
- The set of all outcomes is called the **sample space** of the experiment.

We denote the sample space with the letter Ω , so

$$\Omega = \{\text{all possible outcomes}\}$$

Example

A football match is a random experiment, where:

- Outcome is the score of the game.
- For example, one possible outcomes is "Pyunik 2 - 1 Alashkert". One way to denote it is $(2, 1)$.
- The sample space is:

$$\Omega = \{(0, 0); (0, 1); (1, 0); (1, 1); (2, 0); \dots\}$$

Example

A football match is a random experiment, where:

- Outcome is the score of the game.
- For example, one possible outcomes is "Pyunik 2 - 1 Alashkert". One way to denote it is $(2, 1)$.
- The sample space is:

$$\Omega = \{(0, 0); (0, 1); (1, 0); (1, 1); (2, 0); \dots\}$$

Example

Tossing a coin is a random experiment, where:

- Outcomes are **Heads** and **Tails**,
- The sample space is:

$$\Omega = \{\mathbf{Heads}, \mathbf{Tails}\} = \{H, T\}$$

Example

A football match is a random experiment, where:

- Outcome is the score of the game.
- For example, one possible outcomes is "Pyunik 2 - 1 Alashkert". One way to denote it is $(2, 1)$.
- The sample space is:

$$\Omega = \{(0, 0); (0, 1); (1, 0); (1, 1); (2, 0); \dots\}$$

Example

Tossing a coin is a random experiment, where:

- Outcomes are **Heads** and **Tails**,
- The sample space is:

$$\Omega = \{\mathbf{Heads}, \mathbf{Tails}\} = \{H, T\}$$

Informally, we can say that both outcomes are equally likely: 50/50.

Outcomes

Let's observe an example where the sample space contains infinite values.

Outcomes

Let's observe an example where the sample space contains infinite values.

Example

A student waits for the bus in the station. How much will she wait?

- Outcome is the time until the bus arrives.
- Practically, it can be *any* positive value, perhaps under 60 minutes.
E.g. 2.3497 minutes is a possible outcome.
- The sample space is:

$$\Omega = [0, 60]$$

Outcomes

Let's observe an example where the sample space contains infinite values.

Example

A student waits for the bus in the station. How much will she wait?

- Outcome is the time until the bus arrives.
- Practically, it can be *any* positive value, perhaps under 60 minutes.
E.g. 2.3497 minutes is a possible outcome.
- The sample space is:

$$\Omega = [0, 60]$$

What is the probability that the bus arrives at 20.230911 minutes?

Outcomes

Let's observe an example where the sample space contains infinite values.

Example

A student waits for the bus in the station. How much will she wait?

- Outcome is the time until the bus arrives.
- Practically, it can be *any* positive value, perhaps under 60 minutes.
E.g. 2.3497 minutes is a possible outcome.
- The sample space is:

$$\Omega = [0, 60]$$

What is the probability that the bus arrives at 20.230911 minutes?
How about exactly 20.0 minutes?

Outcomes

Let's observe an example where the sample space contains infinite values.

Example

A student waits for the bus in the station. How much will she wait?

- Outcome is the time until the bus arrives.
- Practically, it can be *any* positive value, perhaps under 60 minutes. E.g. 2.3497 minutes is a possible outcome.
- The sample space is:

$$\Omega = [0, 60]$$

What is the probability that the bus arrives at 20.230911 minutes?
How about exactly 20.0 minutes? The probability is **zero**.

Outcomes

Let's observe an example where the sample space contains infinite values.

Example

A student waits for the bus in the station. How much will she wait?

- Outcome is the time until the bus arrives.
- Practically, it can be *any* positive value, perhaps under 60 minutes. E.g. 2.3497 minutes is a possible outcome.
- The sample space is:

$$\Omega = [0, 60]$$

What is the probability that the bus arrives at 20.230911 minutes?

How about exactly 20.0 minutes? The probability is **zero**.

Whenever Ω is an interval (e.g. $(0, 1)$, $[1, 10]$, $[0, +\infty)$), the probability of each outcome is 0.

(of course, this does not mean that they are impossible – [watch this!](#))

Events

Let's consider rolling a die.

Events

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?
 - Is it a natural number?

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?
 - Is it a natural number?
 - Is it any number?

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?
 - Is it a natural number?
 - Is it any number?
 - Was there no outcome after rolling the die?

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?
 - Is it a natural number?
 - Is it any number?
 - Was there no outcome after rolling the die?

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?
 - Is it a natural number?
 - Is it any number?
 - Was there no outcome after rolling the die?

To formulate the questions which are of interest to us, we represent them by *sets*. For example, the third question can be represented by the set $\{2, 3, 5\}$.

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?
 - Is it a natural number?
 - Is it any number?
 - Was there no outcome after rolling the die?

To formulate the questions which are of interest to us, we represent them by *sets*. For example, the third question can be represented by the set $\{2, 3, 5\}$.

- The sets of outcomes of interest to us are called **events**. The set $\{2, 3, 5\}$ above is an event.

Let's consider rolling a die.

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- Questions we might be interested in are:
 - Is the outcome equal to 4?
 - Is it smaller than 3?
 - Is it a prime number?
 - Is it a natural number?
 - Is it any number?
 - Was there no outcome after rolling the die?

To formulate the questions which are of interest to us, we represent them by *sets*. For example, the third question can be represented by the set $\{2, 3, 5\}$.

- The sets of outcomes of interest to us are called **events**. The set $\{2, 3, 5\}$ above is an event.
- The set of all events is called the **event space** and denoted by \mathcal{F} .

Example

If you roll a die, the event that a number smaller than 3 shows up is the set:

$$A = \{1, 2\} \in \mathcal{F}$$

Example

If you roll a die, the event that a number smaller than 3 shows up is the set:

$$A = \{1, 2\} \in \mathcal{F}$$

The event of having an odd number is:

$$B = \{1, 3, 5\}$$

Example

If you roll a die, the event that a number smaller than 3 shows up is the set:

$$A = \{1, 2\} \in \mathcal{F}$$

The event of having an odd number is:

$$B = \{1, 3, 5\}$$

and the event of an even number is:

$$C = \{2, 4, 6\}$$

Example

If you roll a die, the event that a number smaller than 3 shows up is the set:

$$A = \{1, 2\} \in \mathcal{F}$$

The event of having an odd number is:

$$B = \{1, 3, 5\}$$

and the event of an even number is:

$$C = \{2, 4, 6\}$$

Can A and B happen at the same time?

Example

If you roll a die, the event that a number smaller than 3 shows up is the set:

$$A = \{1, 2\} \in \mathcal{F}$$

The event of having an odd number is:

$$B = \{1, 3, 5\}$$

and the event of an even number is:

$$C = \{2, 4, 6\}$$

Can A and B happen at the same time? How would you denote the event of both A and B happening?

Example

If you roll a die, the event that a number smaller than 3 shows up is the set:

$$A = \{1, 2\} \in \mathcal{F}$$

The event of having an odd number is:

$$B = \{1, 3, 5\}$$

and the event of an even number is:

$$C = \{2, 4, 6\}$$

Can A and B happen at the same time? How would you denote the event of both A and B happening? What about B and C ?

Definition

Two events A and B of the same experiment are called **disjoint** or **mutually exclusive** if $A \cap B = \emptyset$.

In other words, events are disjoint if they cannot occur at the same time.

Definition

Two events A and B of the same experiment are called **disjoint** or **mutually exclusive** if $A \cap B = \emptyset$.

In other words, events are disjoint if they cannot occur at the same time.

Example

When rolling a die, the events $A = \{1, 4\}$ and $B = \{2, 5\}$ are disjoint, while A and $C = \{3, 4, 5\}$ are not.

Example

When waiting for a bus, the events $A = [0, 20]$ and $B = [30, 40]$ are disjoint, but none of them is disjoint with $C = [10, 40]$.

Question

How can we measure the possibilities of different events?

Question

How can we measure the possibilities of different events?

It depends on the problem/experiment itself.

Question

How can we measure the possibilities of different events?

It depends on the problem/experiment itself.

Suppose you roll a fair die. Since each of the outcomes has the same likelihood as the others, you would expect the probability of rolling, say, 3 to be equal to $\frac{1}{6}$:

$$\mathbb{P}(3) = \frac{1}{6}$$

Question

How can we measure the possibilities of different events?

It depends on the problem/experiment itself.

Suppose you roll a fair die. Since each of the outcomes has the same likelihood as the others, you would expect the probability of rolling, say, 3 to be equal to $\frac{1}{6}$:

$$\mathbb{P}(3) = \frac{1}{6}$$

Similarly, each of the outcomes $\{1, 2, 3, 4, 5, 6\}$ also has the probability

$$\mathbb{P}(1) = \mathbb{P}(2) = \cdots = \mathbb{P}(6) = \frac{1}{6}$$

of showing up.

Question

How can we measure the possibilities of different events?

It depends on the problem/experiment itself.

Suppose you roll a fair die. Since each of the outcomes has the same likelihood as the others, you would expect the probability of rolling, say, 3 to be equal to $\frac{1}{6}$:

$$\mathbb{P}(3) = \frac{1}{6}$$

Similarly, each of the outcomes $\{1, 2, 3, 4, 5, 6\}$ also has the probability

$$\mathbb{P}(1) = \mathbb{P}(2) = \dots = \mathbb{P}(6) = \frac{1}{6}$$

of showing up.

Experiments like this (where all outcomes have the same probability of occurring) are called **equiprobable**.

Probability

The probability of an event, for example, $A = \{3, 5, 6\}$, will be

$$\mathbb{P}(A) =$$

Probability

The probability of an event, for example, $A = \{3, 5, 6\}$, will be

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}$$

since A has 3 elements and the total number of outcomes is 6.

In equiprobable case, the probability of any event $A \in \mathcal{F}$ is:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{total number of outcomes}} = \frac{|A|}{|\Omega|}$$

Probability

The probability of an event, for example, $A = \{3, 5, 6\}$, will be

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}$$

since A has 3 elements and the total number of outcomes is 6.

In equiprobable case, the probability of any event $A \in \mathcal{F}$ is:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{total number of outcomes}} = \frac{|A|}{|\Omega|}$$

Example

Now we roll two fair dice. Our sample space will be

$$\Omega = \{(x, y) \mid 1 \leq x, y \leq 6\}.$$

Assuming that all of 36 outcomes are equally likely to show up, the probability of each (x, y) outcome is:

Probability

The probability of an event, for example, $A = \{3, 5, 6\}$, will be

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}$$

since A has 3 elements and the total number of outcomes is 6.

In equiprobable case, the probability of any event $A \in \mathcal{F}$ is:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{total number of outcomes}} = \frac{|A|}{|\Omega|}$$

Example

Now we roll two fair dice. Our sample space will be

$$\Omega = \{(x, y) \mid 1 \leq x, y \leq 6\}.$$

Assuming that all of 36 outcomes are equally likely to show up, the probability of each (x, y) outcome is:

$$\mathbb{P}((x, y)) = \frac{1}{36} \quad \text{for any } (x, y) \in \Omega$$

Question

What if instead of rolling a die, we go outside, choose a random person and check if they are left-handed? Would the probability still be $\frac{1}{2}$?

Question

What if instead of rolling a die, we go outside, choose a random person and check if they are left-handed? Would the probability still be $\frac{1}{2}$?

Since only about 10% of people are left-handed, it is 9 times less likely to pick a lefty than a righty, hence the probability is about $\frac{1}{10}$, not $\frac{1}{2}$.

Question

What if instead of rolling a die, we go outside, choose a random person and check if they are left-handed? Would the probability still be $\frac{1}{2}$?

Since only about 10% of people are left-handed, it is 9 times less likely to pick a lefty than a righty, hence the probability is about $\frac{1}{10}$, not $\frac{1}{2}$.

In this case, the experiment was not equiprobable. Instead, we had 2 outcomes, say,

$$\Omega = \{L, R\}$$

Question

What if instead of rolling a die, we go outside, choose a random person and check if they are left-handed? Would the probability still be $\frac{1}{2}$?

Since only about 10% of people are left-handed, it is 9 times less likely to pick a lefty than a righty, hence the probability is about $\frac{1}{10}$, not $\frac{1}{2}$.

In this case, the experiment was not equiprobable. Instead, we had 2 outcomes, say,

$$\Omega = \{L, R\}$$

with their probabilities being

$$\mathbb{P}(L) = \frac{1}{10}, \quad \mathbb{P}(R) = \frac{9}{10}$$

Question

What if instead of rolling a die, we go outside, choose a random person and check if they are left-handed? Would the probability still be $\frac{1}{2}$?

Since only about 10% of people are left-handed, it is 9 times less likely to pick a lefty than a righty, hence the probability is about $\frac{1}{10}$, not $\frac{1}{2}$.

In this case, the experiment was not equiprobable. Instead, we had 2 outcomes, say,

$$\Omega = \{L, R\}$$

with their probabilities being

$$\mathbb{P}(L) = \frac{1}{10}, \quad \mathbb{P}(R) = \frac{9}{10}$$

As it shows, probability can also be not equiprobable.

Probability

So in general, in different problems the *probability measure* \mathbb{P} can be different, and it depends on the specifics of the problem how it is computed.

There are, however, three properties which the probability measure \mathbb{P} always satisfies:

- 1 $\mathbb{P}(A) \geq 0$ for any event $A \in \mathcal{F}$,

Probability

So in general, in different problems the *probability measure* \mathbb{P} can be different, and it depends on the specifics of the problem how it is computed.

There are, however, three properties which the probability measure \mathbb{P} always satisfies:

- 1 $\mathbb{P}(A) \geq 0$ for any event $A \in \mathcal{F}$,
- 2 $\mathbb{P}(\Omega) = 1$,

Probability

So in general, in different problems the *probability measure* \mathbb{P} can be different, and it depends on the specifics of the problem how it is computed.

There are, however, three properties which the probability measure \mathbb{P} always satisfies:

- 1 $\mathbb{P}(A) \geq 0$ for any event $A \in \mathcal{F}$,
- 2 $\mathbb{P}(\Omega) = 1$,
- 3 For any disjoint events A_1, A_2, \dots ,

$$\mathbb{P}(A_1 \cup A_2 \cup \dots) = \mathbb{P}(A_1) + \mathbb{P}(A_2) + \dots$$

i.e.

$$\mathbb{P}\left(\bigcup_n A_n\right) = \sum_n \mathbb{P}(A_n)$$

Other useful properties

- 1 For any event A , $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$

Other useful properties

- 1 For any event A , $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- 2 $\mathbb{P}(\emptyset) = 0$

Other useful properties

- 1 For any event A , $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- 2 $\mathbb{P}(\emptyset) = 0$
- 3 If $A \subset B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$

Other useful properties

- 1 For any event A , $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- 2 $\mathbb{P}(\emptyset) = 0$
- 3 If $A \subset B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$
- 4 For any (maybe *not* disjoint) events A_1, A_2, \dots :

$$\mathbb{P}\left(\bigcup_n A_n\right) \leq \sum_n \mathbb{P}(A_n)$$

Other useful properties

- ① For any event A , $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- ② $\mathbb{P}(\emptyset) = 0$
- ③ If $A \subset B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$
- ④ For any (maybe *not* disjoint) events A_1, A_2, \dots :

$$\mathbb{P}\left(\bigcup_n A_n\right) \leq \sum_n \mathbb{P}(A_n)$$

- ⑤ **(Inclusion-Exclusion Rule)** For any events A and B :

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$

Other useful properties

- ① For any event A , $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- ② $\mathbb{P}(\emptyset) = 0$
- ③ If $A \subset B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$
- ④ For any (maybe *not* disjoint) events A_1, A_2, \dots :

$$\mathbb{P}\left(\bigcup_n A_n\right) \leq \sum_n \mathbb{P}(A_n)$$

- ⑤ **(Inclusion-Exclusion Rule)** For any events A and B :

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$


Other useful properties

- 1 For any event A , $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- 2 $\mathbb{P}(\emptyset) = 0$
- 3 If $A \subset B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$
- 4 For any (maybe *not* disjoint) events A_1, A_2, \dots :

$$\mathbb{P}\left(\bigcup_n A_n\right) \leq \sum_n \mathbb{P}(A_n)$$

- 5 **(Inclusion-Exclusion Rule)** For any events A and B :

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$



Conditional Probability

We roll a fair die. Consider these two problems:

Conditional Probability

We role a fair die. Consider these two problems:

- a. What is the probability that the outcome is even?



Conditional Probability

We role a fair die. Consider these two problems:

- a. What is the probability that the outcome is even?



Since there are 6 possible outcomes, out of which only 3 are even,

$$A = \{2, 4, 6\}, \quad \mathbb{P}(A) = \frac{1}{2}$$

Conditional Probability

We role a fair die. Consider these two problems:

- a. What is the probability that the outcome is even?



Since there are 6 possible outcomes, out of which only 3 are even,

$$A = \{2, 4, 6\}, \quad \mathbb{P}(A) = \frac{1}{2}$$

- b. What is the probability that the outcome is even, given that we know that it is prime?



Conditional Probability

We role a fair die. Consider these two problems:

- a. What is the probability that the outcome is even?



Since there are 6 possible outcomes, out of which only 3 are even,

$$A = \{2, 4, 6\}, \quad \mathbb{P}(A) = \frac{1}{2}$$

- b. What is the probability that the outcome is even, given that we know that it is prime?



Here we know that the outcome **cannot be** 1, 4 or 6.

Conditional Probability

We role a fair die. Consider these two problems:

- a. What is the probability that the outcome is even?



Since there are 6 possible outcomes, out of which only 3 are even,

$$A = \{2, 4, 6\}, \quad \mathbb{P}(A) = \frac{1}{2}$$

- b. What is the probability that the outcome is even, given that we know that it is prime?



Here we know that the outcome **cannot be** 1, 4 or 6. Therefore we are left with only three possible outcomes $B = \{2, 3, 5\}$, out of which only 2 is even.

Conditional Probability

We role a fair die. Consider these two problems:

- a. What is the probability that the outcome is even?



Since there are 6 possible outcomes, out of which only 3 are even,

$$A = \{2, 4, 6\}, \quad \mathbb{P}(A) = \frac{1}{2}$$

- b. What is the probability that the outcome is even, given that we know that it is prime?



Here we know that the outcome **cannot be** 1, 4 or 6. Therefore we are left with only three possible outcomes $B = \{2, 3, 5\}$, out of which only 2 is even. So in this case the probability of being even is $\frac{1}{3}$.


Conditional Probability

Definition

For any events A and B (if $\mathbb{P}(B) \neq 0$), the following number:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

is called the **conditional probability** of A **given** B (or the probability of A under the condition of B).




Conditional Probability

Definition

For any events A and B (if $\mathbb{P}(B) \neq 0$), the following number:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

is called the **conditional probability** of A **given** B
(or the probability of A under the condition of B).



more information
is given


Conditional Probability

Definition

For any events A and B (if $\mathbb{P}(B) \neq 0$), the following number:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

is called the **conditional probability** of A **given** B (or the probability of A under the condition of B).




Conditional Probability

Definition

For any events A and B (if $\mathbb{P}(B) \neq 0$), the following number:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

is called the **conditional probability** of A **given** B
(or the probability of A under the condition of B).



In our problem, if A means being even and B means being prime, we had:

$$\mathbb{P}(\text{even} \mid \text{given that prime}) = \mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{1/6}{3/6} = \frac{1}{3}$$

Conditional Probability

Example

Suppose we roll two fair dice. What is the probability that the first one is 2, given that their sum is no greater than 5?

Conditional Probability

Example

Suppose we roll two fair dice. What is the probability that the first one is 2, given that their sum is no greater than 5?

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

Usually, there would be 36 total outcomes, but since we *know* the sum is 5, they are only 10 possible outcomes left. Out of them only three outcomes ("2-1", "2-2" and "2-3") are desired. So the probability is $\frac{3}{10}$.

Question

In some university, $3/5$ of all students are women and the rest are men. It is known that 15% of men are over left-handed, while only 10% of women are. If you choose a random student, what is the probability of the student being left-handed?


Question

In some university, $3/5$ of all students are women and the rest are men. It is known that 15% of men are over left-handed, while only 10% of women are. If you choose a random student, what is the probability of the student being left-handed?

Let B denote the event that the randomly selected student is a woman, and B^c that he is a man.


With another letter, say A , let us denote the event of being left-handed.

Law of Total Probability




$$\mathbb{P}(A) =$$

Law of Total Probability




$$\mathbb{P}(A) =$$

Law of Total Probability



$$\mathbb{P}(A) = \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^c) =$$

Law of Total Probability



$$\mathbb{P}(A) = \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^c) = \mathbb{P}(B) \cdot \mathbb{P}(A|B) + \mathbb{P}(B^c) \cdot \mathbb{P}(A|B^c)$$

Law of Total Probability


What we get is known as the Law of Total Probability:

Theorem

If A and B are some events such that $\mathbb{P}(B) \neq 0$, then

$$\mathbb{P}(A) = \mathbb{P}(B) \cdot \mathbb{P}(A|B) + \mathbb{P}(B^c) \cdot \mathbb{P}(A|B^c)$$

Law of Total Probability



We can also generalize it to the case of three or more subgroups:

Theorem

If B_1, B_2, \dots, B_n are some disjoint events such that $A \subset \bigcup_{k=1}^n B_k$, then

$$\mathbb{P}(A) = \mathbb{P}(B_1) \cdot \mathbb{P}(A|B_1) + \dots + \mathbb{P}(B_n) \cdot \mathbb{P}(A|B_n)$$

Law of Total Probability

Example

There are 52 cards in a deck. One of the cards is randomly removed from the deck, 51 are left. What is the probability that if we take a card, it will be a diamond?

Law of Total Probability

Example

There are 52 cards in a deck. One of the cards is randomly removed from the deck, 51 are left. What is the probability that if we take a card, it will be a diamond?

If we denote by A the probability of taking a diamond, and by B the probability that the removed card was a diamond, then either:

Law of Total Probability

Example

There are 52 cards in a deck. One of the cards is randomly removed from the deck, 51 are left. What is the probability that if we take a card, it will be a diamond?

If we denote by A the probability of taking a diamond, and by B the probability that the removed card was a diamond, then either:

- the taken card was a diamond, and there are 12 left,
- the taken card wasn't a diamond, and there are 13 left.

Law of Total Probability

Example

There are 52 cards in a deck. One of the cards is randomly removed from the deck, 51 are left. What is the probability that if we take a card, it will be a diamond?

If we denote by A the probability of taking a diamond, and by B the probability that the removed card was a diamond, then either:

- the taken card was a diamond, and there are 12 left,
- the taken card wasn't a diamond, and there are 13 left.

The total probability will be:

$$\mathbb{P}(A) = \mathbb{P}(B) \cdot \mathbb{P}(A|B) + \mathbb{P}(B^c) \cdot \mathbb{P}(A|B^c)$$

$$\mathbb{P}(A) = \frac{13}{52} \cdot \frac{12}{51} + \frac{39}{52} \cdot \frac{13}{51} = \frac{1}{4}$$

Bayes Rule

One more simple yet powerful tool is the so called Bayes Rule:

Theorem

If A and B are some events (with non-zero probabilities), then

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B)\mathbb{P}(B)}{\mathbb{P}(A)}$$

Bayes Rule

One more simple yet powerful tool is the so called Bayes Rule:

Theorem

If A and B are some events (with non-zero probabilities), then

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B)\mathbb{P}(B)}{\mathbb{P}(A)}$$

Example

In nature, dangerous fires are rare (about 1% chance) but smoke is fairly common (10% chance) due to barbecues, and 90% of dangerous fires make smoke.

Bayes Rule

One more simple yet powerful tool is the so called Bayes Rule:

Theorem

If A and B are some events (with non-zero probabilities), then

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B)\mathbb{P}(B)}{\mathbb{P}(A)}$$

Example

In nature, dangerous fires are rare (about 1% chance) but smoke is fairly common (10% chance) due to barbecues, and 90% of dangerous fires make smoke. You see a smoke. What is the probability that there is dangerous fire?

Bayes Rule

One more simple yet powerful tool is the so called Bayes Rule:

Theorem

If A and B are some events (with non-zero probabilities), then

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B)\mathbb{P}(B)}{\mathbb{P}(A)}$$

Example

In nature, dangerous fires are rare (about 1% chance) but smoke is fairly common (10% chance) due to barbecues, and 90% of dangerous fires make smoke. You see a smoke. What is the probability that there is dangerous fire?

Let F denote dangerous fire, S smoke. Then:

$$\mathbb{P}(F|S) = \frac{\mathbb{P}(F) \cdot \mathbb{P}(S|F)}{\mathbb{P}(S)} = \frac{1}{100} \cdot \frac{90}{100} : \frac{10}{100} = 0.09$$

Independence

Now let's consider another simple experiment: flipping a fair coin and rolling a six-sided die. The probability of getting **Heads** on the coin flip is independent of the probability of rolling a specific number on the die, as these two events *do not affect* each other.

Independence

Now let's consider another simple experiment: flipping a fair coin and rolling a six-sided die. The probability of getting **Heads** on the coin flip is independent of the probability of rolling a specific number on the die, as these two events *do not affect* each other.

Definition

Events A and B of the same experiment are called **independent** if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

Independence

Now let's consider another simple experiment: flipping a fair coin and rolling a six-sided die. The probability of getting **Heads** on the coin flip is independent of the probability of rolling a specific number on the die, as these two events *do not affect* each other.

Definition

Events A and B of the same experiment are called **independent** if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

or, equivalently,

$$\mathbb{P}(A|B) = \mathbb{P}(A).$$

Independence

Now let's consider another simple experiment: flipping a fair coin and rolling a six-sided die. The probability of getting **Heads** on the coin flip is independent of the probability of rolling a specific number on the die, as these two events *do not affect* each other.

Definition

Events A and B of the same experiment are called **independent** if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

or, equivalently,

$$\mathbb{P}(A|B) = \mathbb{P}(A).$$

The events are called **dependent** if they are not independent.

Independence

Now let's consider another simple experiment: flipping a fair coin and rolling a six-sided die. The probability of getting **Heads** on the coin flip is independent of the probability of rolling a specific number on the die, as these two events *do not affect* each other.

Definition

Events A and B of the same experiment are called **independent** if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

or, equivalently,

$$\mathbb{P}(A|B) = \mathbb{P}(A).$$

The events are called **dependent** if they are not independent.

This definition makes sense because it means that the probability of both A and B occurring together is simply the product of their individual probabilities: the outcome of one event has no effect on the other.

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) =$$

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2},$$

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).

Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}, \quad \mathbb{P}(B) =$$

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}, \quad \mathbb{P}(B) = \frac{3}{6} = \frac{1}{2}$$

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}, \quad \mathbb{P}(B) = \frac{3}{6} = \frac{1}{2}$$

$$\mathbb{P}(A \cap B) =$$

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}, \quad \mathbb{P}(B) = \frac{3}{6} = \frac{1}{2}$$

$$\mathbb{P}(A \cap B) = \frac{9}{36} = \frac{1}{4}$$

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}, \quad \mathbb{P}(B) = \frac{3}{6} = \frac{1}{2}$$

$$\mathbb{P}(A \cap B) = \frac{9}{36} = \frac{1}{4} = \frac{1}{2} \cdot \frac{1}{2}$$

Independence

	1	2	3	4	5	6
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

Example

Suppose we are rolling two fair dice (for the last time).
Event A = the first die is even, B = the second die is odd.

$$\mathbb{P}(A) = \frac{3}{6} = \frac{1}{2}, \quad \mathbb{P}(B) = \frac{3}{6} = \frac{1}{2}$$

$$\mathbb{P}(A \cap B) = \frac{9}{36} = \frac{1}{4} = \frac{1}{2} \cdot \frac{1}{2} = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- 1 If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- 1 If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)
- 2 If A and B are independent events, then

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- ① If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)
- ② If A and B are independent events, then
 - A and B^c

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- ① If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)
- ② If A and B are independent events, then
 - A and B^c
 - A^c and B

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- ① If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)
- ② If A and B are independent events, then
 - A and B^c
 - A^c and B
 - A^c and B^c

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- ① If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)
- ② If A and B are independent events, then
 - A and B^c
 - A^c and B
 - A^c and B^c

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- ① If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)
- ② If A and B are independent events, then
 - A and B^c
 - A^c and B
 - A^c and B^care also independent.

Independence

Suppose A and B are some events with $\mathbb{P}(A) \neq 0$ and $\mathbb{P}(B) \neq 0$.

Properties

- ① If A and B are independent, they cannot be disjoint.
(Hence disjoint events are dependent.)
- ② If A and B are independent events, then
 - A and B^c
 - A^c and B
 - A^c and B^care also independent.

When solving a problem, principles like total probability, Bayes rule, and independence help us a lot. Let's explore one more useful problem solving technique.

Geometric Probability

When thinking about probabilities of some events, it is often easier for us to think about them geometrically.

Geometric Probability

When thinking about probabilities of some events, it is often easier for us to think about them geometrically.

Example


Your bus is coming at a random time between 12 pm and 1 pm. If you show up at 12:30 pm, how likely are you to catch the bus?

Geometric Probability

When thinking about probabilities of some events, it is often easier for us to think about them geometrically.

Example

Your bus is coming at a random time between 12 pm and 1 pm. If you show up at 12:30 pm, how likely are you to catch the bus?




If we draw the hours on a number line and mark the time points for which we will miss or catch the bus, we see that the *length* of the segment for catching the bus is half the total length:

$$\mathbb{P}(\text{catching the bus}) = \frac{30}{30 + 30} = \frac{1}{2}$$

Geometric Probability

Example

A selection is to be made between points G and J as seen below




The probability that selection falls in HF is

Geometric Probability

Example

A selection is to be made between points G and J as seen below




The probability that selection falls in HF is $\frac{HF}{GH} = \frac{5}{11}$.

Geometric Probability

Example

A selection is to be made between points G and J as seen below



The probability that selection falls in HF is $\frac{HF}{GH} = \frac{5}{11}$.

So in general, we draw the sample space and the set of desired outcomes as lines or line segments, and divide the length of the "desired outcomes" by the length of the "sample space" to get the probability.

Geometric Probability


Imagine now that we are playing darts with two circles, and the smaller circle has two times smaller radius than the bigger one.



A dart is thrown at a random. What is the probability that it lands in the smaller circle?

Geometric Probability

Imagine now that we are playing darts with two circles, and the smaller circle has two times smaller radius than the bigger one.



A dart is thrown at a random. What is the probability that it lands in the smaller circle?

In this case, since the sample space and the set of desired outcomes is 2-dimensional, we should divide the *area* of the smaller circle by the *area* of the bigger circle.

The probability that the dart falls in the smaller circle, is $\frac{\pi r^2}{\pi (2r)^2} = \frac{1}{4}$.

Geometric Probability

One of the most powerful uses of geometric probability is applying it to problems that are not inherently geometric.

Geometric Probability

One of the most powerful uses of geometric probability is applying it to problems that are not inherently geometric.

Example

Both the bus and you get to the bus stop at random times between 12 pm and 1 pm. When the bus arrives, it waits for 5 minutes before leaving. When you arrive, you wait for 20 minutes before leaving if the bus doesn't come. What is the probability that you catch the bus?

Geometric Probability

One of the most powerful uses of geometric probability is applying it to problems that are not inherently geometric.

Example

Both the bus and you get to the bus stop at random times between 12 pm and 1 pm. When the bus arrives, it waits for 5 minutes before leaving. When you arrive, you wait for 20 minutes before leaving if the bus doesn't come. What is the probability that you catch the bus?

Let b denote the time that the bus arrives (after 12 pm), and y , the time you arrive. The set of all possible outcomes will be $[0, 60] \times [0, 60]$.


Geometric Probability

One of the most powerful uses of geometric probability is applying it to problems that are not inherently geometric.

Example


Both the bus and you get to the bus stop at random times between 12 pm and 1 pm. When the bus arrives, it waits for 5 minutes before leaving. When you arrive, you wait for 20 minutes before leaving if the bus doesn't come. What is the probability that you catch the bus?

Let b denote the time that the bus arrives (after 12 pm), and y , the time you arrive. The set of all possible outcomes will be $[0, 60] \times [0, 60]$.




Geometric Probability

Then, we need to determine the region of "success"; that is, the points where we catch the bus. Since the bus will wait for 5 minutes, you need to arrive within 5 minutes of the bus' arrival, so $y \leq b + 5$.




Geometric Probability

Then, we need to determine the region of "success"; that is, the points where we catch the bus. Since the bus will wait for 5 minutes, you need to arrive within 5 minutes of the bus' arrival, so $y \leq b + 5$.




However, you only wait for 20 minutes, so you can't arrive more than 20 minutes before the bus, so $y \geq b - 20$.




Geometric Probability

Combining our two conditions, we can draw the region of success:




Geometric Probability

Combining our two conditions, we can draw the region of success:




Now, we just need to find the area of this region. A simple method is to find the remaining area, and then subtract that from the total area:




Geometric Probability

Combining our two conditions, we can draw the region of success:



Now, we just need to find the area of this region. A simple method is to find the remaining area, and then subtract that from the total area:



$$\mathbb{P}(\text{catching the bus}) = \frac{60^2 - \frac{55^2}{2} - \frac{40^2}{2}}{60^2} = \frac{103}{288}$$