

Eigenvalues, Eigendecomposition, SVD

Hayk Aprikyan, Hayk Tarkhanyan

Motivation

Some matrices are easier to understand than others.

Motivation

Some matrices are easier to understand than others.

Question

What does the matrix $A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ do to the vectors of \mathbb{R}^2 ?

Motivation

Some matrices are easier to understand than others.

Question

What does the matrix $A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ do to the vectors of \mathbb{R}^2 ?

Well, it just stretches the vectors in x -direction by 2 times and in y -direction by 3 times – no rotation, no shearing, easy peasy.

Motivation

Some matrices are easier to understand than others.

Question

What does the matrix $A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ do to the vectors of \mathbb{R}^2 ?

Well, it just stretches the vectors in x -direction by 2 times and in y -direction by 3 times – no rotation, no shearing, easy peasy.

In general, a diagonal matrix stretches the space in i^{th} direction by d_i times:

$$\begin{bmatrix} d_1 & 0 & \cdots & 0 \\ 0 & d_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_n \end{bmatrix}$$

Let's see if we can make any matrix look like this.

Eigenvalues and Eigenvectors

Our secret weapon will be the *eigenvalues* and *eigenvectors* of a matrix.

Eigenvalues and Eigenvectors

Our secret weapon will be the *eigenvalues* and *eigenvectors* of a matrix.


Usually, when we multiply a vector by a square matrix, both its length and direction change.

Eigenvalues and Eigenvectors

Our secret weapon will be the *eigenvalues* and *eigenvectors* of a matrix.

Usually, when we multiply a vector by a square matrix, both its length and direction change.

However, sometimes there are very special vectors which **do not change their direction** when being multiplied/transformed by that matrix:



but only get stretched or squished (or flipped) by some factor.

Eigenvalues and Eigenvectors

In other words, when you multiply A with such a special vector \mathbf{v} :

$$A\mathbf{v}$$

Eigenvalues and Eigenvectors

In other words, when you multiply A with such a special vector \mathbf{v} :

$$A\mathbf{v} = \lambda\mathbf{v}$$

you get the same vector \mathbf{v} back, just scaled by some factor λ .

Eigenvalues and Eigenvectors

In other words, when you multiply A with such a special vector \mathbf{v} :

$$A\mathbf{v} = \lambda\mathbf{v}$$

you get the same vector \mathbf{v} back, just scaled by some factor λ .

Such vectors are called the *eigenvectors* of the matrix, and the factor by which they get stretched/squished is called the *eigenvalue*.

Eigenvalues and Eigenvectors

In other words, when you multiply A with such a special vector \mathbf{v} :

$$A\mathbf{v} = \lambda\mathbf{v}$$

you get the same vector \mathbf{v} back, just scaled by some factor λ .

Such vectors are called the *eigenvectors* of the matrix, and the factor by which they get stretched/squished is called the *eigenvalue*.

Definition

If for some number λ and some non-zero vector \mathbf{v}

$$A\mathbf{v} = \lambda\mathbf{v}$$

then we say

- λ is an **eigenvalue** of A ,
- \mathbf{v} is an **eigenvector** of A corresponding to the eigenvalue λ .

Eigenvalues and Eigenvectors

In other words, when you multiply A with such a special vector \mathbf{v} :

$$A\mathbf{v} = \lambda\mathbf{v}$$

you get the same vector \mathbf{v} back, just scaled by some factor λ .

Such vectors are called the *eigenvectors* of the matrix, and the factor by which they get stretched/squished is called the *eigenvalue*.

Definition

If for some number λ and some non-zero vector \mathbf{v}

$$A\mathbf{v} = \lambda\mathbf{v}$$

then we say

- λ is an **eigenvalue** of A ,
- \mathbf{v} is an **eigenvector** of A corresponding to the eigenvalue λ .

Note that λ can be any number (including negative numbers and zero).

Eigenvalues and Eigenvectors

Example

For the matrix $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$ and vector $\mathbf{v} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}$,

$$A\mathbf{v} = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 20 \\ 4 \end{bmatrix} = 4\mathbf{v}$$

so $\lambda = 4$ is an eigenvalue of A with eigenvector \mathbf{v} .

Eigenvalues and Eigenvectors

Example

For the matrix $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$ and vector $\mathbf{v} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}$,

$$A\mathbf{v} = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 20 \\ 4 \end{bmatrix} = 4\mathbf{v}$$

so $\lambda = 4$ is an eigenvalue of A with eigenvector \mathbf{v} .

Question

Can you name another eigenvector of A ?

Eigenvalues and Eigenvectors

Example

For the matrix $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$ and vector $\mathbf{v} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}$,

$$A\mathbf{v} = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 20 \\ 4 \end{bmatrix} = 4\mathbf{v}$$

so $\lambda = 4$ is an eigenvalue of A with eigenvector \mathbf{v} .

Question

Can you name another eigenvector of A ?

If \mathbf{v} is an eigenvector of A , any multiple of \mathbf{v} is also an eigenvector!

$$A(c \cdot \mathbf{v}) = c \cdot (A\mathbf{v}) = c \cdot (4\mathbf{v}) = 4 \cdot (c\mathbf{v})$$

Eigenvalues and Eigenvectors

Example

For the matrix $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$ and vector $\mathbf{v} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}$,

$$A\mathbf{v} = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 20 \\ 4 \end{bmatrix} = 4\mathbf{v}$$

so $\lambda = 4$ is an eigenvalue of A with eigenvector \mathbf{v} .

Question

Can you name another eigenvector of A ?

If \mathbf{v} is an eigenvector of A , any multiple of \mathbf{v} is also an eigenvector!

$$A(c \cdot \mathbf{v}) = c \cdot (A\mathbf{v}) = c \cdot (4\mathbf{v}) = 4 \cdot (c\mathbf{v})$$

Do all matrices have eigenvalues and eigenvectors?

Eigenvalues and Eigenvectors

Note that:

- **Not all matrices have eigenvalues and eigenvectors.**
For example, the rotation matrices do not have any eigenvectors.

Eigenvalues and Eigenvectors

Note that:

- **Not all matrices have eigenvalues and eigenvectors.**
For example, the rotation matrices do not have any eigenvectors.
- In fact, **only some square matrices** have eigenvectors.

Eigenvalues and Eigenvectors

Note that:

- **Not all matrices have eigenvalues and eigenvectors.**
For example, the rotation matrices do not have any eigenvectors.
- In fact, **only some square matrices** have eigenvectors.
- For each fixed eigenvalue, there are **infinitely many eigenvectors** belonging to the same line.

Eigenvalues and Eigenvectors

Note that:

- **Not all matrices have eigenvalues and eigenvectors.**
For example, the rotation matrices do not have any eigenvectors.
- In fact, **only some square matrices** have eigenvectors.
- For each fixed eigenvalue, there are **infinitely many eigenvectors** belonging to the same line.
- It can also be that there are two or more *linearly independent* eigenvectors corresponding to the same eigenvalue.

Eigenvalues and Eigenvectors

Note that:

- **Not all matrices have eigenvalues and eigenvectors.**
For example, the rotation matrices do not have any eigenvectors.
- In fact, **only some square matrices** have eigenvectors.
- For each fixed eigenvalue, there are **infinitely many eigenvectors** belonging to the same line.
- It can also be that there are two or more *linearly independent* eigenvectors corresponding to the same eigenvalue.

Eigenvalues and Eigenvectors

Note that:

- **Not all matrices have eigenvalues and eigenvectors.**
For example, the rotation matrices do not have any eigenvectors.
- In fact, **only some square matrices** have eigenvectors.
- For each fixed eigenvalue, there are **infinitely many eigenvectors** belonging to the same line.
- It can also be that there are two or more *linearly independent* eigenvectors corresponding to the same eigenvalue.

In fact, for each fixed eigenvalue λ , the set of all eigenvectors corresponding to λ , together with the zero vector $\mathbf{0}$, is a vector subspace of \mathbb{R}^n , called the *eigenspace* corresponding to λ :

$$E_\lambda = \{\mathbf{v} \in \mathbb{R}^n \mid A\mathbf{v} = \lambda\mathbf{v}\}$$

Eigenvalues and Eigenvectors

Note that:

- **Not all matrices have eigenvalues and eigenvectors.**
For example, the rotation matrices do not have any eigenvectors.
- In fact, **only some square matrices** have eigenvectors.
- For each fixed eigenvalue, there are **infinitely many eigenvectors** belonging to the same line.
- It can also be that there are two or more *linearly independent* eigenvectors corresponding to the same eigenvalue.

In fact, for each fixed eigenvalue λ , the set of all eigenvectors corresponding to λ , together with the zero vector $\mathbf{0}$, is a vector subspace of \mathbb{R}^n , called the *eigenspace* corresponding to λ :

$$E_\lambda = \{\mathbf{v} \in \mathbb{R}^n \mid A\mathbf{v} = \lambda\mathbf{v}\}$$

The set of all eigenvalues of A is called the *spectrum* of A .

Eigenvalues and Eigenvectors

How can we actually find if a given square matrix has eigenvalues and eigenvectors, and if so, how to find them?

Eigenvalues and Eigenvectors

How can we actually find if a given square matrix has eigenvalues and eigenvectors, and if so, how to find them?

Well, suppose $A \in \mathbb{R}^{n \times n}$ and we want to find $\lambda \in \mathbb{R}$ and $\mathbf{v} \neq \mathbf{0}$ such that:

$$A\mathbf{v} = \lambda\mathbf{v}$$

Eigenvalues and Eigenvectors

How can we actually find if a given square matrix has eigenvalues and eigenvectors, and if so, how to find them?

Well, suppose $A \in \mathbb{R}^{n \times n}$ and we want to find $\lambda \in \mathbb{R}$ and $\mathbf{v} \neq \mathbf{0}$ such that:

$$A\mathbf{v} = \lambda\mathbf{v}$$

$$A\mathbf{v} = \lambda(I\mathbf{v})$$

Eigenvalues and Eigenvectors

How can we actually find if a given square matrix has eigenvalues and eigenvectors, and if so, how to find them?

Well, suppose $A \in \mathbb{R}^{n \times n}$ and we want to find $\lambda \in \mathbb{R}$ and $\mathbf{v} \neq \mathbf{0}$ such that:

$$A\mathbf{v} = \lambda\mathbf{v}$$

$$A\mathbf{v} = \lambda(I\mathbf{v})$$

$$A\mathbf{v} - \lambda I\mathbf{v} = \mathbf{0}$$

Eigenvalues and Eigenvectors

How can we actually find if a given square matrix has eigenvalues and eigenvectors, and if so, how to find them?

Well, suppose $A \in \mathbb{R}^{n \times n}$ and we want to find $\lambda \in \mathbb{R}$ and $\mathbf{v} \neq \mathbf{0}$ such that:

$$A\mathbf{v} = \lambda\mathbf{v}$$

$$A\mathbf{v} = \lambda(I\mathbf{v})$$

$$A\mathbf{v} - \lambda I\mathbf{v} = \mathbf{0}$$

$$(A - \lambda I)\mathbf{v} = \mathbf{0}$$

Eigenvalues and Eigenvectors

How can we actually find if a given square matrix has eigenvalues and eigenvectors, and if so, how to find them?

Well, suppose $A \in \mathbb{R}^{n \times n}$ and we want to find $\lambda \in \mathbb{R}$ and $\mathbf{v} \neq \mathbf{0}$ such that:

$$A\mathbf{v} = \lambda\mathbf{v}$$

$$A\mathbf{v} = \lambda(I\mathbf{v})$$

$$A\mathbf{v} - \lambda I\mathbf{v} = \mathbf{0}$$

$$\underbrace{(A - \lambda I)}_{\text{takes } \mathbf{v} \neq \mathbf{0} \text{ to zero}} \mathbf{v} = \mathbf{0}$$

Eigenvalues and Eigenvectors

How can we actually find if a given square matrix has eigenvalues and eigenvectors, and if so, how to find them?

Well, suppose $A \in \mathbb{R}^{n \times n}$ and we want to find $\lambda \in \mathbb{R}$ and $\mathbf{v} \neq \mathbf{0}$ such that:

$$A\mathbf{v} = \lambda\mathbf{v}$$

$$A\mathbf{v} = \lambda(I\mathbf{v})$$

$$A\mathbf{v} - \lambda I\mathbf{v} = \mathbf{0}$$

$$\underbrace{(A - \lambda I)}_{\text{takes } \mathbf{v} \neq \mathbf{0} \text{ to zero}} \mathbf{v} = \mathbf{0}$$

The matrix $(A - \lambda I)$ squishes down some non-zero vector \mathbf{v} to zero, which means its determinant must be zero:

$$\det(A - \lambda I) = 0$$

Eigenvalues and Eigenvectors

Now here:

$$\det(A - \lambda I) = 0$$

Eigenvalues and Eigenvectors

Now here:

$$\det(A - \lambda I) = 0$$

the expression on the left is a polynomial (where λ is the unknown):

$$A - \lambda I = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{bmatrix}$$

Eigenvalues and Eigenvectors

Now here:

$$\det(A - \lambda I) = 0$$

the expression on the left is a polynomial (where λ is the unknown):

$$A - \lambda I = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{bmatrix}$$

so the determinant is:

$$\det(A - \lambda I) = \dots = \lambda^2 - (a_{11} + a_{22})\lambda + (a_{11}a_{22} - a_{12}a_{21})$$

Eigenvalues and Eigenvectors

Now here:

$$\det(A - \lambda I) = 0$$

the expression on the left is a polynomial (where λ is the unknown):

$$A - \lambda I = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{bmatrix}$$

so the determinant is:

$$\det(A - \lambda I) = \dots = \lambda^2 - (a_{11} + a_{22})\lambda + (a_{11}a_{22} - a_{12}a_{21})$$

In general, for any $n \times n$ matrix A , the expression $\det(A - \lambda I)$ is a polynomial of degree n in λ .

Eigenvalues and Eigenvectors

Now here:

$$\det(A - \lambda I) = 0$$

the expression on the left is a polynomial (where λ is the unknown):

$$A - \lambda I = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{bmatrix}$$

so the determinant is:

$$\det(A - \lambda I) = \dots = \lambda^2 - (a_{11} + a_{22})\lambda + (a_{11}a_{22} - a_{12}a_{21})$$

In general, for any $n \times n$ matrix A , the expression $\det(A - \lambda I)$ is a polynomial of degree n in λ .

We call it the *characteristic polynomial* of A and denote it by $p_A(\lambda)$. The roots of this polynomial are the eigenvalues of A .

Eigenvalues and Eigenvectors

Example

Let's find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$:

Eigenvalues and Eigenvectors

Example

Let's find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$:

$$A - \lambda I =$$

Eigenvalues and Eigenvectors

Example

Let's find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$:

$$A - \lambda I = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} =$$

Eigenvalues and Eigenvectors

Example

Let's find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$:

$$A - \lambda I = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 5 \\ 1 & -1 - \lambda \end{bmatrix}$$

Eigenvalues and Eigenvectors

Example

Let's find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$:

$$A - \lambda I = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 5 \\ 1 & -1 - \lambda \end{bmatrix}$$

The characteristic polynomial is:

$$p_A(\lambda) = \det(A - \lambda I) = (3 - \lambda)(-1 - \lambda) - 5 = \lambda^2 - 2\lambda - 8$$

Eigenvalues and Eigenvectors

Example

Let's find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$:

$$A - \lambda I = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 5 \\ 1 & -1 - \lambda \end{bmatrix}$$

The characteristic polynomial is:

$$p_A(\lambda) = \det(A - \lambda I) = (3 - \lambda)(-1 - \lambda) - 5 = \lambda^2 - 2\lambda - 8$$

the roots of which are $\lambda = 4$ and $\lambda = -2$.

Eigenvalues and Eigenvectors

Example

Let's find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix}$:

$$A - \lambda I = \begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 5 \\ 1 & -1 - \lambda \end{bmatrix}$$

The characteristic polynomial is:

$$p_A(\lambda) = \det(A - \lambda I) = (3 - \lambda)(-1 - \lambda) - 5 = \lambda^2 - 2\lambda - 8$$

the roots of which are $\lambda = 4$ and $\lambda = -2$.

We found the eigenvalues of A ! Now, let's find the corresponding eigenvectors.

Eigenvalues and Eigenvectors

Example

For $\lambda = 4$, we need to find the non-zero solutions of:

$$A\mathbf{v} = 4\mathbf{v}$$

Eigenvalues and Eigenvectors

Example

For $\lambda = 4$, we need to find the non-zero solutions of:

$$A\mathbf{v} = 4\mathbf{v}$$

Denote $\mathbf{v} = [x \ y]^T$, then:

$$\begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 4 \begin{bmatrix} x \\ y \end{bmatrix}$$

Eigenvalues and Eigenvectors

Example

For $\lambda = 4$, we need to find the non-zero solutions of:

$$A\mathbf{v} = 4\mathbf{v}$$

Denote $\mathbf{v} = [x \ y]^T$, then:

$$\begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 4 \begin{bmatrix} x \\ y \end{bmatrix}$$

or equivalently:

$$\begin{bmatrix} 3x + 5y \\ x - y \end{bmatrix} = \begin{bmatrix} 4x \\ 4y \end{bmatrix}$$

Eigenvalues and Eigenvectors

Example

For $\lambda = 4$, we need to find the non-zero solutions of:

$$A\mathbf{v} = 4\mathbf{v}$$

Denote $\mathbf{v} = [x \ y]^T$, then:

$$\begin{bmatrix} 3 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 4 \begin{bmatrix} x \\ y \end{bmatrix}$$

or equivalently:

$$\begin{bmatrix} 3x + 5y \\ x - y \end{bmatrix} = \begin{bmatrix} 4x \\ 4y \end{bmatrix}$$

from which we get $y = \frac{1}{5}x$, i.e. all the vectors on the line $y = \frac{1}{5}x$ are eigenvectors corresponding to $\lambda = 4$.

Example

In other words, the eigenspace E_4 is the line $y = \frac{1}{5}x$ – the vectors on this line do not change their direction when multiplied by A , but only get stretched by 4 times.

Eigenvalues and Eigenvectors

Example

In other words, the eigenspace E_4 is the line $y = \frac{1}{5}x$ – the vectors on this line do not change their direction when multiplied by A , but only get stretched by 4 times.

Similarly we can show that E_{-2} is the line $y = -x$.

Eigenvalues and Eigenvectors

Example

In other words, the eigenspace E_4 is the line $y = \frac{1}{5}x$ – the vectors on this line do not change their direction when multiplied by A , but only get stretched by 4 times.

Similarly we can show that E_{-2} is the line $y = -x$.

Again, what we are concerned with, is the *concept*, not the computation.

Eigenvalues and Eigenvectors

Example

In other words, the eigenspace E_4 is the line $y = \frac{1}{5}x$ – the vectors on this line do not change their direction when multiplied by A , but only get stretched by 4 times.

Similarly we can show that E_{-2} is the line $y = -x$.

Again, what we are concerned with, is the *concept*, not the computation.

Question

What do you think happens to the eigenvalues of a matrix if we transpose it?

Eigenvalues and Eigenvectors

Sometimes, an eigenvalue can appear more than once as a root of the characteristic polynomial.

Eigenvalues and Eigenvectors

Sometimes, an eigenvalue can appear more than once as a root of the characteristic polynomial. For example, the matrix

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 0 & 3 & 0 \\ 0 & 2 & 5 \end{bmatrix}$$

has characteristic polynomial

$$p_A(\lambda) = \underbrace{(\lambda - 3)^2}_{\text{2 times}} \underbrace{(\lambda - 5)}_{\text{1 time}}$$

Eigenvalues and Eigenvectors

Sometimes, an eigenvalue can appear more than once as a root of the characteristic polynomial. For example, the matrix

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 0 & 3 & 0 \\ 0 & 2 & 5 \end{bmatrix}$$

has characteristic polynomial

$$p_A(\lambda) = \underbrace{(\lambda - 3)^2}_{\text{2 times}} \underbrace{(\lambda - 5)}_{\text{1 time}}$$

In this case, we say that eigenvalue $\lambda = 3$ has *algebraic multiplicity* 2, and $\lambda = 5$ has algebraic multiplicity 1.

Eigenvalues and Eigenvectors

As a bonus, we have a beautiful and surprising theorem:

Theorem

The determinant of a matrix is equal to the product of its eigenvalues:

$$\det(A) = \lambda_1 \cdot \lambda_2 \cdot \dots \cdot \lambda_n$$

and the trace of a matrix is equal to the sum of its eigenvalues:

$$\operatorname{tr}(A) = \lambda_1 + \lambda_2 + \dots + \lambda_n$$

Note that the eigenvalues are counted **as many times** as their algebraic multiplicities here.

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Since there are n of them, they form a basis of \mathbb{R}^n .

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Since there are n of them, they form a basis of \mathbb{R}^n . Therefore any vector $\mathbf{x} \in \mathbb{R}^n$ can be expressed as a linear combination of these eigenvectors:

$$\mathbf{x} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n$$

for some coefficients c_1, c_2, \dots, c_n .

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Since there are n of them, they form a basis of \mathbb{R}^n . Therefore any vector $\mathbf{x} \in \mathbb{R}^n$ can be expressed as a linear combination of these eigenvectors:

$$\mathbf{x} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n$$

for some coefficients c_1, c_2, \dots, c_n .

Question

What happens when we multiply \mathbf{x} by A ?

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Since there are n of them, they form a basis of \mathbb{R}^n . Therefore any vector $\mathbf{x} \in \mathbb{R}^n$ can be expressed as a linear combination of these eigenvectors:

$$\mathbf{x} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n$$

for some coefficients c_1, c_2, \dots, c_n .

Question

What happens when we multiply \mathbf{x} by A ?

$$A\mathbf{x} =$$

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Since there are n of them, they form a basis of \mathbb{R}^n . Therefore any vector $\mathbf{x} \in \mathbb{R}^n$ can be expressed as a linear combination of these eigenvectors:

$$\mathbf{x} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n$$

for some coefficients c_1, c_2, \dots, c_n .

Question

What happens when we multiply \mathbf{x} by A ?

$$A\mathbf{x} = c_1 A\mathbf{v}_1 + c_2 A\mathbf{v}_2 + \dots + c_n A\mathbf{v}_n =$$

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Since there are n of them, they form a basis of \mathbb{R}^n . Therefore any vector $\mathbf{x} \in \mathbb{R}^n$ can be expressed as a linear combination of these eigenvectors:

$$\mathbf{x} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n$$

for some coefficients c_1, c_2, \dots, c_n .

Question

What happens when we multiply \mathbf{x} by A ?

$$A\mathbf{x} = c_1 A\mathbf{v}_1 + c_2 A\mathbf{v}_2 + \dots + c_n A\mathbf{v}_n = c_1 \lambda_1 \mathbf{v}_1 + c_2 \lambda_2 \mathbf{v}_2 + \dots + c_n \lambda_n \mathbf{v}_n$$

Eigendecomposition

Suppose now $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix with n different eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (and corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$).

We can prove that since A is symmetric, its eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$$

are orthogonal to each other.

Since there are n of them, they form a basis of \mathbb{R}^n . Therefore any vector $\mathbf{x} \in \mathbb{R}^n$ can be expressed as a linear combination of these eigenvectors:

$$\mathbf{x} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n$$

for some coefficients c_1, c_2, \dots, c_n .

Question

What happens when we multiply \mathbf{x} by A ?

$$A\mathbf{x} = c_1 A\mathbf{v}_1 + c_2 A\mathbf{v}_2 + \dots + c_n A\mathbf{v}_n = c_1 \lambda_1 \mathbf{v}_1 + c_2 \lambda_2 \mathbf{v}_2 + \dots + c_n \lambda_n \mathbf{v}_n$$

It has the same effect as stretching in each direction of \mathbf{v}_i by λ_i times!

Eigendecomposition

To make it more precise, suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix with n eigenvalues and n **linearly independent** eigenvectors.

Eigendecomposition

To make it more precise, suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix with n eigenvalues and n **linearly independent** eigenvectors.

For each of its eigenvalues λ_i and corresponding eigenvectors \mathbf{v}_i , we have:

$$A\mathbf{v}_i = \lambda_i\mathbf{v}_i$$

Eigendecomposition

To make it more precise, suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix with n eigenvalues and n **linearly independent** eigenvectors.

For each of its eigenvalues λ_i and corresponding eigenvectors \mathbf{v}_i , we have:

$$A\mathbf{v}_i = \lambda_i\mathbf{v}_i$$

Or, if we put all the eigenvectors together as columns of a matrix P and all the eigenvalues as entries of a diagonal matrix Λ :

$$P = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_n \end{bmatrix}, \quad \Lambda = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix}$$

then

$$AP = P\Lambda$$

Eigendecomposition

To make it more precise, suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix with n eigenvalues and n **linearly independent** eigenvectors.

For each of its eigenvalues λ_i and corresponding eigenvectors \mathbf{v}_i , we have:

$$A\mathbf{v}_i = \lambda_i\mathbf{v}_i$$

Or, if we put all the eigenvectors together as columns of a matrix P and all the eigenvalues as entries of a diagonal matrix Λ :

$$P = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_n \end{bmatrix}, \quad \Lambda = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix}$$

then

$$AP = P\Lambda$$

or equivalently:

$$A = P\Lambda P^{-1}$$

Eigendecomposition

To make it more precise, suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix with n eigenvalues and n **linearly independent** eigenvectors.

For each of its eigenvalues λ_i and corresponding eigenvectors \mathbf{v}_i , we have:

$$A\mathbf{v}_i = \lambda_i\mathbf{v}_i$$

Or, if we put all the eigenvectors together as columns of a matrix P and all the eigenvalues as entries of a diagonal matrix Λ :

$$P = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_n \end{bmatrix}, \quad \Lambda = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix}$$

then


$$AP = P\Lambda$$

or equivalently:

$$A = P\Lambda P^{-1}$$

This is called the *eigendecomposition* (or *spectral decomposition*) of A .


Eigendecomposition



Geometrically, the eigendecomposition means that multiplying by A is:

- 1 rotating the basis to the eigenvector basis (*multiplying by P^{-1}*)


Eigendecomposition



Geometrically, the eigendecomposition means that multiplying by A is:

- 1 rotating the basis to the eigenvector basis (*multiplying by P^{-1}*)
- 2 stretching in each direction of \mathbf{v}_i by λ_i times (*multiplying by Λ*)


Eigendecomposition



Geometrically, the eigendecomposition means that multiplying by A is:

- 1 rotating the basis to the eigenvector basis (*multiplying by P^{-1}*)
- 2 stretching in each direction of \mathbf{v}_i by λ_i times (*multiplying by Λ*)
- 3 rotating the basis back to the standard basis (*multiplying by P*)


Eigendecomposition



Geometrically, the eigendecomposition means that multiplying by A is:

- 1 rotating the basis to the eigenvector basis (*multiplying by P^{-1}*)
- 2 stretching in each direction of \mathbf{v}_i by λ_i times (*multiplying by Λ*)
- 3 rotating the basis back to the standard basis (*multiplying by P*)

Eigendecomposition




Geometrically, the eigendecomposition means that multiplying by A is:

- ① rotating the basis to the eigenvector basis (*multiplying by P^{-1}*)
- ② stretching in each direction of \mathbf{v}_i by λ_i times (*multiplying by Λ*)
- ③ rotating the basis back to the standard basis (*multiplying by P*)

Computationally, the eigendecomposition is useful because multiplying by a diagonal matrix is very easy.

Eigendecomposition



Geometrically, the eigendecomposition means that multiplying by A is:

- 1 rotating the basis to the eigenvector basis (*multiplying by P^{-1}*)
- 2 stretching in each direction of \mathbf{v}_i by λ_i times (*multiplying by Λ*)
- 3 rotating the basis back to the standard basis (*multiplying by P*)

Computationally, the eigendecomposition is useful because multiplying by a diagonal matrix is very easy.

Question

How would you compute A^{10} ?

Singular Value Decomposition (SVD)

The eigendecomposition is very useful, but it only works for square symmetric matrices with enough linearly independent eigenvectors. What if we want to decompose a non-square matrix?

Singular Value Decomposition (SVD)

The eigendecomposition is very useful, but it only works for square symmetric matrices with enough linearly independent eigenvectors. What if we want to decompose a non-square matrix?

The answer is the **Singular Value Decomposition (SVD)**:

Theorem

Any matrix $A \in \mathbb{R}^{m \times n}$ can be written as a product of three matrices:

$$A = U \Sigma V^T$$

where

- $U \in \mathbb{R}^{m \times m}$ is a rotation in \mathbb{R}^m , and $U^T U = I$
- $V \in \mathbb{R}^{n \times n}$ is a rotation in \mathbb{R}^n , and $V^T V = I$
- $\Sigma \in \mathbb{R}^{m \times n}$ is a diagonal matrix (with non-negative numbers)

Singular Value Decomposition (SVD)

The eigendecomposition is very useful, but it only works for square symmetric matrices with enough linearly independent eigenvectors. What if we want to decompose a non-square matrix?

The answer is the **Singular Value Decomposition (SVD)**:

Theorem

Any matrix $A \in \mathbb{R}^{m \times n}$ can be written as a product of three matrices:


$$A = U \Sigma V^T$$

where

- $U \in \mathbb{R}^{m \times m}$ is a rotation in \mathbb{R}^m , and $U^T U = I$
- $V \in \mathbb{R}^{n \times n}$ is a rotation in \mathbb{R}^n , and $V^T V = I$
- $\Sigma \in \mathbb{R}^{m \times n}$ is a diagonal matrix (with non-negative numbers)

The numbers in Σ are called the *singular values* of A .


Singular Value Decomposition (SVD)

$$A = U \Sigma V^T$$


The diagram shows the matrix equation $A = U \Sigma V^T$ with dimension labels. Matrix A is represented by a blue square with a vertical dimension of m and a horizontal dimension of n . Matrix U is a blue square with a vertical dimension of m and a horizontal dimension of m . Matrix Σ is a blue rectangle with a vertical dimension of m and a horizontal dimension of n . Matrix V^T is a blue square with a vertical dimension of n and a horizontal dimension of n . The dimensions are indicated by arrows and labels above and to the left of each matrix.

When A is symmetric, the singular values coincide with the eigenvalues of A , and this decomposition (essentially) coincides with the eigendecomposition.

Singular Value Decomposition (SVD)

$$A = U \Sigma V^T$$


The diagram shows the matrix equation $A = U \Sigma V^T$ with dimension labels. Matrix A is represented by a blue square with a vertical double-headed arrow on the left labeled m and a horizontal double-headed arrow on top labeled n . To its right is an equals sign. Matrix U is a blue square with a vertical double-headed arrow on the left labeled m and a horizontal double-headed arrow on top labeled m . To its right is matrix Σ , a blue rectangle with a vertical double-headed arrow on the left labeled m and a horizontal double-headed arrow on top labeled n . To the right of Σ is matrix V^T , a blue square with a vertical double-headed arrow on the left labeled n and a horizontal double-headed arrow on top labeled n .

When A is symmetric, the singular values coincide with the eigenvalues of A , and this decomposition (essentially) coincides with the eigendecomposition.

We won't go into the details of how to compute the SVD, but we will see some of its applications later in the course.

Positive and Negative Definiteness

One last thing about eigenvalues. Turns out,

Theorem

A symmetric matrix $A \in \mathbb{R}^{n \times n}$ always has n eigenvalues (not necessarily distinct).

Positive and Negative Definiteness

One last thing about eigenvalues. Turns out,

Theorem

A symmetric matrix $A \in \mathbb{R}^{n \times n}$ always has n eigenvalues (not necessarily distinct).

If all of its eigenvalues are positive, we say that the matrix is **positive definite** and denote:

$$A \succ 0$$

Positive and Negative Definiteness

One last thing about eigenvalues. Turns out,

Theorem

A symmetric matrix $A \in \mathbb{R}^{n \times n}$ always has n eigenvalues (not necessarily distinct).

If all of its eigenvalues are positive, we say that the matrix is **positive definite** and denote:

$$A \succ 0$$

This is actually equivalent to saying that for any vector $\mathbf{x} \neq \mathbf{0}$:

$$\mathbf{x}^T A \mathbf{x} > 0$$

Positive and Negative Definiteness

One last thing about eigenvalues. Turns out,

Theorem

A symmetric matrix $A \in \mathbb{R}^{n \times n}$ always has n eigenvalues (not necessarily distinct).

If all of its eigenvalues are positive, we say that the matrix is **positive definite** and denote:

$$A \succ 0$$

This is actually equivalent to saying that for any vector $\mathbf{x} \neq \mathbf{0}$:

$$\mathbf{x}^T A \mathbf{x} > 0$$

Similarly, if all eigenvalues are negative (or equivalently, if $\mathbf{x}^T A \mathbf{x} < 0$ for all $\mathbf{x} \neq \mathbf{0}$), we say that the matrix is **negative definite**:

$$A \prec 0$$

Positive and Negative Definiteness

Example

$A = \begin{bmatrix} 4 & -4 \\ -4 & 5 \end{bmatrix}$ is positive definite because its eigenvalues are

$$\lambda_1 \approx 0.47, \quad \lambda_2 \approx 8.53$$

Positive and Negative Definiteness

Example

$A = \begin{bmatrix} 4 & -4 \\ -4 & 5 \end{bmatrix}$ is positive definite because its eigenvalues are

$$\lambda_1 \approx 0.47, \quad \lambda_2 \approx 8.53$$

or equivalently, for any $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \neq \mathbf{0}$:

$$\mathbf{x}^T A \mathbf{x} = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 4 & -4 \\ -4 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 4x_1^2 - 8x_1x_2 + 5x_2^2 > 0$$

Positive and Negative Definiteness

Example

$A = \begin{bmatrix} 4 & -4 \\ -4 & 5 \end{bmatrix}$ is positive definite because its eigenvalues are

$$\lambda_1 \approx 0.47, \quad \lambda_2 \approx 8.53$$

or equivalently, for any $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \neq \mathbf{0}$:

$$\mathbf{x}^T A \mathbf{x} = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 4 & -4 \\ -4 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 4x_1^2 - 8x_1x_2 + 5x_2^2 > 0$$

Can we check if a matrix is positive/negative definite without computing its eigenvalues?

Positive and Negative Definiteness

Suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nn} \end{bmatrix}$$

Positive and Negative Definiteness

Suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nn} \end{bmatrix}$$

and look at its upper-left submatrices:

$$A_1 = [a_{11}], \quad A_2 = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad A_3 = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}, \quad \dots$$

Positive and Negative Definiteness

Suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nn} \end{bmatrix}$$

and look at its upper-left submatrices:

$$A_1 = [a_{11}], \quad A_2 = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad A_3 = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}, \quad \dots$$

$A \succ 0$ if and only if the determinants of all these submatrices are positive:

$$\det(A_1) > 0, \quad \det(A_2) > 0, \quad \det(A_3) > 0, \quad \dots$$

Positive and Negative Definiteness

Suppose $A \in \mathbb{R}^{n \times n}$ is a **symmetric** matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nn} \end{bmatrix}$$

and look at its upper-left submatrices:

$$A_1 = [a_{11}], \quad A_2 = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad A_3 = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}, \quad \dots$$

$A \succ 0$ if and only if the determinants of all these submatrices are positive:

$$\det(A_1) > 0, \quad \det(A_2) > 0, \quad \det(A_3) > 0, \quad \dots$$

and $A \prec 0$ if and only if they alternate in sign:

$$\det(A_1) < 0, \quad \det(A_2) > 0, \quad \det(A_3) < 0, \quad \dots$$