

Taylor Series, Integral

Hayk Aprikyan, Hayk Tarkhanyan

Convex and Concave Functions

We know that the sign of the derivative tells us whether a function is increasing or decreasing:

$f' > 0 \Rightarrow$ increasing,

$f' < 0 \Rightarrow$ decreasing


Convex and Concave Functions

We know that the sign of the derivative tells us whether a function is increasing or decreasing:

$$f' > 0 \Rightarrow \text{increasing,}$$

$$f' < 0 \Rightarrow \text{decreasing}$$

However, a function can increase like this




Convex and Concave Functions

We know that the sign of the derivative tells us whether a function is increasing or decreasing:


$$f' > 0 \Rightarrow \text{increasing,}$$

$$f' < 0 \Rightarrow \text{decreasing}$$

However, a function can increase like this



or like this




Convex and Concave Functions

We know that the sign of the derivative tells us whether a function is increasing or decreasing:


$$f' > 0 \Rightarrow \text{increasing,}$$

$$f' < 0 \Rightarrow \text{decreasing}$$

However, a function can increase like this





or like this



How can you determine which way it is?

Convex and Concave Functions



Definition

We say that $f(x)$ is **convex** on some interval if for any two points on its graph, the line connecting them always lies **above** the graph of $f(x)$.

Definition

We say that $f(x)$ is **concave** on some interval if for any two points on its graph, the line connecting them always lies **below** the graph of $f(x)$.

Convex and Concave Functions

More technically, a function is

- convex if

$$f(\alpha p + (1 - \alpha)q) \leq \alpha f(p) + (1 - \alpha)f(q)$$

- concave if

$$f(\alpha p + (1 - \alpha)q) \geq \alpha f(p) + (1 - \alpha)f(q)$$

for any two points p and q , and for any number $0 \leq \alpha \leq 1$.

Convex and Concave Functions

More technically, a function is

- convex if

$$f(\alpha p + (1 - \alpha)q) \leq \alpha f(p) + (1 - \alpha)f(q)$$

- concave if

$$f(\alpha p + (1 - \alpha)q) \geq \alpha f(p) + (1 - \alpha)f(q)$$

for any two points p and q , and for any number $0 \leq \alpha \leq 1$.

We will not prove this

Convex and Concave Functions

More technically, a function is

- convex if

$$f(\alpha p + (1 - \alpha)q) \leq \alpha f(p) + (1 - \alpha)f(q)$$

- concave if

$$f(\alpha p + (1 - \alpha)q) \geq \alpha f(p) + (1 - \alpha)f(q)$$

for any two points p and q , and for any number $0 \leq \alpha \leq 1$.

We will not prove this but open the link and play with α to get a feeling of what this means:

- www.desmos.com/calculator/ujoh0mh59d

Convex and Concave Functions

So, how to know if a function is convex or concave?

Convex and Concave Functions

So, how to know if a function is convex or concave?

Again, derivative is all you need!

Theorem

If $f(x)$ is twice-differentiable (i.e. there exists $f''(x)$), then:

- f is convex if and only if $f''(x) \geq 0$
- f is concave if and only if $f''(x) \leq 0$

Convex and Concave Functions

So, how to know if a function is convex or concave?

Again, derivative is all you need!

Theorem


If $f(x)$ is twice-differentiable (i.e. there exists $f''(x)$), then:

- f is convex if and only if $f''(x) \geq 0$
- f is concave if and only if $f''(x) \leq 0$

Examples

- 1 $f(x) = x^2$ is convex on $(-\infty, \infty)$.
 $f''(x) = 2 > 0$
- 2 $f(x) = -x^2$ is concave on $(-\infty, \infty)$.
 $f''(x) = -2 < 0$
- 3 $f(x) = x$ is both convex and concave on $(-\infty, \infty)$.
 $f''(x) = 0$

Convex and Concave Functions



Taylor Series

As we saw, the derivatives of a function tell so much about them. Can they be used to approximate the function?

Taylor Series

As we saw, the derivatives of a function tell so much about them. Can they be used to approximate the function?

Suppose we have a function $f(x)$. Let us construct another function $g(x)$ which, at some point a has:

As we saw, the derivatives of a function tell so much about them. Can they be used to approximate the function?

Suppose we have a function $f(x)$. Let us construct another function $g(x)$ which, at some point a has:

- the same value as f

Taylor Series

As we saw, the derivatives of a function tell so much about them. Can they be used to approximate the function?

Suppose we have a function $f(x)$. Let us construct another function $g(x)$ which, at some point a has:

- the same value as f
- the same rate of change as f

As we saw, the derivatives of a function tell so much about them. Can they be used to approximate the function?

Suppose we have a function $f(x)$. Let us construct another function $g(x)$ which, at some point a has:

- the same value as f
- the same rate of change as f
- the same rate of the rate of change as f

As we saw, the derivatives of a function tell so much about them. Can they be used to approximate the function?

Suppose we have a function $f(x)$. Let us construct another function $g(x)$ which, at some point a has:

- the same value as f
- the same rate of change as f
- the same rate of the rate of change as f
- ...

As we saw, the derivatives of a function tell too much about them. Can they be used to approximate the function?

Suppose we have a function $f(x)$. Let us construct another function $g(x)$ which, at some point a has (or, equivalently):

- $g(a) = f(a)$
- $g'(a) = f'(a)$
- $g''(a) = f''(a)$
- ...

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Question

Which simple function has the value 1 at $x = 0$, and derivative 1 at $x = 0$?

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Question

Which simple function has the value 1 at $x = 0$, and derivative 1 at $x = 0$?

The polynomial $g(x) = 1 + x$ has just that properties. Moreover, it is easy to work with, since it is a polynomial!

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Question

Which simple function has the value 1 at $x = 0$, and derivative 1 at $x = 0$?

The polynomial $g(x) = 1 + x$ has just that properties. Moreover, it is easy to work with, since it is a polynomial!

Definition

If there exist derivatives $f', f'', \dots, f^{(k)}$, then the polynomial

$$P_k(x) = f(a)$$

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Question

Which simple function has the value 1 at $x = 0$, and derivative 1 at $x = 0$?

The polynomial $g(x) = 1 + x$ has just that properties. Moreover, it is easy to work with, since it is a polynomial!

Definition

If there exist derivatives f' , f'' , \dots , $f^{(k)}$, then the polynomial

$$P_k(x) = f(a) + f'(a)(x - a)$$

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Question

Which simple function has the value 1 at $x = 0$, and derivative 1 at $x = 0$?

The polynomial $g(x) = 1 + x$ has just that properties. Moreover, it is easy to work with, since it is a polynomial!

Definition

If there exist derivatives $f', f'', \dots, f^{(k)}$, then the polynomial

$$P_k(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2$$

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Question

Which simple function has the value 1 at $x = 0$, and derivative 1 at $x = 0$?

The polynomial $g(x) = 1 + x$ has just that properties. Moreover, it is easy to work with, since it is a polynomial!

Definition

If there exist derivatives $f', f'', \dots, f^{(k)}$, then the polynomial

$$P_k(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \dots + \frac{f^{(k)}(a)}{k!}(x - a)^k$$

is called the **Taylor polynomial** of order k of the function $f(x)$ about the point a .

Taylor Series

For example, suppose $f(x) = e^x$. Then $f(0) = 1$ and $f'(0) = 1$.

Question

Which simple function has the value 1 at $x = 0$, and derivative 1 at $x = 0$?

The polynomial $g(x) = 1 + x$ has just that properties. Moreover, it is easy to work with, since it is a polynomial!

Definition

If there exist derivatives $f', f'', \dots, f^{(k)}$, then the polynomial

$$P_k(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \dots + \frac{f^{(k)}(a)}{k!}(x - a)^k$$

is called the **Taylor polynomial** of order k of the function $f(x)$ about the point a .

If we take $k = \infty$, we will get an infinite sum which is called the **Taylor series** of $f(x)$ about a .

Taylor Series

Taylor's Theorem

If a function $f(x)$ is k times differentiable at the point a , then

$$P_k(x) \rightarrow f(x), \quad k \rightarrow \infty$$

around some (maybe small) interval around a .

Taylor Series

Taylor's Theorem

If a function $f(x)$ is k times differentiable at the point a , then

$$P_k(x) \rightarrow f(x), \quad k \rightarrow \infty$$

around some (maybe small) interval around a .

Example


- The Taylor series for any polynomial is the polynomial itself.

- $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + x^4 + \dots$

- $\ln(1+x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$


- $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$

Taylor Series




Taylor Series

1




Taylor Series

$$1 + x$$




Taylor Series

$$1 + x + \frac{x^2}{2}$$




Taylor Series

$$1 + x + \frac{x^2}{2} + \frac{x^3}{6}$$




Taylor Series

$$1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24}$$




Taylor Series

$$1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{5!}$$




Taylor Series

$$1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{5!} + \frac{x^6}{6!}$$



Taylor Series

$$1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{5!} + \frac{x^6}{6!} + \frac{x^7}{7!}$$



- Try another function yourself!

Indefinite Integral

$2x$ is the derivative of x^2

Indefinite Integral

$2x$ is the derivative of x^2

So what is x^2 ?

Indefinite Integral

$2x$ is the derivative of x^2

So what is x^2 ?

x^2 is the *antiderivative* of $2x$

Indefinite Integral

$2x$ is the derivative of x^2

So what is x^2 ?

x^2 is the *antiderivative* of $2x$

Definition

F is called an **antiderivative** or **indefinite integral** of f if $F'(x) = f(x)$.

Indefinite Integral

$2x$ is the derivative of x^2

So what is x^2 ?

x^2 is the *antiderivative* of $2x$

Definition

F is called an **antiderivative** or **indefinite integral** of f if $F'(x) = f(x)$.

The indefinite integral is denoted like this:

$$\int f(x) dx$$

Indefinite Integral

So in this notation,

$$\int 2x \, dx = x^2$$

Can you name another function that also has derivative $2x$?

Indefinite Integral

So in this notation,

$$\int 2x \, dx = x^2$$

Can you name another function that also has derivative $2x$?

- $(x^2 + 1)' = 2x$

Indefinite Integral

So in this notation,

$$\int 2x \, dx = x^2$$

Can you name another function that also has derivative $2x$?

- $(x^2 + 1)' = 2x$
- $(x^2 + 2)' = 2x$

Indefinite Integral

So in this notation,

$$\int 2x \, dx = x^2$$

Can you name another function that also has derivative $2x$?

- $(x^2 + 1)' = 2x$
- $(x^2 + 2)' = 2x$
- $(x^2 + c)' = 2x$

Indefinite Integral

So in this notation,

$$\int 2x \, dx = x^2$$

Can you name another function that also has derivative $2x$?

- $(x^2 + 1)' = 2x$
- $(x^2 + 2)' = 2x$
- $(x^2 + c)' = 2x$

If $F'(x) = f(x)$, then $(F(x) + c)' = f(x)$ for any number c , so actually:

$$\int 2x \, dx = x^2 + C$$

for any number C .

Indefinite Integral

So in this notation,

$$\int 2x \, dx = x^2$$

Can you name another function that also has derivative $2x$?

- $(x^2 + 1)' = 2x$
- $(x^2 + 2)' = 2x$
- $(x^2 + c)' = 2x$

If $F'(x) = f(x)$, then $(F(x) + c)' = f(x)$ for any number c , so actually:

$$\int 2x \, dx = x^2 + C$$

for any number C .

$\int f(x) \, dx$ is not one function, but a *set* of functions.

Indefinite Integral

Properties

1

$$\int af(x) dx = a \int f(x) dx$$

Indefinite Integral

Properties

1

$$\int af(x) dx = a \int f(x) dx$$

2

$$\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$$

Indefinite Integral

Properties

1

$$\int af(x) dx = a \int f(x) dx$$

2

$$\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$$

3

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx$$

Indefinite Integral

Properties

1

$$\int af(x) dx = a \int f(x) dx$$

2

$$\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$$

3

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx$$

Indefinite Integral

Properties

1

$$\int af(x) dx = a \int f(x) dx$$

2

$$\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$$

3

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx$$

This can also be written as:

$$\int f dg = fg - \int g df,$$

where $\int f dg = \int fg' dx$.

Indefinite Integral

- The antiderivative of any constant $f(x) = a$ is:

$$\int a \, dx = ax + C$$

Indefinite Integral

- The antiderivative of any constant $f(x) = a$ is:

$$\int a \, dx = ax + C$$

- The antiderivative of $f(x) = x$ is:

$$\int x \, dx = \frac{x^2}{2} + C$$

Indefinite Integral

- The antiderivative of any constant $f(x) = a$ is:

$$\int a \, dx = ax + C$$

- The antiderivative of $f(x) = x$ is:

$$\int x \, dx = \frac{x^2}{2} + C$$

- For any constant $n \neq -1$, the antiderivative of $f(x) = x^n$ is:

$$\int x^n \, dx = \frac{x^{n+1}}{n+1} + C$$

Indefinite Integral

- The antiderivative of any constant $f(x) = a$ is:

$$\int a \, dx = ax + C$$

- The antiderivative of $f(x) = x$ is:

$$\int x \, dx = \frac{x^2}{2} + C$$

- For any constant $n \neq 1$, the derivative of $f(x) = x^n$ is:

$$\int x^n \, dx = \frac{x^{n+1}}{n+1} + C$$

- The antiderivative of $f(x) = \frac{1}{x}$ is:

$$\int \frac{1}{x} \, dx = \ln |x| + C$$

Indefinite Integral

- The antiderivative of $f(x) = e^x$ is:

$$\int e^x dx = e^x + C$$

Indefinite Integral

- The antiderivative of $f(x) = e^x$ is:

$$\int e^x dx = e^x + C$$

- The antiderivative of $f(x) = \cos x$ is:

$$\int \cos x dx = \sin x + C$$

Indefinite Integral

- The antiderivative of $f(x) = e^x$ is:

$$\int e^x dx = e^x + C$$

- The antiderivative of $f(x) = \cos x$ is:

$$\int \cos x dx = \sin x + C$$

- The antiderivative of $f(x) = \sin x$ is:

$$\int \sin x dx = -\cos x + C$$

Indefinite Integral

- The antiderivative of $f(x) = e^x$ is:

$$\int e^x dx = e^x + C$$

- The antiderivative of $f(x) = \cos x$ is:

$$\int \cos x dx = \sin x + C$$

- The antiderivative of $f(x) = \sin x$ is:

$$\int \sin x dx = -\cos x + C$$

- The antiderivative of $f(x) = \frac{1}{1+x^2}$ is:

$$\int \frac{1}{1+x^2} dx = \operatorname{arctg} x + C$$

Example

Which function should you differentiate to get $x \cos x$?

Example

Which function should you differentiate to get $x \cos x$?

Notice that $\cos x = (\sin x)'$, so by the Property 3,

$$\int x \cos x \, dx = \int x(\sin x)' \, dx = \int x \, d(\sin x) = x(\sin x) - \int (\sin x) \, dx$$

Example

Which function should you differentiate to get $x \cos x$?

Notice that $\cos x = (\sin x)'$, so by the Property 3,

$$\int x \cos x \, dx = \int x(\sin x)' \, dx = \int x \, d(\sin x) = x(\sin x) - \int (\sin x) \, dx$$

and the last integral $= \cos x + C$ (as we know).

Example

Which function should you differentiate to get $x \cos x$?

Notice that $\cos x = (\sin x)'$, so by the Property 3,

$$\int x \cos x \, dx = \int x(\sin x)' \, dx = \int x \, d(\sin x) = x(\sin x) - \int (\sin x) \, dx$$

and the last integral $= \cos x + C$ (as we know).

All of this is, of course, computation and tricks (no need to memorize).

Example

Which function should you differentiate to get $x \cos x$?

Notice that $\cos x = (\sin x)'$, so by the Property 3,

$$\int x \cos x \, dx = \int x(\sin x)' \, dx = \int x \, d(\sin x) = x(\sin x) - \int (\sin x) \, dx$$


and the last integral $= \cos x + C$ (as we know).

All of this is, of course, computation and tricks (no need to memorize).

How about we make an *actual* use of this?


Definite Integral

Suppose we are given the velocity of a car at each timepoint. How can we calculate the distance travelled by the car?



Definite Integral

Suppose we are given the velocity of a car at each timepoint. How can we calculate the distance travelled by the car?



According to physics, distance = area under the velocity curve.

Definite Integral

Question




Suppose you have a continuous function f . How can we calculate the area under its graph?

Definite Integral

Question

Suppose you have a continuous function f . How can we calculate the area under its graph?

By dividing it into tiny rectangles and adding up their areas.



Definite Integral

Take the interval $[a, b]$ and divide (partition) it into n small parts with points $\{x_0, x_1, \dots, x_n\}$. Let $\Delta x_i = x_i - x_{i-1}$ denote the length of $[x_{i-1}, x_i]$.

Definite Integral

Take the interval $[a, b]$ and divide (partition) it into n small parts with points $\{x_0, x_1, \dots, x_n\}$. Let $\Delta x_i = x_i - x_{i-1}$ denote the length of $[x_{i-1}, x_i]$.

Definition

The **Riemann sum** of a function $f(x)$ is given by:

$$R_n = \sum_{i=1}^n f(c_i) \Delta x_i$$

where c_i is any point from $[x_{i-1}, x_i]$.

Definite Integral

Take the interval $[a, b]$ and divide (partition) it into n small parts with points $\{x_0, x_1, \dots, x_n\}$. Let $\Delta x_i = x_i - x_{i-1}$ denote the length of $[x_{i-1}, x_i]$.

Definition

The **Riemann sum** of a function $f(x)$ is given by:

$$R_n = \sum_{i=1}^n f(c_i) \Delta x_i$$

where c_i is any point from $[x_{i-1}, x_i]$.

Making the Δ s smaller, this sum will get closer to the area.

Definite Integral

Take the interval $[a, b]$ and divide (partition) it into n small parts with points $\{x_0, x_1, \dots, x_n\}$. Let $\Delta x_i = x_i - x_{i-1}$ denote the length of $[x_{i-1}, x_i]$.

Definition

The **Riemann sum** of a function $f(x)$ is given by:

$$R_n = \sum_{i=1}^n f(c_i) \Delta x_i$$

where c_i is any point from $[x_{i-1}, x_i]$.

Making the Δ s smaller, this sum will get closer to the area.


Definition

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x_i$$

is called the **definite integral** of the function $f(x)$ on $[a, b]$.

Definite Integral

The definite integral $\int_a^b f(x) dx$ represents the **signed area** between the graph of $f(x)$ and the x -axis over the interval $[a, b]$.



- Play with Riemann sums!

Definite Integral

How can we calculate the definite integral without limits?

Definite Integral

How can we calculate the definite integral without limits?

Theorem (very important)

Suppose that $f(x)$ is continuous on the interval $[a, b]$. If $F(x)$ is an antiderivative of $f(x)$, then

$$\int_a^b f(x) dx = F(b) - F(a)$$

Definite Integral

How can we calculate the definite integral without limits?

Theorem (very important)

Suppose that $f(x)$ is continuous on the interval $[a, b]$. If $F(x)$ is an antiderivative of $f(x)$, then

$$\int_a^b f(x) dx = F(b) - F(a)$$

Example

- $$\int_0^2 x^2 dx = \frac{1}{3} \cdot x^3 \Big|_0^2 = \frac{1}{3}(2^3 - 0^3) = \frac{8}{3}$$

- $$\int_0^\pi \sin x dx = -\cos x \Big|_0^\pi = -\cos \pi - (-\cos 0) = 2$$

Definite Integral

1

$$\int_a^b f(x) dx = - \int_b^a f(x) dx, \quad \int_a^a f(x) dx = 0$$

Definite Integral

1

$$\int_a^b f(x) dx = - \int_b^a f(x) dx, \quad \int_a^a f(x) dx = 0$$

2

$$\int_a^b cf(x) dx = c \int_a^b f(x) dx$$

Definite Integral

1

$$\int_a^b f(x) dx = - \int_b^a f(x) dx, \quad \int_a^a f(x) dx = 0$$

2

$$\int_a^b cf(x) dx = c \int_a^b f(x) dx$$

3

$$\int_a^b (f(x) \pm g(x)) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$$

Definite Integral

1

$$\int_a^b f(x) dx = - \int_b^a f(x) dx, \quad \int_a^a f(x) dx = 0$$

2

$$\int_a^b cf(x) dx = c \int_a^b f(x) dx$$

3

$$\int_a^b (f(x) \pm g(x)) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$$

4

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

Definite Integral

1

$$\int_a^b f(x) dx = - \int_b^a f(x) dx, \quad \int_a^a f(x) dx = 0$$

2

$$\int_a^b cf(x) dx = c \int_a^b f(x) dx$$

3

$$\int_a^b (f(x) \pm g(x)) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$$

4

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

5

$$\int_a^b f(x) dx = \int_a^b f(y) dy$$

i.e. the name of the variable does not matter.