

Random Variables

Hayk Aprikyan, Hayk Tarkhanyan

April 15, 2025

Random Variables

Recap:

During the previous lecture we had a problem like this:

Example

Both the bus and you get to the bus stop at random times between 12 pm and 1 pm. When the bus arrives, it waits for 5 minutes before leaving. When you arrive, you wait for 20 minutes before leaving if the bus doesn't come. What is the probability that you catch the bus?

Here we did not know the **exact values** of the arrival times, so in order to compare them, we **denoted them** by y and b – and calculated their probabilities.

In this case, we say that y and b are **random variables**.

Random Variables

Definition

A quantity whose value is unknown and depends on a random experiment is called a **random variable**.

Random Variables

Definition

A quantity whose value is unknown and depends on a random experiment is called a **random variable**.

Example

You throw a fair dice and win \$1 if it is prime, lose \$1 if it is composite (4 or 6), and stay even otherwise. In other words, if X is your profit, then

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

Random Variables

Definition

A quantity whose value is unknown and depends on a random experiment is called a **random variable**.

Example

You throw a fair dice and win \$1 if it is prime, lose \$1 if it is composite (4 or 6), and stay even otherwise. In other words, if X is your profit, then

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

Since the exact value you win/lose cannot be known, it is a random variable.

Random Variables

Definition

A quantity whose value is unknown and depends on a random experiment is called a **random variable**.

Example

You throw a fair dice and win \$1 if it is prime, lose \$1 if it is composite (4 or 6), and stay even otherwise. In other words, if X is your profit, then

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

Since the exact value you win/lose cannot be known, it is a random variable.

What is the probability that $X = 1$?

Random Variables

Definition

A quantity whose value is unknown and depends on a random experiment is called a **random variable**.

Example

You throw a fair dice and win \$1 if it is prime, lose \$1 if it is composite (4 or 6), and stay even otherwise. In other words, if X is your profit, then

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

Since the exact value you win/lose cannot be known, it is a random variable.

What is the probability that $X = 1$? (We'll get to this later.)

Random Variables

Example

Two fair dice are thrown. The sum of their numbers is a random variable.

Random Variables

Example

Two fair dice are thrown. The sum of their numbers is a random variable.

Example

Two teams play football.

- The number of goals they score is a random variable.
- The difference between their scores is also a random variable.

Random Variables

Example

Two fair dice are thrown. The sum of their numbers is a random variable.

Example

Two teams play football.

- The number of goals they score is a random variable.
- The difference between their scores is also a random variable.

Example

You call a random number and ask the number of children in their family.

Random Variables

Example

Two fair dice are thrown. The sum of their numbers is a random variable.

Example

Two teams play football.

- The number of goals they score is a random variable.
- The difference between their scores is also a random variable.

Example

You call a random number and ask the number of children in their family.

Example

You are waiting for the bus 62. The time until it arrives is a random variable.

Random Variables

Example

Two fair dice are thrown. The sum of their numbers is a random variable.

Example

Two teams play football.

- The number of goals they score is a random variable.
- The difference between their scores is also a random variable.

Example

You call a random number and ask the number of children in their family.

Example

You are waiting for the bus 62. The time until it arrives is a random variable.

What values can each of the random variables above take?

Random Variables (optional)

Not a random variable

Two fair dice are thrown,

- who threw the dice,
- the color of the sky,
- what will show up if you had thrown a third die

are not random variables, as their values *do not depend* on the outcome of the dice (even if you knew the outcome, you could not answer these).

Random Variables (optional)

Not a random variable

Two fair dice are thrown,

- who threw the dice,
- the color of the sky,
- what will show up if you had thrown a third die

are not random variables, as their values *do not depend* on the outcome of the dice (even if you knew the outcome, you could not answer these).

In practice, every unknown thing in the given problem is a random variable.

Technically speaking, although, the value of the random variable should somehow depend on the outcome ω of the experiment. In other words, if you ask "When is $X = 4$?", there should be a way to measure that probability, i.e. $\{X = 4\}$ should be an event.

In your textbook you will find it as $\{\omega \text{ for which } X = 4\} \in \mathcal{F}$, but we skip the technicalities.

Random Variables

Notice that in some cases the possible values look like $\{2, 3, 4, \dots, 12\}$ or $\{0, 1, 2, \dots\}$, while in others they are intervals like $(0, +\infty)$.

Random Variables

Notice that in some cases the possible values look like $\{2, 3, 4, \dots, 12\}$ or $\{0, 1, 2, \dots\}$, while in others they are intervals like $(0, +\infty)$.

Definition

If the values of X can be represented as a list (say, x_1, x_2, x_3, \dots with probabilities $0.2, 0.1, 0.4, \dots$), it is called a **discrete random variable**.

Random Variables

Notice that in some cases the possible values look like $\{2, 3, 4, \dots, 12\}$ or $\{0, 1, 2, \dots\}$, while in others they are intervals like $(0, +\infty)$.

Definition

If the values of X can be represented as a list (say, x_1, x_2, x_3, \dots with probabilities $0.2, 0.1, 0.4, \dots$), it is called a **discrete random variable**.

Otherwise,

Definition

If the values of X cannot be represented as a list (e.g. they are an interval), it is called a **continuous random variable**.

Of course, what we are interested in the most, is the probability that X takes a certain value.

For example:

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

and the question is:

$$\mathbb{P}(X = 1) = ?$$

Of course, what we are interested in the most, is the probability that X takes a certain value.

For example:

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

and the question is:

$$\mathbb{P}(X = 1) = ?$$

Since $X = 1$ corresponds to the outcomes $\{2, 3, 5\}$, the probability $\mathbb{P}(X = 1)$ is the same as $\mathbb{P}(\{2, 3, 5\})$

Of course, what we are interested in the most, is the probability that X takes a certain value.

For example:

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

and the question is:

$$\mathbb{P}(X = 1) = ?$$

Since $X = 1$ corresponds to the outcomes $\{2, 3, 5\}$, the probability $\mathbb{P}(X = 1)$ is the same as $\mathbb{P}(\{2, 3, 5\}) = \frac{3}{6} = 0.5$.

Of course, what we are interested in the most, is the probability that X takes a certain value.

For example:

$$X = \begin{cases} 1, & \text{if } \omega \in \{2, 3, 5\} \\ -1, & \text{if } \omega \in \{4, 6\} \\ 0, & \text{if } \omega = 1 \end{cases}$$

and the question is:

$$\mathbb{P}(X = 1) = ?$$

Since $X = 1$ corresponds to the outcomes $\{2, 3, 5\}$, the probability $\mathbb{P}(X = 1)$ is the same as $\mathbb{P}(\{2, 3, 5\}) = \frac{3}{6} = 0.5$.

The thing above, $\mathbb{P}(X = \text{something})$, is called the **probability mass function** or just the **PMF** of X .

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam.

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) =$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15}.$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}(X = 1) =$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}(X = 1) = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}(X = 1) = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}(X = 1) = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

$$\mathbb{P}(X = 2) =$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}(X = 1) = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

$$\mathbb{P}(X = 2) = \frac{4}{15} \cdot \frac{3}{14}$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}(X = 1) = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

$$\mathbb{P}(X = 2) = \frac{4}{15} \cdot \frac{3}{14} = \frac{6}{105} \quad (\text{or just } 1 - \frac{11}{21} - \frac{44}{105})$$

Example

A pitiful student knows 4 out of 15 topics, and there will be 2 questions on the exam. Let X = number of questions the student gets correct.

$$\mathbb{P}(X = 0) = \frac{11}{15} \cdot \frac{10}{14} = \frac{11}{21} \quad (\text{rip student})$$

$$\mathbb{P}(X = 1) = \frac{4}{15} \cdot \frac{11}{14} + \frac{11}{15} \cdot \frac{4}{14} = \frac{44}{105}$$

$$\mathbb{P}(X = 2) = \frac{4}{15} \cdot \frac{3}{14} = \frac{6}{105} \quad (\text{or just } 1 - \frac{11}{21} - \frac{44}{105})$$

k (right answers)	$\mathbb{P}(X = k)$
0	11/21
1	44/105
2	6/105

While interested in gaining knowledge and stuff, one thing the student cares most is if they pass or not, i.e.

- Not only the probabilities that $X = 0$ or $X = 1$ or $X = 2$,

While interested in gaining knowledge and stuff, one thing the student cares most is if they pass or not, i.e.

- Not only the probabilities that $X = 0$ or $X = 1$ or $X = 2$,
- But importantly, the probability that $X > 0$,

While interested in gaining knowledge and stuff, one thing the student cares most is if they pass or not, i.e.

- Not only the probabilities that $X = 0$ or $X = 1$ or $X = 2$,
- But importantly, the probability that $X > 0$,
(or the opposite, $X \leq 2$ or $X < 2$).

While interested in gaining knowledge and stuff, one thing the student cares most is if they pass or not, i.e.

- Not only the probabilities that $X = 0$ or $X = 1$ or $X = 2$,
- But importantly, the probability that $X > 0$,
(or the opposite, $X \leq 2$ or $X < 2$).

In our case, $\mathbb{P}(X < 2) = \mathbb{P}(X = 0) + \mathbb{P}(X = 1) = \frac{99}{105}$.

While interested in gaining knowledge and stuff, one thing the student cares most is if they pass or not, i.e.

- Not only the probabilities that $X = 0$ or $X = 1$ or $X = 2$,
- But importantly, the probability that $X > 0$,
(or the opposite, $X \leq 2$ or $X < 2$).

In our case, $\mathbb{P}(X < 2) = \mathbb{P}(X = 0) + \mathbb{P}(X = 1) = \frac{99}{105}$.

If there were, say, 5 questions and more than 3 points needed to pass, then we would be interested in

$$\mathbb{P}(X \leq 3) \quad (\text{probability of failing})$$

or

$$\mathbb{P}(X > 3) \quad (\text{probability of passing})$$

The quantity

$$\mathbb{P}(X \leq k) \quad (\text{where } k \text{ is any number})$$

is called the **cumulative distribution function** or the **CDF** of X .

The quantity

$$\mathbb{P}(X \leq k) \quad (\text{where } k \text{ is any number})$$

is called the **cumulative distribution function** or the **CDF** of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}(X \leq k)$$

The quantity

$$\mathbb{P}(X \leq k) \quad (\text{where } k \text{ is any number})$$

is called the **cumulative distribution function** or the **CDF** of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}(X \leq k)$$

Properties

- $0 \leq F_X(x) \leq 1$, for any $x \in \mathbb{R}$

The quantity

$$\mathbb{P}(X \leq k) \quad (\text{where } k \text{ is any number})$$

is called the **cumulative distribution function** or the **CDF** of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}(X \leq k)$$

Properties

- $0 \leq F_X(x) \leq 1$, for any $x \in \mathbb{R}$
- $F_X(x)$ is a non-decreasing function

The quantity

$$\mathbb{P}(X \leq k) \quad (\text{where } k \text{ is any number})$$

is called the **cumulative distribution function** or the **CDF** of X .

It is conveniently denoted by $F_X(k)$ or $F(k)$:

$$F_X(k) = \mathbb{P}(X \leq k)$$

Properties

- $0 \leq F_X(x) \leq 1$, for any $x \in \mathbb{R}$
- $F_X(x)$ is a non-decreasing function
- $\lim_{x \rightarrow -\infty} F_X(x) = 0$ and $\lim_{x \rightarrow +\infty} F_X(x) = 1$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) =$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) =$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

$$F_X(5.9) =$$

Example

Let X indicate the number on a fair die. Then

$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

$$F_X(5.9) = \frac{5}{6}$$

Example

Let X indicate the number on a fair die. Then


$$F_X(6.1) = 1$$

$$F_X(6) = 1$$

$$F_X(5.9) = \frac{5}{6}$$

$$F_X(x) = \begin{cases} 0, & \text{if } x < 1, \\ \frac{1}{6}, & \text{if } 1 \leq x < 2, \\ \frac{2}{6}, & \text{if } 2 \leq x < 3, \\ \frac{3}{6}, & \text{if } 3 \leq x < 4, \\ \frac{4}{6}, & \text{if } 4 \leq x < 5, \\ \frac{5}{6}, & \text{if } 5 \leq x < 6, \\ 1, & \text{if } x \geq 6 \end{cases}$$

Graphs of CDFs usually look like this:



Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

- While the students may be asked different questions,
- \Rightarrow it may happen that X and Y are not equal,
- still Y has the same PMF and CDF as X (check it!)

In this case, we say that X and Y are identically distributed:

Definition

Two random variables are said to be **identically distributed** if their PMFs/CDFs are equal.

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

- While the students may be asked different questions,
- \Rightarrow it may happen that X and Y are not equal,
- still Y has the same PMF and CDF as X (check it!)

In this case, we say that X and Y are identically distributed:

Definition

Two random variables are said to be **identically distributed** if their PMFs/CDFs are equal.

Yet, identically distributed \neq equal. E.g. if you toss a fair coin, and set

$$X = \begin{cases} 0, & \omega = \mathbf{H}, \\ 1, & \omega = \mathbf{T}, \end{cases} \quad Y = \begin{cases} 1, & \omega = \mathbf{H}, \\ 0, & \omega = \mathbf{T}, \end{cases}$$

They both have the save PMFs:

$$\mathbb{P}(X = 0) = \mathbb{P}(X = 1) = \mathbb{P}(Y = 0) = \mathbb{P}(Y = 1) = \frac{1}{2}$$

Suppose we have a second pitiful student who has also studied for 4 topics out of 15. Let Y = number of questions the second student gets correct.

- While the students may be asked different questions,
- \Rightarrow it may happen that X and Y are not equal,
- still Y has the same PMF and CDF as X (check it!)

In this case, we say that X and Y are identically distributed:

Definition

Two random variables are said to be **identically distributed** if their PMFs/CDFs are equal.

Yet, identically distributed \neq equal. E.g. if you toss a fair coin, and set

$$X = \begin{cases} 0, & \omega = \mathbf{H}, \\ 1, & \omega = \mathbf{T}, \end{cases} \quad Y = \begin{cases} 1, & \omega = \mathbf{H}, \\ 0, & \omega = \mathbf{T}, \end{cases}$$

They both have the same PMFs:

$$\mathbb{P}(X = 0) = \mathbb{P}(X = 1) = \mathbb{P}(Y = 0) = \mathbb{P}(Y = 1) = \frac{1}{2} \quad \text{but } X \neq Y$$

Finally, what if X is a continuous random variable?




Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}(X = 0.5)$?

Finally, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}(X = 0.5)$? What about $\mathbb{P}(X = 0.589)$?

Finally, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}(X = 0.5)$? What about $\mathbb{P}(X = 0.589)$? As previously, it is 0.

Finally, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}(X = 0.5)$? What about $\mathbb{P}(X = 0.589)$? As previously, it is 0.

What can we say about the PMF of X ?

Finally, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}(X = 0.5)$? What about $\mathbb{P}(X = 0.589)$? As previously, it is 0.

What can we say about the PMF of X ? It is always zero!

Finally, what if X is a continuous random variable?



Say you randomly put your finger on some number X in $(0, 10)$. What is the probability that $\mathbb{P}(X = 0.5)$? What about $\mathbb{P}(X = 0.589)$? As previously, it is 0.

What can we say about the PMF of X ? It is always zero!

So another question we might ask is:

Question

What do you think $\mathbb{P}(X \leq 5)$ is?

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}(X \leq 5) = \frac{1}{2}$$

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}(X \leq 5) = \frac{1}{2}$$

Fortunately, we have another tool in case of continuous random variables.

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}(X \leq 5) = \frac{1}{2}$$

Fortunately, we have another tool in case of continuous random variables.

Definition

If X is a continuous random variable, then there exists a nonnegative function $f(x)$ such that for any $c \in \mathbb{R}$,

$$F_X(c) = \mathbb{P}(X \leq c) = \int_{-\infty}^c f(t) dt$$

The function f is called the **probability density function** or **PDF** of X .

While for discrete random variables both PMF and CDF make sense, for continuous random variables PMF is useless, but still can rely on CDF:

$$\mathbb{P}(X \leq 5) = \frac{1}{2}$$

Fortunately, we have another tool in case of continuous random variables.

Definition

If X is a continuous random variable, then there exists a nonnegative function $f(x)$ such that for any $c \in \mathbb{R}$,

$$F_X(c) = \mathbb{P}(X \leq c) = \int_{-\infty}^c f(t) dt$$

The function f is called the **probability density function** or **PDF** of X .


The PDF of a continuous random variable plays essentially the same role as the PMF of a discrete random variable.

Just like the density of an object measures the concentration of mass (per unit volume), the probability density function captures the density of *probability* at point x :

$$f_X(x) = \lim_{h \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + h)}{h}$$

Just like the density of an object measures the concentration of mass (per unit volume), the probability density function captures the density of *probability* at point x :

$$f_X(x) = \lim_{h \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + h)}{h}$$



Properties

① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Properties

① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

② $\int_{-\infty}^{\infty} f_X(x) dx = 1$

Properties

- ① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$
- ② $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- ③ If F_X is continuous at x , $F'_X(x) = f_X(x)$

Properties

- ① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$
- ② $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- ③ If F_X is continuous at x , $F'_X(x) = f_X(x)$
- ④ $\mathbb{P}(a < X \leq b) = F_X(b) - F_X(a) = \int_a^b f_X(x) dx$

Properties

- ① $f_X(x) \geq 0$ for all $x \in \mathbb{R}$
- ② $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- ③ If F_X is continuous at x , $F'_X(x) = f_X(x)$
- ④ $\mathbb{P}(a < X \leq b) = F_X(b) - F_X(a) = \int_a^b f_X(x) dx$
- ⑤ $\mathbb{P}(X = a) = \mathbb{P}(X \leq a) - \mathbb{P}(X < a)$

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

For $x < a$, $F_X(x) = \mathbb{P}(X \leq x) = 0$,

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.

Let's find the CDF and PDF of X .

For $x < a$, $F_X(x) = \mathbb{P}(X \leq x) = 0$,

For $x \geq b$, $F_X(x) = \mathbb{P}(X \leq x) = 1$,

Example

Ani chooses a random real number X uniformly from the interval $[a, b]$.

By "uniformly" we mean that for any two intervals of the same length (e.g. $(1.3, 1.5)$ and $(4.7, 4.9)$) X can belong to them with the same probability.


Let's find the CDF and PDF of X .


For $x < a$, $F_X(x) = \mathbb{P}(X \leq x) = 0$,

For $x \geq b$, $F_X(x) = \mathbb{P}(X \leq x) = 1$,

For $a \leq x \leq b$, we have:

$$F_X(x) = \mathbb{P}(X \leq x) = \mathbb{P}(X \in [a, x]) = \frac{x - a}{b - a}$$





Thus,


$$F_X(x) = \begin{cases} 0 & \text{if } x < a \\ \frac{x-a}{b-a} & \text{if } a \leq x \leq b \\ 1 & \text{if } x > b \end{cases}$$

To find $f_X(x)$, we take the derivative of $F_X(x)$:

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a < x < b \\ 0 & \text{if } x < a \text{ or } x > b \end{cases}$$

To find $f_X(x)$, we take the derivative of $F_X(x)$:

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a < x < b \\ 0 & \text{if } x < a \text{ or } x > b \end{cases}$$



Independence

Recall that two events A and B are called independent if they do not affect each other, i.e.

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

Independence

Recall that two events A and B are called independent if they do not affect each other, i.e.

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

Similarly, we will say that two random variables X and Y are independent if their values do not affect each other (i.e. if you know the value of X , it gives you no information about the value of Y).

Independence

Recall that two events A and B are called independent if they do not affect each other, i.e.

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

Similarly, we will say that two random variables X and Y are independent if their values do not affect each other (i.e. if you know the value of X , it gives you no information about the value of Y).

Definition

X and Y are called **independent** if

$$\mathbb{P}(X \leq a \text{ and } Y \leq b) = \mathbb{P}(X \leq a) \cdot \mathbb{P}(Y \leq b)$$

for any $a, b \in \mathbb{R}$.

So the probability of both X and Y simultaneously being less than some numbers is just their *separate* probabilities multiplied together.