## In The Name of God. The Merciful, The Compassionate.

## Symmetric Matrices, Positive Definiteness

notes on Gilbert Strang videos, Lecture 25

## 1 Symmetric Matrices: $A = A^T$

1. The eigenvalues are real:

$$Ax = \lambda x \Rightarrow \bar{A}\bar{x} = \bar{\lambda}\bar{x} = A\bar{x} = \bar{\lambda}\bar{x}$$

$$\Rightarrow \bar{x}^T A^T = \bar{x}^T \bar{\lambda} = \bar{x}^T A$$

$$\Rightarrow \bar{x}^T A x = \bar{\lambda}\bar{x}^T x$$

$$Ax = \lambda x \Rightarrow \bar{x}^T A x = \lambda \bar{x}^T x$$

$$\Rightarrow \lambda \bar{x}^T x = \bar{\lambda}\bar{x}^T x$$

$$\Rightarrow \lambda = \bar{\lambda} \Rightarrow \lambda \text{ is real}$$

- Note that  $\bar{x}^T x$  is  $\geq 0$ .
- If A is complex: the proof is still working if  $\bar{A}^T = A$
- $\bullet$  Good matrices are conjugate transpose.
- 2. The eigenvectors are (can be chosen) orthogonal  $A=S\Lambda S^{-1}\to Q\Lambda Q^T \text{ Assuming orthonormal eigenvectors}$

$$A = Q\Lambda Q^{T} = \begin{bmatrix} | & | \\ q_{1} & \dots & q_{n} \\ | & | \end{bmatrix} \begin{bmatrix} \lambda_{1} & 0 \\ \vdots & \ddots & \vdots \\ 0 & \lambda_{n} \end{bmatrix} \begin{bmatrix} q_{1}^{T} \\ \vdots \\ q_{n}^{T} \end{bmatrix}$$
$$= \lambda_{1}q_{1}q_{1}^{T} + \lambda_{2}q_{2}q_{2}^{T} + \dots$$