

NUMERICAL OPTIMIZATION

TUTO 1: GRADIENTS AND MINIMIZATION

L. DESBAT & F. IUTZELER

A. DIFFERENTIABILITY, MINIMA, AND CONVEXITY

Exercise 1 (Quadratic functions).

- a. In \mathbb{R}^n , compute the gradient of the squared Euclidean norm $\|\cdot\|_2^2$ at a generic point $x \in \mathbb{R}^n$.
- b. Let A be an $m \times n$ real matrix and b a size- m real vector. We define $f(x) = \|Ax - b\|_2^2$. For a generic vector $a \in \mathbb{R}^n$, compute the gradient $\nabla f(a)$ and Hessian $H_f(a)$.
- c. Let C be an $n \times n$ real matrix, d a size- n real vector, and $e \in \mathbb{R}$. We define $g(x) = x^T C x + d^T x + e$. For a generic vector $a \in \mathbb{R}^n$, compute the gradient $\nabla g(a)$ and Hessian $H_g(a)$.
- d. Can all functions of the form of f and be written in the form of g ? And conversely?

Exercise 2 (Basic Differential calculus). Use the composition lemma to compute the gradients of:

- a. $f_1(x) = \|Ax - b\|_2^2$.
- b. $f_2(x) = \|x\|_2$.


Exercise 3 (Preparing the Lab). In the first lab, we will consider the following toy functions:

$$\begin{aligned}
 f : \quad \mathbb{R}^2 &\rightarrow \mathbb{R} \\
 (x_1, x_2) &\mapsto 4(x_1 - 3)^2 + 2(x_2 - 1)^2 \\
 g : \quad \mathbb{R}^2 &\rightarrow \mathbb{R} \\
 (x_1, x_2) &\mapsto \log(1 + \exp(4(x_1 - 3)^2) + \exp(2(x_2 - 1)^2)) - \log(3) \\
 r : \quad \mathbb{R}^2 &\rightarrow \mathbb{R} \\
 (x_1, x_2) &\mapsto (1 - x_1)^2 + 100(x_2 - x_1^2)^2 \\
 t : \quad \mathbb{R}^2 &\rightarrow \mathbb{R} \\
 (x_1, x_2) &\mapsto (0.6x_1 + 0.2x_2)^2 ((0.6x_1 + 0.2x_2)^2 - 4(0.6x_1 + 0.2x_2) + 4) + (-0.2x_1 + 0.6x_2)^2 \\
 p : \quad \mathbb{R}^2 &\rightarrow \mathbb{R} \\
 (x_1, x_2) &\mapsto |x_1 - 3| + 2|x_2 - 1|.
 \end{aligned}$$


- a. From the 3D plots of A.1, which functions are visibly non-convex.
- b. For all five functions, show that they are convex or give an argument for their non-convexity.
- c. For functions f, g, r, t , compute their gradient.
- d. For functions f, g , compute their Hessian.

Exercise 4 (Fundamentals of convexity).


- a. Let f and g be two convex functions. Show that $m(x) = \max(f(x), g(x))$ is convex.
- b. Show that $f_1(x) = \max(x^2 - 1, 0)$ is convex.
- c. Let f be a convex function and g be a convex, non-decreasing function. Show that $c(x) = g(f(x))$ is convex.
- d. Show that $f_2(x) = \exp(x^2)$ is convex. What about $f_3(x) = \exp(-x^2)$?
- e. Justify why the 1-norm, the 2 norm, and the squared 2-norm are convex.




(A) a *simple* function: f




(B) some *harder* function: g



(C) *Rosenbrock's* function: r



(D) *two pits* function: t



(E) *polyhedral* function: p

FIGURE A.1. 3D plots of the considered functions

Exercise 5 (Strict and strong convexity). A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is said

- *strictly convex* if for any $x \neq y \in \mathbb{R}^n$ and any $\alpha \in]0, 1[$

$$f(\alpha x + (1 - \alpha)y) < \alpha f(x) + (1 - \alpha)f(y)$$

- *strongly convex* if there exists $\beta > 0$ such that $f - \frac{\beta}{2} \|\cdot\|_2^2$ is convex.

- a. For a strictly convex function f , show that the problem

$$\begin{cases} \min f(x) \\ x \in C \end{cases}$$

where C is a convex set admits at most one solution.

- b. Show that a strongly convex function is also strictly convex.

(hint: use the identity $\|\alpha x + (1 - \alpha)y\|^2 = \alpha\|x\|^2 + (1 - \alpha)\|y\|^2 - \alpha(1 - \alpha)\|x - y\|^2$.)

Exercise 6 (Optimality conditions). Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice differentiable function and $\bar{x} \in \mathbb{R}^n$. We suppose that f admits a local minimum at \bar{x} that is $f(x) \geq f(\bar{x})$ for all x in a neighborhood¹ of \bar{x} .

- For any direction $u \in \mathbb{R}^n$, we define the $\mathbb{R} \rightarrow \mathbb{R}$ function $q(t) = f(\bar{x} + tu)$. Compute $q'(t)$.
- By using the first order Taylor expansion of q at 0, show that $\nabla f(\bar{x}) = 0$.
- Compute $q''(t)$. By using the second order Taylor expansion of q at 0, show that $\nabla^2 f(\bar{x})$ is positive semi-definite.

B. THE GRADIENT ALGORITHM

Exercise 7 (Descent lemma). A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be L -smooth if it is differentiable and its gradient ∇f is L -Lipchitz continuous, that is

$$\forall x, y \in \mathbb{R}^n, \quad \|\nabla f(x) - \nabla f(y)\| \leq L\|x - y\|.$$

The goal of the exercise is to prove that if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is L -smooth, then for all $x, y \in \mathbb{R}^n$,

$$f(x) \leq f(y) + (x - y)^T \nabla f(y) + \frac{L}{2} \|x - y\|^2$$

- a. Starting from fundamental theorem of calculus stating that for all $x, y \in \mathbb{R}^n$,

$$f(x) - f(y) = \int_0^1 (x - y)^T \nabla f(y + t(x - y)) dt$$

prove the descent lemma.

- b. Give a function for which the inequality is tight and one for which it is not.

Exercise 8 (Smooth functions). Consider the constant stepsize gradient algorithm $x_{k+1} = x_k - \gamma \nabla f(x_k)$ on an L -smooth function f with some minimizer (i.e. some x^* such that $f(x) \geq f(x^*)$ for all x).

- Use the *descent lemma* to prove convergence of the sequence $(f(x_k))_k$ when $\gamma \leq 2/L$.
- Did you use at some point that the function was convex? Conclude about the convergence of the gradient algorithm on smooth non-convex functions.

¹Formally, one would write $\forall x \in \mathbb{R}^n$ such that $\|x - \bar{x}\| \leq \varepsilon$ for $\varepsilon > 0$ and some norm $\|\cdot\|$.